Online ISSN: 2249-4626
Print ISSN: 0975-5896

Global Journal

of Science Frontier Research: D

Agriculture and Veterinary



F1 Friesian X Boran

Effective Micro Organism

Highlights

Aquatic Biodiversity Analysis

Performance of Broiler Breeders

Discovering Thoughts, Inventing Future

VOLUME 14 ISSUE 9 VERSION 1.0

© 2001-2014 by Global Journal of Science Frontier Research, USA

Volume 14 Issue 40 was originally issued as D9

Global Journal of Science Frontier Research: D
Agriculture & Veterinary

Volume 14 Issue 40
Originally issued as D9

Global Journal of Science Frontier Research: D
Agriculture & Veterinary

Volume 14 Issue 9 (Ver. 1.0)

Open Association of Research Society

© Global Journal of Science Frontier Research. 2014.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Science Frontier Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Science Frontier Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Science Frontier Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website http://globaljournals.us/terms-and-condition/menu-id-1463/

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089
License No.: 42125/022010/1186
Registration No.: 430374
Import-Export Code: 1109007027
Employer Identification Number (EIN):
USA Tax ID: 98-0673427

Global Journals Inc.

*(A Delaware USA Incorporation with "Good Standing"; **Reg. Number: 0423089**)*

Sponsors: Open Association of Research Society
Open Scientific Standards

Publisher's Headquarters office

Global Journals Headquarters
301st Edgewater Place Suite, 100 Edgewater Dr.-Pl,
Wakefield MASSACHUSETTS, Pin: 01880,
United States of America
USA Toll Free: +001-888-839-7392
USA Toll Free Fax: +001-888-839-7392

Offset Typesetting

Global Journals Incorporated
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,
Pin: CR9 2ER, United Kingdom

Packaging & Continental Dispatching

Global Journals
E-3130 Sudama Nagar, Near Gopur Square,
Indore, M.P., Pin:452009, India

Find a correspondence nodal officer near you

To find nodal officer of your country, please email us at *local@globaljournals.org*

eContacts

Press Inquiries: *press@globaljournals.org*
Investor Inquiries: *investors@globaljournals.org*
Technical Support: *technology@globaljournals.org*
Media & Releases: *media@globaljournals.org*

Pricing (Including by Air Parcel Charges):

For Authors:
22 USD (B/W) & 50 USD (Color)
Yearly Subscription (Personal & Institutional):
200 USD (B/W) & 250 USD (Color)

Integrated Editorial Board
(Computer Science, Engineering, Medical, Management, Natural Science, Social Science)

John A. Hamilton,"Drew" Jr.,
Ph.D., Professor, Management
Computer Science and Software Engineering
Director, Information Assurance Laboratory
Auburn University

Dr. Henry Hexmoor
IEEE senior member since 2004
Ph.D. Computer Science, University at Buffalo
Department of Computer Science
Southern Illinois University at Carbondale

Dr. Osman Balci, Professor
Department of Computer Science
Virginia Tech, Virginia University
Ph.D.and M.S.Syracuse University, Syracuse, New York
M.S. and B.S. Bogazici University, Istanbul, Turkey

Yogita Bajpai
M.Sc. (Computer Science), FICCT
U.S.A.Email:
yogita@computerresearch.org

Dr. T. David A. Forbes
Associate Professor and Range Nutritionist
Ph.D. Edinburgh University - Animal Nutrition
M.S. Aberdeen University - Animal Nutrition
B.A. University of Dublin- Zoology

Dr. Wenying Feng
Professor, Department of Computing & Information Systems
Department of Mathematics
Trent University, Peterborough,
ON Canada K9J 7B8

Dr. Thomas Wischgoll
Computer Science and Engineering,
Wright State University, Dayton, Ohio
B.S., M.S., Ph.D.
(University of Kaiserslautern)

Dr. Abdurrahman Arslanyilmaz
Computer Science & Information Systems Department
Youngstown State University
Ph.D., Texas A&M University
University of Missouri, Columbia
Gazi University, Turkey

Dr. Xiaohong He
Professor of International Business
University of Quinnipiac
BS, Jilin Institute of Technology; MA, MS, PhD,. (University of Texas-Dallas)

Burcin Becerik-Gerber
University of Southern California
Ph.D. in Civil Engineering
DDes from Harvard University
M.S. from University of California, Berkeley & Istanbul University

Dr. Bart Lambrecht
Director of Research in Accounting and FinanceProfessor of Finance
Lancaster University Management School
BA (Antwerp); MPhil, MA, PhD (Cambridge)

Dr. Carlos García Pont
Associate Professor of Marketing
IESE Business School, University of Navarra
Doctor of Philosophy (Management), Massachusetts Institute of Technology (MIT)
Master in Business Administration, IESE, University of Navarra
Degree in Industrial Engineering, Universitat Politècnica de Catalunya

Dr. Fotini Labropulu
Mathematics - Luther College
University of ReginaPh.D., M.Sc. in Mathematics
B.A. (Honors) in Mathematics
University of Windso

Dr. Lynn Lim
Reader in Business and Marketing
Roehampton University, London
BCom, PGDip, MBA (Distinction), PhD, FHEA

Dr. Mihaly Mezei
ASSOCIATE PROFESSOR
Department of Structural and Chemical Biology, Mount Sinai School of Medical Center
Ph.D., Etvs Lornd University
Postdoctoral Training,
New York University

Dr. Söhnke M. Bartram
Department of Accounting and FinanceLancaster University Management SchoolPh.D. (WHU Koblenz)
MBA/BBA (University of Saarbrücken)

Dr. Miguel Angel Ariño
Professor of Decision Sciences
IESE Business School
Barcelona, Spain (Universidad de Navarra)
CEIBS (China Europe International Business School).
Beijing, Shanghai and Shenzhen
Ph.D. in Mathematics
University of Barcelona
BA in Mathematics (Licenciatura)
University of Barcelona

Philip G. Moscoso
Technology and Operations Management
IESE Business School, University of Navarra
Ph.D in Industrial Engineering and Management, ETH Zurich
M.Sc. in Chemical Engineering, ETH Zurich

Dr. Sanjay Dixit, M.D.
Director, EP Laboratories, Philadelphia VA Medical Center
Cardiovascular Medicine - Cardiac Arrhythmia
Univ of Penn School of Medicine

Dr. Han-Xiang Deng
MD., Ph.D
Associate Professor and Research
Department Division of Neuromuscular Medicine
Davee Department of Neurology and Clinical NeuroscienceNorthwestern University
Feinberg School of Medicine

Dr. Pina C. Sanelli
Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic Radiology
M.D., State University of New York at Buffalo,School of Medicine and Biomedical Sciences

Dr. Roberto Sanchez
Associate Professor
Department of Structural and Chemical Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University

Dr. Wen-Yih Sun
Professor of Earth and Atmospheric SciencesPurdue University Director
National Center for Typhoon and Flooding Research, Taiwan
University Chair Professor
Department of Atmospheric Sciences, National Central University, Chung-Li, TaiwanUniversity Chair Professor
Institute of Environmental Engineering, National Chiao Tung University, Hsin-chu, Taiwan.Ph.D., MS The University of Chicago, Geophysical Sciences
BS National Taiwan University, Atmospheric Sciences
Associate Professor of Radiology

Dr. Michael R. Rudnick
M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and Hypertension Division (PMC)
Penn Medicine, University of Pennsylvania
Presbyterian Medical Center, Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of Internal Medicine

Dr. Bassey Benjamin Esu
B.Sc. Marketing; MBA Marketing; Ph.D Marketing
Lecturer, Department of Marketing, University of Calabar
Tourism Consultant, Cross River State Tourism Development Department
Co-ordinator , Sustainable Tourism Initiative, Calabar, Nigeria

Dr. Aziz M. Barbar, Ph.D.
IEEE Senior Member
Chairperson, Department of Computer Science
AUST - American University of Science & Technology
Alfred Naccash Avenue – Ashrafieh

President Editor (HON.)

Dr. George Perry, (Neuroscientist)
Dean and Professor, College of Sciences
Denham Harman Research Award (American Aging Association)
ISI Highly Cited Researcher, Iberoamerican Molecular Biology Organization
AAAS Fellow, Correspondent Member of Spanish Royal Academy of Sciences
University of Texas at San Antonio
Postdoctoral Fellow (Department of Cell Biology)
Baylor College of Medicine
Houston, Texas, United States

Chief Author (HON.)

Dr. R.K. Dixit
M.Sc., Ph.D., FICCT
Chief Author, India
Email: authorind@computerresearch.org

Dean & Editor-in-Chief (HON.)

Vivek Dubey(HON.)
MS (Industrial Engineering),
MS (Mechanical Engineering)
University of Wisconsin, FICCT
Editor-in-Chief, USA
editorusa@computerresearch.org

Sangita Dixit
M.Sc., FICCT
Dean & Chancellor (Asia Pacific)
deanind@computerresearch.org

Suyash Dixit
(B.E., Computer Science Engineering), FICCTT
President, Web Administration and
Development , CEO at IOSRD
COO at GAOR & OSS

Er. Suyog Dixit
(M. Tech), BE (HONS. in CSE), FICCT
SAP Certified Consultant
CEO at IOSRD, GAOR & OSS
Technical Dean, Global Journals Inc. (US)
Website: www.suyogdixit.com
Email:suyog@suyogdixit.com

Pritesh Rajvaidya
(MS) Computer Science Department
California State University
BE (Computer Science), FICCT
Technical Dean, USA
Email: pritesh@computerresearch.org

Luis Galárraga
J!Research Project Leader
Saarbrücken, Germany

Contents of the Issue

i. Copyright Notice
ii. Editorial Board Members
iii. Chief Author and Dean
iv. Contents of the Issue
v. Research and Review Papers

1. Growth Performance of F1 Friesian X Boran Crossbred Dairy Calves Supplemented with Effective Micro Organism (EM) Fermented Wheat Bran (Bokashi) in the Central Highlands of Ethiopia. *1-6*
2. Role of Camel's Milk in Growth Rate of Chroococcus Sp. *7-9*
3. Aquatic Biodiversity Analysis of Four Rice Terraces Clusters in the Cordillera Region, Northern Philippines. *11-16*
4. The Effects of Feeding Time and Regimen on the Performance of Broiler Breeders: A Review. *17-19*

vi. Fellows and Auxiliary Memberships
vii. Process of Submission of Research Paper
viii. Preferred Author Guidelines
ix. Index

Global Journal of Science Frontier Research: D
Agriculture and Veterinary
Volume 14 Issue 9 Version 1.0 Year 2014
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Growth Performance of F1 Friesian X Boran Crossbred Dairy Calves Supplemented with Effective Micro Organism (EM) Fermented Wheat Bran (Bokashi) in the Central Highlands of Ethiopia

By Aemiro Kehaliew, Getu kitaw, Getnet Assefa, Dereje Fekadu & Zewdie Wondatir

Ethiopian Institute of Agricultural Research, Ethiopia

Abstract- This experiment was designed to investigate the biological and economical response of feeding effective microorganisms treated wheat bran provided at different level to young crossbred calves. Twenty crossbred Frisian*Boran calves weighing (100± 10.85) kg and 6 months of age were selected and divided randomly in to 4 similar treatment groups (5 animals each group) in 100 days feeding trial. All the calves in the different groups were fed similar basal diet of native pasture hay. The first group (T1, control) fed native hay as a basal diet and concentrate mixture as a supplement. Group 2, 3 and 4 (T2, T3 and T4) fed the same basal diet and concentrate mixture replaced by 33, 66 and 100 % EM treated wheat bran, respectively.

Keywords: calves, efficiency, effective microorganisms, weight gain.

GJSFR-D Classification : FOR Code: 830302



Strictly as per the compliance and regulations of :



© 2014. Aemiro Kehaliew, Getu kitaw, Getnet Assefa, Dereje Fekadu & Zewdie Wondatir. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Growth Performance of F1 Friesian X Boran Crossbred Dairy Calves Supplemented with Effective Micro Organism (EM) Fermented Wheat Bran (Bokashi) in the Central Highlands of Ethiopia

Aemiro Kehaliew [α], Getu kitaw [σ], Getnet Assefa [ρ], Dereje Fekadu [ω] & Zewdie Wondatir [¥]

Abstract- This experiment was designed to investigate the biological and economical response of feeding effective microorganisms treated wheat bran provided at different level to young crossbred calves. Twenty crossbred Frisian*Boran calves weighing (100± 10.85) kg and 6 months of age were selected and divided randomly in to 4 similar treatment groups (5 animals each group) in 100 days feeding trial. All the calves in the different groups were fed similar basal diet of native pasture hay. The first group (T1, control) fed native hay as a basal diet and concentrate mixture as a supplement. Group 2, 3 and 4 (T2, T3 and T4) fed the same basal diet and concentrate mixture replaced by 33, 66 and 100 % EM treated wheat bran, respectively. Based on these treatments, the response of growing calves in terms of dry matter and nutrient intake, daily weight gain, feed conversion efficiency and economic feasibility were evaluated. The CP content of effective microorganisms (EM) treated wheat bran was lower than that of the concentrate mixture and wheat bran. However there was no significant difference in organic matter digestibility between the EM treated wheat bran and the concentrate mixture (65.26 Vs 65.61%), respectively. There was also no significant (P>0.05) difference in live weight change, dry matter and metabolizable energy intake among the different treatments. The highest daily body weight gain and better feed conversion efficiency was observed in T2. Among the different treatments the lowest feed cost per daily intake was encountered in T2. In general EM treated feed supplementation had positive effect on growth performance of calves.

Keywords: calves, efficiency, effective microorganisms, weight gain.

I. Introduction

Successful strategies for increasing the efficiency of utilization of low quality roughages include supplementation, chemical treatment, biological treatment and manipulation of the rumen ecosystem. Among the biological treatment methods, use of effective microorganism is one of the options for improving feed quality. Effective Microorganisms (EM) are a mixed culture of aerobic and anaerobic types of microbes living symbiotically with each other. These microorganisms are beneficial types, natural, free-living and safe. They can withstand a wide variation of environmental conditions making them consistent and stable. Modern animal production requires use of safe and effective additives to stimulate feed consumption and destroy harmful microorganisms of the diet in addition to be used as rumen manipulators to increase animal productivity.

According to Saili et al., 2010, fermentation of feeds with microorganisms had resulted in increased live weight gain. Probiotics are live microorganisms that often result in increased feed conversion efficiency and live weight gain (Sissons, 1989). They also introduce beneficial microorganisms into the gut which act to maintain optimal conditions within the gastrointestinal tract and inhibit the growth of pathogenic or other undesirable bacteria. Several functions of probiotics have been proposed, including the protection of young animals against enteropathic disorders such as diarrhea by inhibiting the colonization of the gut by coliform bacteria and an increase in feed conversion efficiency and live weight gain in growing animals (Fuller, 1990 and Saili et al., 2010).

Studies have shown that large mixtures of microorganisms can be used with superior results to single bacteria strain or simple mixtures of few types of bacteria (Mead and Impey, 1986; Stavric et al., 1991). The same authors also reported that the success of complex mixtures appears to be due to interactive effects of anaerobes and facultative anaerobes. As a mixed culture of beneficial and naturally occurring microorganisms, Effective Microorganisms (EM) contain selected species of microorganisms including predominant populations of lactic acid bacteria and yeasts and smaller numbers of photosynthetic bacteria, actinomycetes and other types of organisms. The use of EM in animal husbandry nowadays is very well identified in many parts of the world. In a study conducted in Belarus by Konoplya and Higa (2000), EM was successfully used in poultry and swine units as feed

Author α σ ω ¥: Holetta Agricultural Research Center.
Author ρ: Ethiopian Institute of Agricultural Research.
e-mail: zewbt2006@yahoo.com

© 2014 Global Journals Inc. (US)

constituent and sanitation spray. In South Africa EM was used to increase productivity in integrated animal units and poultry farms (Hanekon et al., 2001). In a sudy conducted by Maqbool etal.,2001, the use of EM treated rice straw proved to be beneficial for the prevention of disease probably by virtue of its ability to check the growth of mycotoxin producing fungi. This agent not only controls the disease but also increases the growth rate, feed consumption and feed efficacy. Similar results were also recorded in broilers by Khan et al. (1992) and Haddadin et al. (1996).

Thus, the objectives of the present study was to evaluate the effect of the supplemental value of different levels of EM treated wheat bran on feed intake, growth performance and economic benefit of F1 crossbred Friesian* Boran calves .

II. Material and Methods

a) Location and treatment description

This study was carried out at Holetta Agricultural Research Center (9^{O}N latitude; 38^{O}E longitude). The experimental site lies at an altitude of 2400 meters above seas level. In this study, twenty Fresian* Boran crossbred calves with an average live weight of 100 ±10.85 kg and six month of age were used. The experimental calves were divided randomly in to four treatment groups (five animals in each group). Calves of the first group (TI, control) fed native hay as a basal diet and concentrate mixture of 43% noug seed cake (Guzotia abyssinica), 55% wheat bran and 2% salt. Calves of the 2^{nd}, 3^{rd} and 4^{th} treatment group received the same basal diet with 33, 66 and 100% of the concentrate mixture replaced by effective microorganism (EM) treated wheat bran, respectively.

b) Experimental treatments

1. *Treatment 1:* native hay adlibtium + concentrate mixture(control)
2. *Treatment 2:* native hay adlibtium + 33% of the Concentrate mixture replaced by EM treated wheat bran
3. *Treatment 3:* native hay adlibtium + 66% of the Concentrate mixture replaced by EM treated wheat bran
4. *Treatment 4:* native hay adlibtium + 100% of the Concentrate mixture replaced by EM treated wheat bran

c) Activated solution and EM treatment preparation procedures

Bokashi (40% moisture) is an EM treated wheat bran that was prepared from; 1 lt EM1(stock solution) + lt molasses + 50 g table sugar + 18 lt chlorine free water + 50 kg wheat bran. The mix was allowed to ferment for about three weeks in a large plastic barrel of capacity 150kg before it was being fed to the animal. The EM stock solution was provided by Woljjejji private limited company. Bokashi (EM treated wheat bran) was self prepared based on the procedures as follows:

1. One liter of molasses and 50 gm of sugar was added in to 18 liters of water and it was stirred very well.
2. One liter of EM-1(stock solution) was added in to the mixture
3. Stirring continue until a homogeneous solution was attained
4. The container was covered with the plastic sheet and it was very well tied to prevent the entry of air.
5. It was stored in shaded place with ambient temperature for 20 days.
6. After 20 days the resulting mixture known as EM Activated Solution (EMAS) was ready for use.
7. EMAS was viable for 15 to 20 days after fermentation. Always keep in anaerobic condition.

d) Treatment of wheat bran with Activated solution

- Molasses was dissolved in the 20 liter of water.
- EMAS was added into the above prepared molasses solution.
- The diluted EM solution was poured into 50kg of wheat bran and mixed well. EM diluted solution was added gradually and mixed well while checking the moisture content. The moisture content should be about 30-40%.
- The mixture was put into a plastic container that did not permit air entry to maintain an anaerobic condition
- The plastic container was stored in a place away from direct sunlight
- It was allowed to ferment for 21 days.
- The Bokashi was ready for use when it gives a sweet fermented smell. If it produces a rotten smell, it is a failure.
- The Bokashi should be used soon after preparation. If storage is required, spread it on a concrete floor, dry well in the shade and then put into vinyl bag.

e) Feeding management

Twenty calves were assigned in to four experimental treatments in RCBD design. Each calf was kept in a separate pen in a well ventilated barn. Native hay and water was provided ad libitum. Concentrate mixture including the EM treated wheat bran was provided twice a day early in the morning and late in the afternoon. The concentrate mixture (43% noug seed cake, 55% wheat bran and 2% salt) was formulated to meet the daily nutrient requirement of calves according to Kearl, (1982). The feed offered and refusal was recorded daily. Data on live weight change was recorded fortnightly.

© 2014 Global Journals Inc. (US)

f) Analysis of feed chemical composition and digestibility

Samples of native hay, wheat bran, noug seed cake, EM treated wheat bran and concentrate mixture were analyzed for dry matter (DM), crude protein (CP) and total ash using procedures described by AOAC, (1990). The neutral detergent fiber (NDF) and acid detergent fiber (ADF) concentrations in feed samples were determined according to the procedure of Van Soest and Robertson (1985). In-vitro organic matter digestibility (IVOMD) was determined according to the procedure of Tilley and Terry, (1963)

g) Data analysis

The calves were randomly assigned to the different dietary treatments in Randomized Complete Block (RCBD) design. The data was analyzed using SAS statistical software (SAS, 2002) as per the following model.

$Y_{ij} = \mu + B_i + T_j + e_{ijk}$ .Where, Y_{ij} = Response variables, μ = Overall mean, Bi = The effect of i^{th} block, T_j = The effect of j^{th} treatment and e_{ijk} = Random error.

III. Results and Discussion

a) Chemical composition of experimental feeds

The chemical compositions of experimental feeds are shown in Table 1. Dry matter content of experimental feeds was in the range of 89.39 to 91.69%. Among the experimental feeds, Bokashi (EM treated wheat bran) had lower dry matter content. The CP content of EM treated wheat bran was lower than that of the concentrate mixture and wheat bran. In a similar study by Bruchem, (1998) CP and ME content of EM treated silage was slightly lower than that of untreated silage. This might be due to the fact that part of the nutrients is utilized by the microbes for their growth and multiplication. However there was no difference in organic matter digestibility between the EM treated wheat bran and the concentrate mixture (65.26 Vs 65.61%), respectively. EM treated wheat bran had higher NDF content than un treated wheat bran and concentrate mixture. The current result is also in agreement with the report of Bruchem, (1998). On the other hand higher ADF content encountered in concentrate mixture than EM treated and un treated wheat bran.

Table 1 : Chemical compositions of experimental feed ingredients

Feed ingredient	Chemical compositions (% DM)						
	DM	Ash	CP	NDF	ADF	Lignin	DOMD
Noug seed cake	91.69	8.93	37.23	38.63	25.52	4.71	56.19
Wheat bran	88.97	4.77	16.27	43.48	12.66	3.35	70.89
Bokashi	88.39	6.75	15.39	46.65	13.66	3.53	65.26
Concentrate	89.26	8.26	24.79	36.09	21.93	4.30	65.61
Natural pasture hay	90.65	8.58	6.10	74.40	43.83	7.24	46.02

b) Dry matter and nutrient intake

As shown in Table 2, total dry matter (DM) intakes of calves were in the range of 3.68 to 4.43 kg/day. In a similar study by Khin et al., (2008), the DM intake was in the range of 3.75 to 3.92 kg/day. The highest and the lowest DM intake was observed in treatment 1and 4, respectively. There was no significant ($P>0.05$) difference in dry matter intake among the different treatments. Among the EM treated feed supplemented groups 100% replacement had lower DM intake than the other groups.

There was significant ($P<0.05$) difference in crude protein intake (CPI) among the different supplements. The highest CP intake was obtained in concentrate mixture supplemented group. Among the EM treated feed supplemented groups 100% supplemented ones had the lowest CPI. The difference in CPI was due to replacement of the protein sources of the concentrate mixture at different level by EM treated wheat bran. Metabolizable energy intake (MEI) was not significantly ($P>0.05$) varied among the different treatments.

© 2014 Global Journals Inc. (US)

Table 2 : Voluntary feed dry matter and nutrient intake Feed DM Intake

Feed DM Intake	Treatment				Mean	SE
	1	2	3	4		
Concentrate (kg/d)	2.08	1.27	0.596	0	1.32	
Bokashi (kg/d)	0	0.717	1.30	1.88	1.30	
Hay (kg/d)	2.34[a]	2.07[a]	2.05[a]	1.80[a]	2.07	0.217
TDMI (kg/d)	4.43	4.06	3.95	3.68	4.03	0.449
Hay intake, %BWt.	1.58	1.52	1.63	1.46	1.55	
TDMI, %BWt.	2.99	2.98	3.15	2.99	3.03	
CPI (gm/d)	657[a]	550[ab]	471[ab]	396[b]	519	0.06
CPI(gm/kg MBW)	15.47	13.79	12.75	10.72	13.25	
MEI (MJ/d)	38.39	35.65	34.73	32.82	35.40	4.01
MEI(MJ/kg MBW)	0.90	0.89	0.93	0.89	0.90	

[a-b] ,means in the same row with different superscripts are significantly different each other (P<0.05), kg/d: kilogram per day, BWt: body weight, TDMI: total dry matter intake, CPI: crude protein intake, MEI: Metabolizable energy intake, MJ/d: mega joule per day, gm/d: gram per day, SE: Standard error, gm/kgMBW: gram per kilo gram metabolic body weight, MJ/Kg MBW: maga joule per kilo gram metabolic body weight

c) Live weight change

There was no significant (P>0.05) difference in live weigh change among the different treatments (Table 3). Daily live weight gain was in the range of 696.7 to 823.3 gm/day. The highest daily body weight gain was obtained in T2, where 33% of the concentrate supplement was replaced by EM treated wheat bran (Bokashi). A similar work reported by Maqbool et al., (2001) indicated that 39.2 percent of body weight increase (600 gm/day) in live weigh gain was encountered in calves fed EM treated straw compared to other forms of feeding. Previous finding by Bruchem (1998) also indicated that higher growth rate was obtained from EM treated feed. The finding of this study indicated that in spite of replacing the protein source in the concentrate mixture by 100% EM treated wheat bran, daily weight gain was not affected markedly compared to concentrate mix supplemented groups. This implies that EM treatment improved the quality of the treated feed particularly CP content of the ration as was also recorded by Maqbool et al., (1997). In another study, Khin et al., (2008) obtained a weight gain of 300 gm/calf/day with the supplementation of 6 cc probiotics. These findings showed that supplementation of probiotics had positive effect on growth performance of young calves. EM treatment in addition to increasing growth rate, feed consumption and feed efficacy, it also helps to control disease (Maqbool et al..,(2001). Similar results were also reported in broilers by Khan et al., (1992) and Haddadin et al., (1996). According to Saili et al., (2010) treatment of cocoa-pods with micro organisms (A. niger) also result in increased live weight gain.

Feed conversion efficiency (FCE) was not significantly (P>0.05) different among the different treatments as shown in Table 3. Relatively higher FCE was observed in T2 (5.70 g feed/g gain) and T4 (5.73 g feed/g gain) where 33 and 100% of the concentrate mixture replaced by EM treated wheat bran, respectively. A similar study by Khin, et al., (2008) showed a feed conversion efficiency of 13.08 kg feed /kg gain which was lower than the present study.

The use of EM in animal production have demonstrated beneficial effects not only on health and production but also on environmental pollution. According to Li et. al., 1994 the naturally occurring microorganisms contained in EM after they enter into the body create more effective intestinal microflora with a greater synthetic capability i.e. one that can synthesize vitamins, hormones and enzyme systems that improve digestion, enhance growth, provide disease resistance, suppress malodor, inhibit pathogen and improves product quality. In all cases the improvement were associated to a greater flow of microbial protein and aminoacid to the duodenum, increased number of rumem cellulytic bacteria which improves fiber degradation, changes in the volatile fatty acids contents, greater glucose availability and reduced mobilization of fatty acids based on glucose, insulin, non-esterified fatty acids and ketones concentration in the blood.

© 2014 Global Journals Inc. (US)

Table 3 : Live weight changes

Growth paramete	Treatment				Mean	SE
	1	2	3	4		
Initial weight (kg)	112.40	101.7	91.1	95.9	100.28	10.85
Final weight (kg)	185.6	175.8	161.8	158.6	170.45	19.30
1	2	3	4			64.33
112.40	101.7	91.1	95.9	100.28	10.85	0.72
185.6	175.8	161.8	158.6	170.45	19.30	
813.33	823.33	785.56	696.67	779.72	64.33	

DWG = daily weight gain, FCE = feed conversion efficiency, SE = Standard error

d) Daily feed cost of experimental calves

The price in feed cost among the different treatment groups was varied from 6.31 to 7.42 birr/day (Table 4). The lowest feed cost was encountered in treatment 4 where 100% of the concentrate mixture was replaced by EM treated wheat bran. On the other hand the cost of feed was higher in concentrate mixture supplemented group (T1). In all cases, the cost of EM activated solution treated wheat bran was lower than that of concentrate mixture supplementation. This finding also indicated that in areas where the cost of protein sources is expensive the use of EM treatment is the best option.

Table 4 : Daily feed cost of experimental calves

Feed ingredient	T1		T2		T3		T4	
	DMI (kg)	Cost (birr)	DMI (kg)	Cost (birr)	DMI (kg)	Cost (birr)	DMI (kg)	Cost (birr)
Hay	2.34	3.11	2.07	2.75	2.05	2.73	1.80	2.39
Wheat bran	1.144	2.20	0.6985	1.34	0.3278	0.63	0	0
Noug cake	0.8944	1.99	0.5461	1.22	0.2563	0.57	0	0
Bokashi	0	0	0.717	1.44*/1.59**	1.30	2.63*/2.89**	1.88	3.80*/4.17**
Salt	0.0416	0.12	0.0254	0.08	0.0119	0.04	0.0416	0.12
Total		7.42		6.83*/6.98**		6.60*/6.86**		6.31*/6.68**

* The cost for 1kg Bokashi prepared from stock solution activated to 20lt EM-2= 2.02 Eth. birr,

**The cost for 1kg Bokashi prepared from purchased EM-2 solution = 2.22 Eth. birr

Note that water and labor cost were not considered in calculating the daily feed cost

*During the study period the average exchange rate was 17.0 Ethiopian birr = 1 USD

IV. Conclusion

The highest daily body weight gain was obtained in T2 (823 gm/calf/day) where 33% of the concentrate supplement was replaced by EM treated wheat bran. Among the different treatments the lowest feed cost per daily intake was encountered in 100% of the EM treated wheat bran supplemented group. EM treated feed supplementation at different level had positive effect on growth performance of calves. Further study on improvement of poor quality feed resource particularly crop residue with EM treatment is required.

References Références Referencias

1. AOAC (Association of Official Analytical Chemists) (1990). Official methods of analysis (15th edition), Washington, DC.
2. Bruchem JV (1998). Effects of beneficial micro-organisms.
3. Fuller R (1990). Probiotics in agricultures. Ag Biotech News and Info.2(2) : 217.
4. Haddadin MSY, Rahim SMA, Hashlamour ER, Robinson RK (1996). The effect of Lactobacillus on the production and chemical composition of hen's egg. Poult. Sci 75: 491-94.
5. Hanekon D, Prinsloo JF, Schoonbee HJ (2001). A comparison of the effect of anolyte and EM on the faecal bacteria loads in the water and on fish produced in pig com fish integrated production units. In: Y.D.A Senanayake, and U.R Sangakkara (eds). Proceedings of the 6th International Conference on Kyusei Nature Farming. South Africa, 1999.

© 2014 Global Journals Inc. (US)

6. Kearl LC (1982). Nutrient requirement of ruminants in developing countries. Internationalfeedstuffs Institute, Utah state University, Longan, USA.
7. Khan ML, Ullah I, Javed MT (1992). Comparative study of probiotic T.M. 50. Bovine 40 and Albac on the performance of broiler chicks Pak. Vet. J., 12: 145-47.
8. Khin T, Htun S, Bu MTN, Yu SS (2008). Effects of Feeding Probiotics as Supplement on Growth Performance of Young Calves. GMSARN International Conference on Sustainable Development: Issues and Prospects for the GMS.
9. Konoplya EF, Higa T (2000). EM application in Animal Husbandrypoultry farming and its action mechanisms (Paper presented at the International Conference on EM technology and Nature Farming, October 2000, Pyongyang, DPR Korea).
10. Li W. J., Y.Z. Ni and H. Umemura. 1995. Effective Microorganisms for Sustainable Animal Production in China. Proceedings of the Conference on Effective Microorganisms for a Sustainable Agriculture and Environment, Paris, France.
11. Maqbool A, Hayat S, Hussain1T, Ahmad F, Mahmood K, and Mahmood F (2001). Use of Effective Microorganisms against Deg Nala Disease in Buffaloes in the Rice Growing Areas of Punjab Province (Pakistan). Faculty of Veterinary Science and Faculty of Agriculture University of Agriculture, Faisalabad, Pakistan1.Department of L. & D.D, Punjab, Pakistan.
12. Maqbool A, Shafiq MK, Khan IA, Mahmood F (1997). Prevalence. aetiology, chemotherapy and control of Deg Nala disease in buffaloes and cattle. Ind. J. Dairy Sci. 50: 1-5.
13. Mead GC, Impey CS (1986). Current progress in reducing Salmonella colonization of poultry by 'competitive exclusion'. J. Appl. Bacteriol. Symp. Suppl. 61:67-75.
14. SAS (Statistical Analytical System) (2002). SAS Institute Inc., Cary, NC, USA.
15. Saili T, Marsetyo, Poppi DP, Isherwood P, Nafiu L, Quigley SP (2010). Effect of treatment of cocoa-pods with Aspergillus niger on liveweight gain and cocoa-pod intake of Bali (Bos sondaicus) cattle in South-East Sulawesi. Animal Production Science 50(7) 693–698.
16. Sissons, JW (1989). Potential of probiotic organisms to prevent diarrhea and promote digestion in farm animals - a review. J. Sci. Food Agric. 49:1.
17. Stavric S, Gleeson TM, Blanchfield B (1991). Efficacy of undefined and defined bacterial treatment in competitive exclusion of Salmonella from chicks. Pages 323-330.
18. In: Colonization Control of Human Bacteria Enteropathogens in Poultry. L.C.Blankenship, ed. Academic Press, Inc., New York.
19. Tilley JMA, Terry RA (1963). A two-stage technique for in vitro digestion of forage crops. J.Br. Grassland Soc. 18:104.
20. Van Soest PJ, Robertson JB (1985). Analysis of Forage and Fibrous Foods. A laboratory manual for Animal science 613 Cornell University, Ithaca, New York, USA.

© 2014 Global Journals Inc. (US)

Global Journal of Science Frontier Research: D
Agriculture and Veterinary
Volume 14 Issue 9 Version 1.0 Year 2014
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Role of Camel's Milk in Growth Rate of *Chroococcus* Sp

By Khalid G. Al-Fartosi, Roaa Jafar Elkheralla & Auhood Kadhim Zaid

University of Thi-Qar, Iraq

Abstract- This study aimed to examine the effect of camel's milk against growth of *Chroococcus* sp. the algae farms were divided into five groups as following: the first group (I) is the control group, contained 100 ml from *Chroococcus* sp. algae, the second group (II) contained 100 ml of *Chroococcus* sp. alga with 1 ml of camel's milk, the third group (III) contained 100 ml of *Chroococcus* sp. alga with 2 ml of camel's milk, the fourth group (IV) contained 100 ml of *Chroococcus* sp. alga with 3 ml of camel's milk and the fifth group (V) contained100 ml of *Chroococcus* sp. alga with 5 ml of camel's milk. All groups were noted during the time periods (today's first , the third day, the seventh day and the ninth day).

The results showed the camel's milk caused a significant increase($p<0.05$) in the rate of growth of alga over a period the experiment.

Keywords: *cyanobacteria, chroococcus alga, camel's milk.*

GJSFR-D Classification : *FOR Code: 070799p*



Strictly as per the compliance and regulations of :

© 2014. Khalid G. Al-Fartosi, Roaa Jafar Elkheralla & Auhood Kadhim Zaid. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Role of Camel's Milk in Growth Rate of *Chroococcus* Sp

Khalid G. Al-Fartosi [α], Roaa Jafar Elkheralla [σ] & Auhood Kadhim Zaid [ρ]

Abstract- This study aimed to examine the effect of camel's milk against growth of *Chroococcus* sp. the algae farms were divided into five groups as following: the first group (I) is the control group, contained 100 ml from *Chroococcus* sp. algae, the second group (II) contained 100 ml of *Chroococcus* sp. alga with 1 ml of camel's milk, the third group (III) contained 100 ml of *Chroococcus* sp. alga with 2 ml of camel's milk, the fourth group (IV) contained 100 ml of *Chroococcus* sp. alga with 3 ml of camel's milk and the fifth group (V) contained100 ml of *Chroococcus* sp. alga with 5 ml of camel's milk. All groups were noted during the time periods (today's first , the third day, the seventh day and the ninth day).

The results showed the camel's milk caused a significant increase($p<0.05$) in the rate of growth of alga over a period the experiment.

Keywords: cyanobacteria, chroococcus alga, camel's milk.

I. Introduction

Cyanobacteria are a very old group of organisms and represent relics of the oldest photoautotrophic vegetation in the world that occur in freshwater ,marine and terrestrial habitats (Mundt and Teuscher, 1988).

Chroococcus, a unicellular organism that is a genus of cyanobacteria, like all cyanobacteria, *Chroococcus* is a prokaryote and therefore lacks any of the membranous organelles of eukaryotes. Known for its underwater habitat, *Chroococcus* prefers the sludge of lake and river bottoms (Ditty *et al.*, 2003).

Cyanobacterial secondary metabolites represent a vast diversity of structures and have been isolated from a number of cyanobacterial genera from different geographical locations. During the last two decades, cyanobacterial secondary metabolites have attracted the attention of researchers mainly due potential therapeutic use of several secondary metabolites include a range of compounds showing animal toxicity and antibacterial, antifungal, anti-inflammatory, antimalarial, antiprotozoal, antituber-culosis, antiviral, antitumor and cytotoxic activities (Gademann and Portmann, 2008; Mayer *et al.,* 2009)

In addition to the role of cyanobacteria in the application of biotechnology in the production of hydrogen gas and the production of ammonia and nitrogen fixation and production of enzymes and pharmaceutical products and use food protein for the treatment of heavy water and the excretion of toxins and their impact on the environment in addition to its role in the analysis of crude oil and some of its derivatives to simple compounds non-polluting (Kumazawa and Mitsui, 1989; Kostreba, 2001; Raghukumar, 2001)

The camel's milk is characterized immune-ological advantages unique, since it has a good quality of proteins of the preventive efficacy against bacteria and viruses such as lactoperoxidase, lysozyme and as well lactoferrin. It has an important role in reducing the ability of germs because of injury, inhibition of enzyme cysteine protease found in bacteria and viruses as well as some parasites, the milk also contains a high concentration of minerals(Sodium, Potassium ,iron ,copper ,zinc and magnesium) and vitamins A, B2,C and E (Ohashi *et al.*, 2003; Kamal *et al.*, 2007 ; Al-Hashem *et al.*, 2009).

The present study aimed to investigate the role of camel's milk in the growth of moss through by addition of different concentrations of milk to the farmer of *Chroococcus sp.*

II. Materials and Methods

a) Isolation and purification of algae

Isolated and scrubbed algae according to (Stien,1973), as was access to unialgal cultures were then purified for the purpose of obtaining axenic cultures depending on the method of (Al-Arajy, 1996) and then diagnosed based on)Desikachary,1959; Prescott, 1975).

b) Development and propagation of algae

Algae were grown using a the middle Chu-10 axis by (Al-Arajy, 1996) and after obtaining sufficient amounts transferred to the 100 ml bottles filled with 70 ml of the former the middle and incubated at a temperature (25 ±3) ˚ C .

c) Measuring the rate of growth

The growth rate of algae counted directly by Chamber Shidu (Coombs et al., 1986).

d) Camel's milk

Milk samples were collected early in the morning from camel farm in the Al-Nassiriya city, Thi-Qar province, Iraq. Milk was collected from camel by hand milking as normally practiced by the farmers. The

Author α σ: Department of Biology,College of Science, University of Thi-Qar, Iraq. e-mail: Khalidalfartosi@yahoo.com

Author ρ: Department of Chemistry,College of Science, University of Thi-Qar, Iraq.

© 2014 Global Journals Inc. (US)

samples were collected in sterile screw bottles and kept in cool boxes until transported to the laboratory.

e) experimental groups

Different concentrations of camel's milk and added to the algal farms which were divided into five groups as following:

Group I : control group, contained 100 ml from Chroococcus sp. only.

Group II : contained 100 ml of Chroococcus sp. alga with 1 ml of camel's milk (1%).

Group III : contained 100 ml of Chroococcus sp. with 2 ml of camel's milk (2%).

Group IV : contained 100 ml of Chroococcus sp. with 3 ml of camel's milk (3%).

Group V : contained 100 ml of Chroococcus sp. with 5 ml of camel's milk(4%).

III. Results

The results of the present study shown in table (1). The results indicated the camel's milk caused a significant increase($p< 0.05$) in the growth of algae Chroococces for the duration of the experiment with different concentrations of milk (1, 2, 3and 5ml/100ml of farms) compared with the control group . The highest rate of growth of algae on the first day in group(V).

Results showed the first day a significant increase ($p< 0.05$) in the growth of rate of Chroococces in groups (II,III ,IV and V) compared with group (I) after one day of treatment with different concentrations (1 %, 2 %, 3 % and 5 %) of camel's milk . Also , there was a significant increase ($p< 0.05$) in the growth rate groups(III) and (V) compared with group(II) ,While, the results recorded a significant decrease ($p <0.05$) in the growth of rate of Chroococces in group(IV) compared with group(II) the same day .

After third day of treatment , the results ¡ndicated a significant increase ($p< 0.05$) in the growth of rate of Chroococces in groups (II,III ,IV and V) compared with control group (group I) , Also , there was a significant increase ($p< 0.05$) in the growth rate groups(III) and (V) compared with group(II) ,While, the growth rate in group(IV) decreased significantly ($p <0.05$) compared with group(II) .

After fifth day of treatment , the results indicated a significant increase ($p< 0.05$) in the growth of rate of Chroococces in groups (II,III ,IV and V) compared with control group (group I) , Also , there was a significant increase ($p< 0.05$) in the growth rate group (III) compared with group(II).While , the results recorded a significant decrease ($p <0.05$) in the growth of rate of Chroococces in groups (IV) and (V) compared with group(II).

After seventh day of treatment , the results indicated a significant increase ($p> 0.05$) in the growth of rate of Chroococces in groups (II) and (III) compared with control group , While , the results recorded a significant decrease ($p <0.05$) in the growth of rate of Chroococces in groups (IV) and (V) compared with group(I). The growth rate of Chroococces decreased significantly ($p <0.05$) in groups (IV) and (V) compared with group(II). While, the results recorded a significant increase ($p< 0.05$) in group (III) compared with group(II).

After ninth day of treatment , the results indicated a significant increase ($p< 0.05$) in the growth of rate of Chroococces in groups (II) and (III) compared with control group , While , the results recorded a significant decrease ($p< 0.05$) in the growth of rate of Chroococces in groups (IV) and (V) compared with group(I). Also, there was a significant decrease ($p< 0.05$) in the growth of rate of Chroococces in groups (IV) and (V) compared with group(II). While , the results recorded a significant increase ($p< 0.05$) in group (III) compared with group(II).

Table 1 : Role of camel's milk in growth rate of chroococcus sp

treatments	Growth rate (cells\1 ml)				
	1 day	3 days	5 days	7 days	9 days
Group I	93.00 ±1.73 ^{A}	65.33 ±2.91 ^{D}	71.33 ±0.88 ^{D}	86.00 ±3.06 ^{B}	78.00 ±5.77 ^{C}
Group II	125.67 ±2.96 ^{A}	80.00 ±2.89 ^{D}	119. 0 ±0.58 ^{A}	109.67 ±5.78 ^{B}	100.00 ±4.62 ^{C}
Group III	139.67 ±2.91 ^{B}	152.00 ±1.15 ^{A}	145.00 ±2.89 ^{B}	127.33 ±1.45 ^{C}	109.67 ±5.78 ^{D}
Group IV	109.67 ±5.78 ^{A}	78. 0 ±1.15 ^{C}	94.0 ±1.15 ^{B}	73.00 ±1.73 ^{CD}	70.67 ±1.76 ^{D}
Group V	195.67 ±2.96 ^{A}	92.00 ±1.15 ^{B}	91.33 ±0.88 ^{B}	73.33 ±0.88 ^{C}	68.00 ±1.15 ^{D}

Differences letters refers to a significant differences at $p<0.05$

© 2014 Global Journals Inc. (US)

IV. Discussion

Algae are many benefits as a source for the production of many materials economically important, which encouraged the cultivation and production of biomass, has been used for the production of pharmaceuticals, cosmetics different materials other industrial as well as animal feed and food additives for human food (Grima *et al*., 1990). As it raises to the surroundings of many materials, such as simple sugars and multiple, alcohols and enzymes, and some of which secreted material impact of anti-bacterial, fungi and tumors (Moore *et al*., 1996).

The increase of growth rates of *Chroocecces* sp. may be attributed to supply of algae with nutrients and minerals important for growth, where is the camel's milk is rich in minerals (sodium, potassium, iron, copper, zinc, magnesium) and vitamins (A, B2, C, E) also contains a high concentration of a protein similar to insulin (Kamal *et al.,* 2007; Al-Hashem *et al.,* 2009). These materials play a fundamental role in the increase of cell divisions and increase the breadth and elongation of the cells and thus increasing the effectiveness of the physiological processes photosynthesis thus, a supply algae carbohydrate materials manufactured in this process and this in turn improves the rate of growth (Sahaf , 2000)

V. Conclusion

Our results demonstrate that camel milk is able to events noticeable change in the rate of growth of algae. Algae contains many chemical compounds that can be used in medical, economically and industrially. Further studies are required, using a human population, to confirm theses protective effects.

References Références Referencias

1. Al- Hashem, F.; Dallak, M.; Bashir, N. and Abbas, M. Camel's milk protects against cadmium chloride induced toxicity in white albinorats. *American Journal of Pharmacology and Toxicity.;* 4(3):107 – 117 (2009).
2. Kostreba, L. Oil spill remediation efforts in the middle east. *Internet file: IA:* Oil spill remediation efforts in the middle east, (2001).
3. Moore, R.; Smith, D.; Patterson, M. and Susan, L. The search for new anti-tumor drugs from blue-green algae. *Curr. Pharm.* Design, 2:317-330 (1996).
4. Al-Arajy, M. Studies on the mass culture of some micro algae as food for fish Larvae. ph. D. thesis, Basrah. Univ. (1996).
5. Desikachary, T. V. Cyanophyta Indian .Concil of Agricultural Research, New Delhi, India (1959).
6. Ditty, J.; Williams, S. and Golden, S. A cyanobacterial circadian timing mechanism . *Annu. Rev. Genet.;* 37:513 – 43 (2003).
7. Gademann, K. and Portmann, C. Scondary metabolites from cyanobacteria : complex structure and powerful bioactivities. *Curr. Org. Chem.,*12:326-341 (2008).
8. Grima, F.; Fernaandes, F. ; Camacho, F. and Chisti, Y. Bioreactors light regine , mass transfer and scalen . *P. J. Biotech.;* 70:231 – 247 (1990).
9. Kamal , A.; Salama , O. and El-Saied , K. Changes in amino acids profile of camel's milk protein during the early lactation .*Int. J. Dairy.* Sci.,2:226-234 (2007).
10. Kumazawa, S. and Mitsui, A. Photosynthetic Properties of Synchronously grown aerobic nitrogen- fixing unicellular cyanobacterium. In: Abst. Ann. Meeting of *Amer. Soc.* Microbiol. P268 (1989).
11. Mayer, A.; Rodrigue, Z. ; Berlink, R. and Hamann, M. Marine pharmacology in 2005 – 6 : Marine compounds with anthelmintic, antibacterial, anticoagulant, antifungal, anti–inflammatory, antimalarial, antiprotozoal, antituber-culosis and antiviral activities; affecting the cardio-vascular, immune and nervous systems, and other miscellaneous mechanisms of action. *Biochim. Biophys. Acta.,* 1790:283 – 308 (2009).
12. Mundt, S. and Teuscher, E. Blue – green algae as a source of Pharmacologically – active compound. *Pharmazie.;* 43:809 – 815 (1988).
13. Ohashi, A.; Murata , E.; Yamamoto, K.; Majima, E.; Sano, E.; Le, Q. and Katunuma, K. New functions of lactoferrin and beta-casein in mammalian milk as cysteine protease inhibitors. *Biochem. Biophys. Res. Commum.,* 20:98-103 (2003).
14. Prescott, G. Algae of the Western great lake area. Ellion C., Brown Co. pub., Dugugue, Lowa, USA (1975).
15. Raghukumar, C.; Vipparty, V.; David, J. and Chandramohan, D. Degradation of crude oil by marine cyanobacteria. *Abs. Appl. Microbiol. and biotechnol. Vol.* 57: 433-436 (2001).
16. Sahaf, H. Applied Plant Nutrition, Press House of Wisdom, Baghdad University, Ministry of Higher Education and Scientific Research, the Republic of Iraq, (1989).
17. Stein, J. R. Handbook of phycological methods. Cambridge Univ. *press.* Cambridge, UK. (1973).

© 2014 Global Journals Inc. (US)

This page is intentionally left blank

© 2014 Global Journals Inc. (US)

Global Journal of Science Frontier Research: D
Agriculture and Veterinary
Volume 14 Issue 9 Version 1.0 Year 2014
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Aquatic Biodiversity Analysis of Four Rice Terraces Clusters in the Cordillera Region, Northern Philippines

By Robert T. Ngidlo & Nestor M. Baguinon
Ifugao State University, Philippines

Abstract- An aquatic biodiversity analysis was conducted in four rice terraces clusters in the Cordillera region in the northern central part of the Philippines. Data was obtained using the quadrat sampling method. The location of sampling activities are the four rice terraces clusters namely: (1) Asipulo, Ifugao; (2) Natonin, Mt Province; (3) Bagumbayan, Kalinga; and (4) Tanglagan, Apayao. Each place of sampling had two (2) blocks: (1) Rice paddy fields and (2) Irrigation canals/creeks. The number of quadrats randomly taken in each block was ten (10) plots with each quadrat measuring 1m x 1m. The number of plots in all places was 4 sites x 2 blocks x 10 plots = 80 plots. Individuals of shell, crab, fish and frog species circumscribed within plot/quadrat boundaries were counted, identified and sorted according to species. The importance value (IV) of each species in each block was determined. This was done by taking the sum of Relative Density (%) and Relative Frequency (%) of each species in a given block. Importance Value = Relative Density + Relative Frequency. The Shannon Wiener Diversity Index (H), Evenness Index (E) and Simpson's Index (C) was obtained for each species. The higher values of H, E and C means the greater diversity character of the community being evaluated.

Keywords: rice terraces, aquatic biodiversity, biodiversity indices, importance value, evenness index.

GJSFR-D Classification : FOR Code: 050202p, 07999



Strictly as per the compliance and regulations of :

© 2014. Robert T. Ngidlo & Nestor M. Baguinon. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Aquatic Biodiversity Analysis of Four Rice Terraces Clusters in the Cordillera Region, Northern Philippines

Robert T. Ngidlo [α] & Nestor M. Baguinon [σ]

Abstract- An aquatic biodiversity analysis was conducted in four rice terraces clusters in the Cordillera region in the northern central part of the Philippines. Data was obtained using the quadrat sampling method. The location of sampling activities are the four rice terraces clusters namely: (1) Asipulo, Ifugao; (2) Natonin, Mt Province; (3) Bagumbayan, Kalinga; and (4) Tanglagan, Apayao. Each place of sampling had two (2) blocks: (1) Rice paddy fields and (2) Irrigation canals/creeks. The number of quadrats randomly taken in each block was ten (10) plots with each quadrat measuring 1m x 1m. The number of plots in all places was 4 sites x 2 blocks x 10 plots = 80 plots. Individuals of shell, crab, fish and frog species circumscribed within plot/quadrat boundaries were counted, identified and sorted according to species. The importance value (IV) of each species in each block was determined. This was done by taking the sum of Relative Density (%) and Relative Frequency (%) of each species in a given block. Importance Value = Relative Density + Relative Frequency. The Shannon Wiener Diversity Index (H), Evenness Index (E) and Simpson's Index (C) was obtained for each species. The higher values of H, E and C means the greater diversity character of the community being evaluated. The results clearly showed that the four sites differ with respect to diversity values. For comparing Rice Paddy versus Canal/Creek, Kalinga and Apayao have higher canal/creek diversity than the rice paddy diversity, i.e. Canal/Creek (H=1.55, E=.97, C=.78) versus Rice Paddy (H=1.04, E=.94, C=.62). The case of Mt. Province is similar, except that the difference between Canal/Creek and Rice Paddy diversities are not so much, or in other words, the former is slightly more diverse than the latter. However, the Ifugao case demonstrated a reverse where H and C diversity indices are greater in the Rice Paddy than in the Canal/Creek although the E at the Canal/Creek had greater value. Key informants identified four factors causing the decline in the population of aquatic biodiversity such as the use of pesticides, excessive collection for food and/or as animal feed, recurrent drought due to climate change and the presence of introduced invasive species.

Keywords: rice terraces, aquatic biodiversity, biodiversity indices, importance value, evenness index.

I. Introduction

Over the years, traditional agricultural systems steadily gained recognition and prominence in the national and international development discussions. Their recognition came primarily due to the relevance of knowledge reposed in these agricultural systems and their possible contribution to sustainable development. The Cordillera region in the northern central part of the Philippines is home to the largest and most extensive traditional rice terraces cultivation in the Philippines and perhaps throughout the Asia-Pacific region. The region is politically subdivided into six provinces and occupy a total land area of 19,294 km^2 made up of rows of great mountain ranges occupying half of Northern Luzon in the Philippines. Its rugged mountainous backbone contains many peaks exceeding 2,000 meters in height with rolling hills and stretches of river valleys along its flanks. With its rugged mountainous backbone, the early settlers of the areahas no other option but to build rice terraces along the slopes for their own survival. Acknowledged to be more than 2,000 years old, the rice terraces represents an indigenous farming system which reflects the people's history and culture. Its management is governed primarily by indigenous systems and institutions which is purely organic in nature. Past literatures written on the rice terraces invariably described them as stable and resilient agro-ecosystem. Some of the rice terraces clusters have been declared as World Heritage site by the United Nation Education, Scientific and Cultural Organization (UNESCO). However, over the last 50 years, there were a series of changes that had occurred in the rice terraces. Foremost of these changes is that farmers are shifting to modern farming systems (Ngidlo, 2013). Today, the rice terraces represents a miniature of both modern and traditional farming systems. The use of high yielding rice varieties, commercial fertilizers, pesticides and the adoption of hand tractors are the leftovers of modern agriculture replacing the traditional organic system.

Aquatic biodiversity is a major food item in the rice terraces. Local people depend much on these biodiversity's to sustain their diet and protein requirement (Ngidlo and Ngohayon, 2009). The loss of aquatic biodiversity may mean a significant loss in the food security status of the local people. The study was conducted to determine aquatic biodiversity indices and to identify factors contributing to the loss of biodiversity in the rice terraces. In the end, the paper will hopefully

Author α σ : Faculty, Ifugao State University, Potia Campus, A. Lista, Ifugao, Philippines and Faculty, University of the Philippines, Los Banos , Laguna, Philippines. e-mail: r.ngidlo@gmail.com

© 2014 Global Journals Inc. (US)

provide a platform for discussions on traditional agricultural systems among various stakeholders.

II. Materials and Methods

The organisms subjected to biodiversity analysis were mollusks, gastropods (snails) namely (1) Agurong, *Jagora asperata (Lamarck)*, (2) Birabid, *Lymnaea (Bullastrus) cumingiana* (3) Bisukol, *Pila luzonica Reeves* , (4) Golden Apple Snail, *Pomacea canaliculata ,* (5) Liddeg, *Angulyagra oxytropis (Benson)* Geslacht and pelecypods (bivalves) namely (6) Tukmem, *Turbicula manilensis* . The lone arthropod was the crustacean (7) Talangka, *Sundathelphusa cagayana* . Vertebrate species consists of the following fish species: (8) Jojo fish; *Anquilla japonica* (10) Mosquito carp, *Gambusia affinis,* (11) Million fish, *Poecilla reticulata* (12) Frog/tadpole (*Rana* sp).

The location of sampling activities are four rice terraces clusters located in four provinces in the Cordillera region in the northern central part of the Philippines. The rice terraces clusters are: (1) Asipulo, Ifugao; (2) Natonin, Mt Province; (3) Bagumbayan, Kalinga; and (4) Tanglagan, Apayao. Each place of sampling had two (2) blocks: (1) Rice paddy fields and (2) Irrigation canals/creeks. The number of quadrats randomly taken in each block is ten (10) plots with each quadrat measuring 1m x 1m. Hence, the number of plots in all places was 4 sites x 2 blocks x 10 plots = 80 plots. Individuals of shell, crab, fish and frog species circumscribed within plot/quadrat boundaries were counted, identified and sorted according to species. The importance value (IV) of each species in each block was determined. This was done by taking the sum of Relative Density (%) and Relative Frequency (%) of each species in a given block. Importance Value = Relative Density + Relative Frequency.

The Shannon Wiener Diversity Index (H), Evenness Index (E) and Simpson's Index (C) was obtained for each species. The higher values of H,E and C means the greater diversity character of the community being evaluated.

Ten (10) key informants were interviewed to identify factors contributing to the loss or increase of aquatic biodiversity in the rice terraces.

III. Results and Discussion

a) Profile of the Rice Terraces Clusters

The rice terraces of the Cordillera region in the Philippines has evolved in recent years and no longer the same organic system they used to be prior to the 1980's. Demographic pressures associated with the need to produce more food escalated the adoption of modern farming practices for the last 50 to 70 years. Three of the selected study sites shifted to modern farming practices similar to that of lowland agriculture while one site still adhere to the traditional farming practices. The profile of the rice terraces clusters are indicated in Table 1.

Table 1 : Profile of the rice terraces

Study Site	Status of Farming System
Asipulo, Ifugao	Adopted modern farming system in the early 1990's
Natonin, Mt. Province	Still continuing with the traditional farming system
Tanglagan, Apayao	Adopted modern farming system in the 1980's
Bagumbayan, Kalinga	Adopted modern farming system in the 1980's

b) Aquatic Biodiversity Analysis

The result of analysis in Table 2 shows that the four sites differ with respect to diversity values. For comparing Rice Paddy versus Canal/Creek, Kalinga and Apayao have higher canal/creek diversity than the rice paddy diversity, i.e. Canal/Creek (H=1.55, E=.97, C=.78) versus Rice Paddy (H=1.04, E=.94, C=.62). The case of Mt. Province is similar, except that the difference between Canal/Creek and Rice Paddy diversities are not so much, or in other words, the former is slightly more diverse than the latter. However, the Ifugao case demonstrated a reverse, H and C diversity indices is greater in the Rice Paddy than in the Canal/Creek although the E at the Canal/Creek had greater value. These aforementioned results are expected since the Canal/Creek cases approximate the usually richer natural ecosystem as opposed to the Rice Paddy agroecosystem (man-made). Translating the above table into a line graph (Figure 1), the pattern becomes obvious, for example, the values are down with Rice Paddy then up with Canal/Creek for Mt. Province, Kalinga and Apayao. The lesser H and C in rice paddies in Mt. Province, Kalinga and Apayao is indicative of the continuous collection for food and or given as feed to animals.. In the case of Apayao, the higher H and C for the irrigation canal/creek is that during the time of sampling there was a *lapat* in force for almost 3 months covering the whole stretch of the sampling area (almost 2 kms. of creek) thus, aquatic biodiversity was protected from exploitation during that period of time. Rice paddies are not covered by the *lapat* and the lesser value of H and C can be attributed to the continuous spraying of pesticides in rice paddies. The lesser H and C in Ifugao can be attributed to the excessive collection for food and for sale in the market. It may also be attributed to the presence of invasive alien species (e.g. Golden Kuhol) in the Ifugao Rice Paddy having spread into the Canal/Creeks. The Golden Kuhol's presence is highest in Asipulo, Ifugao and

© 2014 Global Journals Inc. (US)

negligibly present in the Rice Paddies of Natonin, Mt. Province and Bagumbayan, Kalinga, but absent in their respective Canals/Creeks. Golden Kuhol was not found in Tanglagan, Apayao. The graph showing the H, E and C of the four study sites is shown in Fig. 1

Table 2 : Importance values (IV) of species in all four sites including corresponding Shannon Wiener Diversity, Evenness and Simpson's Indices and contained IVs of exotic species

Species/Variety	Ifugao		Mt. Province		Kalinga		Apayao	
	Rice Paddy	Canal/Creek	Rice Paddy	Canal/Creek	Rice Paddy	Canal/Creek	Rice Paddy	Canal/Creek
Bisukul	3.43		20.08	35.11	38.67	23.77	52.78	42.13
Agurong	78.07	65.75	39.55	55.32	49.36	49.73	43.65	41.2
Liddig			36.17					
Tukmim						26.49		16.67
Birabid			2.18				3.57	
Tadpole/Frog	54.17	89.01		13.1	87.82	41.67		12.04
Crab				86.9		58.33		12.04
Total	135.67	154.76	97.98	190.43	175.85	199.99	100.00	124.08
Total no. species	3	2	4	4	3	5	3	5
Golden kuhol	18.50	34.25	2.02					
Dojo	20.83	10.99						
Million Fish	25				12.18		100	75.92
Mosquito carp			100					
Total	64.33	45.24	102.02	0.00	12.18	0.00	100.00	75.92
Total no. Exotic species	3	2	2		1		1	1
Shannon Div. (H)	0.7775	0.6818	1.1435	1.2129	1.0364	1.5531	0.8181	1.4552
Evenness (E)	0.7078	0.9836	0.8249	0.875	0.9434	0.965	0.7447	0.9042
Simpson's Index (c)	0.51	0.49	0.66	0.67	0.62	0.78	0.53	0.74




Figure 1 : Graph of Shannon Wiener Diversity, Evenness, and Simpson's Indices of all four sites and corresponding blocks including the sum IV of exotic species per block

The average number of individuals per species per plot was computed and presented in Table 3 and 4. The mollusks were tabulated in Table 3 while the non mollusks in Table 4.

© 2014 Global Journals Inc. (US)

Table 3 : Average number of individuals per plot (e.g. average density/plot) per block among native species of mollusks and exotic Golden Kuhol. (average number individuals/sq.m.)

SPECIES	IFUGAO		MT PROVINCE		KALINGA		APAYAO	
	Rice Paddy	Canal-Creek	Rice Paddy	Canal-Creek	Rice Paddy	Canal-Creek	Rice Paddy	Canal-Creek
Bisukol	0.2	0	4.3	1.9	3.3	1.9	6.3	2.3
Agurong	30.9	18.5	12.7	5.7	6.2	6.2	6.1	2.2
Liddig	0	0	12.9	1.8	0	0	0	0
Tukmim	0	0	0	0	0	3.8	0	0.9
Birabid	0	0	0.2	0	0	0	0.2	0
Golden kuhol	1.9	4.2	0.1	0	1.6	0	0	0

Table 4 : Average number of individuals per plot (e.g. average density/plot) per block among non-mollusk species (tadpole/frog, fish and crab)

Species	Ifugao		Mt. Province		Kalinga		Apayao	
	Rice Paddy	Canl/ Creek	Rice Paddy	Canl/ Creek	Rice Paddy	Canl/ Creek	Rice Paddy	Canal/ Creek
Frog/tadpole	0.7	1.2	0	0.2	1.1	0.2	0	0
Crab	0	0	0	1.9	0.1	0	0	0.2
Jojo	0.2	0	0	0	0.1	0.2	0	0
Million fish	0.3	0.1	0	0	0	0.4	1	1.9
Mosquito carp	0	0	84.2	0	0	0	0	0




Figure 2 : The average density per plot for the mollusks

The values in Table 3 were line graphed and shown in Figure 2. Golden apple snail had the highest values in Asipulo, Ifugao. It is also interesting to note that the Native Snail (bisikul) is very low in both Rice Paddy and Canal/Creek of Asipulo, Ifugao while the population density of Golden apple snail is considerable. The data implies that Golden apple snail is outcompeting the native strains of edible snails (bisikul) and perhaps may push the latter to further endangerment or worst even possibly to extinction. The data shows that as the population of golden apple snail decline, there is a corresponding increase in the population of the native strains of the native edible snails (bisikul). Agurong, *(Jagora asperata Lamarck)* is

© 2014 Global Journals Inc. (US)

dominant in all the sites. Agurong is a prolific breeder and least preferred for food by collectors.

The pattern of non-mollusks show little information about the impact of alien fishes to the densities of native species. This could be explained partly by the difficulty of sampling highly mobile subjects, such as swimming fish species, unlike among mollusks which are quite sedentary. Hence, in Figure 3, there is only little pattern to merit explanation. Frogs/tadpoles are high at Canal/Creeks in Ifugao and Mt. Province than corresponding rice paddies, but the reverse is true in Kalinga while their presence in Apayao is zero. The zero population of frogs/tadpoles in Apayao can be attributed to the negative impacts of pesticides continually used by farmers. There is an extremely high density of Talangka (crabs) at the Canal/Creek in Natonin, Mt Province but in the other cases the density is negligible or absent. The rice terraces clusters in Natonin still carry 100% the traditional farming systems which is purely organic in nature. For mobile organisms, such as fishes and crabs, a zero value does not mean they are absent. The sampling method applied, such as random quadrat or plot sampling, may not be the best procedure because they are mobile and therefore their spatial distribution/dispersion in space could be extremely clumped and not normal. For mobile organisms, ecologists usually employ the so called capture-recapture method to determine their population densities. However, the capture-recapture method is applicable for terrestrial organisms and not for aquatic animals.




Figure 3 : The average density per plot for the non-mollusks

There are no historical data to compare the biodiversity of aquatic animals in the rice terraces before and after the transition to modern farming systems. Testimonial evidences provided by key informants confirmed a more diverse past than what it is today. Farmers ascribed the reduction in the population of aquatic biodiversity to four factors: a) impacts of pesticides applied in the rice terraces b) overexploitation or excessive collection c. frequent drought and d. replacement with introduced invasive species such as golden apple snail. For the rice terraces clusters that adopted modern farming practices, majority claimed that the use of pesticides is the primary factor causing the decline in the population of aquatic biodiversity followed by excessive collection and the impacts of alien invasive species such as golden apple snail. Farmers in Tanglagan, Apayao although they have shifted completely to modern farming systems, they still continue to practice an indigenous form of conservation which the locals call *"lapat system"* wherein a portion of a river system of about 2-3 kilometers in length is declared off limit to exploitation for several months or even years. The "lapat" is declared in honor of a person who died in the community and the length of which depends on the prominence and status of the person who died in the community. Table 5 shows the causal factors for the decline in the population of aquatic biodiversity as viewed by farmers.

© 2014 Global Journals Inc. (US)

Table 5 : Causal factors in the loss of biodiversity in the rice terraces

Causal factors of the loss of aquatic biodiversity	Location				Total
	Asipulo, Ifugao	Natonin, Mt. Province	Bagumbayan, Kalinga	Tanglagan, Apayao	
Use of pesticides	6	0	8	10	24
Excessive collection	3	2	2	0	7
either for human food or animal feed					
Insufficient water supply	1	0	0	0	1
Introduced invasive species (golden apple snail, torachuk)	0	8	0	0	8
Total	10	10	10	10	40

IV. Conclusion and Recommendation

The four rice terraces clusters vary in terms of their biodiversity indices. Canals/creeks are more diverse than the rice paddy system. The canals and creeks are natural ecosystems less disturbed by human activities. The use of pesticides is clearly altering the population of aquatic biodiversity. On the other hand, invasive species such as golden apple snail pose a threat to the population of the native strains most preferred by the local people for food. In areas where golden apple snail is prevalent, it may be necessary to encourage their removal not by chemical means but by manual collection and given to animals as feed materials.

For hundreds of years, the varied aquatic life found in the rice terraces customarily meets the protein requirement of the local people. The loss of these aquatic biodiversity certainly affects the food security status of rice terraces farmers and their families. Efforts should be made to protect aquatic biodiversity by limiting the use of pesticides and other forms of harmful chemicals. The farming community in Tanglagan, Apayao have shown an excellent form of conserving aquatic biodiversity through the *"lapat system"* handed down from past generations and carried over by the present generation. The *"lapat system"* can be replicated to other areas in the Cordillera region where rice terraces cultivation persists. However, since the *lapat system* is indigenous only to Apayao and not to the other provinces in the Cordillera region, their adoption requires the enactment of local policies by the Local Government Units to establish a framework for their adoption and transfer to the other provinces.

V. Acknowledgement

The authors acknowledged with sincerest gratitude the Neys-Van Hoogstraten Foundation (NHF) of the Netherlands for providing the fund that was used in the implementation of the study. Likewise recognition is also due to the Ifugao State University (where the author work) for the support given during the implementation of the research project.

References Références Referencias

1. Ngidlo, R.T. and S.L. Ngohayon. 2009. Conservation and Adaptive Management of the Ifugao Rice Terraces: A Case Analysis. Upland Farm Journal, Ifugao State University, Nayon, Lamut, Ifugao, Philippines Vol. XVIII, No. 1.
2. Ngidlo, R.T. 2013. The Rice Terraces in Ifugao Province, Northern Philippines: Current Scenario, Gaps and Future Directions. Global Journal of Biology, Agriculture and Health Sciences. Global Institute for Research and Education. Vol. 2(4) 151-154.

© 2014 Global Journals Inc. (US)

Global Journal of Science Frontier Research: D
Agriculture and Veterinary
Volume 14 Issue 9 Version 1.0 Year 2014
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

The Effects of Feeding Time and Regimen on the Performance of Broiler Breeders: A review

By Modu, M. A, Benisheikh, A. A., Jibrin, T. A., Marte A. M. & Ali A. M.

Yobe state university, Nigeria

Abstract- Feeding time and regimen have the potentials to influence the performance of broiler breeder flocks and are thus of great importance. A literature review is presented concerning the responses of broiler breeders to feeding time and feeding regimen. There is a potential improvement in shell quality, egg weight and egg production that results from feeding later in the day (afternoon feeding)or from splitting daily feed allocation (split feeding). Broiler breeders are usually given a limited daily feed allowance in the morning. This practice does not supply nutrients to coincide with the hen's need.

Keywords: *feeding, time, broiler, breeders.*

GJSFR-D Classification : *FOR Code: 309999*



Strictly as per the compliance and regulations of :



© 2014. Modu, M. A, Benisheikh, A. A., Jibrin, T. A., Marte A. M. & Ali A. M. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

The Effects of Feeding Time and Regimen on the Performance of Broiler Breeders: A Review

Modu, M. A [α], Benisheikh, A. A. [σ], Jibrin, T. A. [ρ], Marte A. M. [ω] & Ali A. M. [¥]

Abstract- **Feeding time and regimen have the potentials to influence the performance of broiler breeder flocks and are thus of great importance. A literature review is presented concerning the responses of broiler breeders to feeding time and feeding regimen. There is a potential improvement in shell quality, egg weight and egg production that results from feeding later in the day (afternoon feeding)or from splitting daily feed allocation (split feeding). Broiler breeders are usually given a limited daily feed allowance in the morning. This practice does not supply nutrients to coincide with the hen's need.**

Keywords: feeding, time, broiler, breeders.

I. Introduction

Genetic selection for growth parameters in meat type chickens gives rise to a parent stock (broiler breeders) that tend to lack the ability to self-regulate feed in-take. As such, their high body mass is associated with excessive fat deposition, lameness, and high mortality rate (often due to skeletal or cardiovascular disease or both). To regulate weight gain, limit health risks and also maintain high fertility for the parent stock encompass a high degree of feed restriction (Renema and Robinson, 2004). It is a common practice to feed broiler breeder flocks early in the morning, at or near the start of the photoperiod. It has been argued that this is an inefficient practice which does not supply nutrients to coincide with the hen's need Cave, (1981), and that feeding later in the day, or splitting the single feed allocation across more frequent periods throughout the day (split feeding) improves the performance of broiler breeders. This restricted amount of feed is rapidly consumed; therefore, the hens fast for an extended period of time before their next feeding resulting to improved performance. These include laying performance, shell quality, reproductive efficiency, oviposition times, birds' welfare and behavioural pattern.

II. Effects of Feeding Time and Regimen on Egg Production

Onagbesan *et al.,* (2006) reported that hens with free access to feed had comparable egg with those on split fed birds in the early production period (27 to 29 week), but their egg production decreased dramatically thereafter. Decreased egg production in broiler breeders with ad libitum feeding is well documented (Sun *et al.,* 2006, Chen *et al.,* 2006, Robinson *et al.,* 1993).Two classical studies of the effects of feeding time in laying hens Balnave, (1977); Daniel and Balnave, (1981) reported higher egg production in afternoon-fed hens than in morning-fed hens. Simon, (1973) observed a tendency for hens fed in the morning to eat smaller amounts of feed than those fed in the evening. Furthermore, Bougon, (1973) observed a greater feed intake in birds fed two hours separate meals, one in the morning and one in the evening , than birds fed a single four hours meal in the morning. Indeed in the classical studies performed by Balnave (1977) and Daniel and Balnave, (1981), the afternoon-fed hens ate more than their morning fed counter parts. Therefore, the improvements in egg production due to afternoon feeding in these studies were most likely due to increased feed intake by the afternoon-fed hens.

III. Effects of Feeding Time and Regimen on Shell Quality

The provision of a limited daily allowance of feed in the morning may not supply nutrients to coincide with the broiler breeder hen's need Cave, (1981). This is particularly the case with Calcium (Ca), an essential component of egg shells. Shell formation normally commences in the afternoon or evening and thus utilization of dietary Ca in the morning-fed broiler breeder is poor due to the inability of these birds to meter Ca from the crop to the lower digestive tract at a uniform rate Farmer *et. al.*,(1983). Feeding broiler breeders later in the day supplies dietary calcium (Ca) at times that correspond more closely to periods of shell deposition ,resulting in improved calcium utilization, which is usually manifested as an increased in egg specific gravity, shell quality, shell weight and thickness Farmer *et. al.,(*1983). Lewis and Perry showed that split feeding could improve shell quality.

Increased embryonic mortality, and thus reduced hatchability, is associated with eggs with a specific gravity ofless than 1.080, McDaniel *et.al,* (1981). Therefore,employing a later feeding time is often viewed as a way to improve hatchability, due to the positive influence that later times have on shell quality.

Author α: Yobe state university farm and recreational Park, P.M.B. 1144, Damaturu, Nigeria.
Author σ ρ ω ¥: Department of Animal Production Technology, Ramat Polytechnic, Maiduguri, Borno State, Nigeria.
e-mail: mammanagro@yahoo.com

© 2014 Global Journals Inc. (US)

IV. Effects of Feeding Time and Regimen on Egg Weight

Spradley *et al.*, (2008) reported that broiler breeders fed twice a day laid heavier eggs than those fed once a day. They also had better overall body weight uniformity. Harms et.al., (1991) reported that later feeding times or split feeding results in heavier egg weight than the earliest feeding times. Therefore, there is a general trend for egg weight to increase when birds are fed later in the day, or when the daily feed allocation is spread over more frequent feeding periods.

It is unlikely that larger eggs will provide any advantage for hatching egg producers, particularly in the later stages of the production cycle, when attempts should be made to control egg size from older flocks as much as possible. However, some advantage gained from afternoon feeding during the early phases of production in order to increase the number of settable eggs from young flocks.

V. Effects of Feeding time and Regimen on Reproductive Efficiency and Oviposition Time

Farmer *et.al*, (1983) reported that volume of semen and absolute numbers of spermatozoa yielded by males were found to be highest in the afternoon. Correspondingly, mating is more frequent in the afternoon than in the morning, Cave, (1981). The distribution of mating throughout the day is affected by the ovulatory cycle, as the oviduct is not receptive to spermatozoa near the time of oviposition Cave, (1981). Most spermatozoa deposited within 1 to 3 hours prior to or just after oviposition are eliminated by vaginal contractions involved in the process of oviposition Farmer et.al., (1983).

VI. Effects of Feeding Time and Regimen on Welfare

Numerous reports have shown that restricted fed broiler breeders experience hunger and frustration due to a high motivation for feeding Cave, (1981), Balnave, (1977) and Bougon, (1973). Split feeding offers a potential method of reducing frustration and possibly hunger, Samara *et al.*, (1996) concluded that feeding twice a day during rearing significantly improved broiler breeder welfare, at least as indicated by the parameters of hunger and frustration under study.

VII. Conclusion

The current commercial practice of feeding broiler breeders early in the day, at or near lights-on does not supply nutrients to coincide with the broiler hens needs. Afternoon feeding or splitting the daily feed allowance across more frequent regime may be options to consider, as these practices lead to improvements in shell quality, egg weight and egg production.

References Références Referencias

1. *Balnave, D. (1977).* The effect of energy restriction of laying hens given either a single morning or single evening meal. *British poultry science 18: 115-119*
2. *Bougon, M. B. (1973).* When to restrict layers feed for best results (part 2). Poultry world, 15th February: 34-35.
3. *Cave, N. A. (1981).* Effect of diurnal programs of nutrient intake on the performance of broiler breeder hens. *Poultry science 60: 1287-1292.*
4. *Chen, S. E, Mc Murty, J. P. , and Walzem, R. L.* (2006). Over feeding inducedovarian dysfunction in broiler breeder hens is associated with lipotoxicity. *Poultry. Sci. 86:70–81*
5. *Daniel, M. and Balnave, D. (1981).* Responses of crossbred layers fed specific meal times. *British poultry science 22: 347-354.*
6. *Farmer, M. and Roland, D.A. (1982).* Calcium utilization in the laying hen. *Poultry science. 61-1378(Abstract).*
7. *Harms, R. H. (1991).* The influence of changing time of feeding on performance of broiler breeder hens. *Poultry science 62: 456-471.*
8. Mc Daniel, G.R., Brake, J and Eckman, M.K (1981). Factors affecting broiler breeder performance. *Poultry science 58: 10-13.*
9. *Onagbesan, O. M., Metayer, S., Tona, K., Williams, J., Decuypere, E., and Bruggeman, V. (2006).* Effects of genotype and feed allowance on plasma luteinizing hormones, follicle-stimulating hormones, progesterone, estradiol levels, follicle differentiation, and egg productionrates of broiler breeder hens. *Poultry. Sci. 85:1245–1258.*
10. *Renema, R. A., and Robinson, F. E. (2004).* Defining normal: Comparison of feed restriction and full feeding of female broiler breeders. *World's Poultry. Sci. J. 60:508–522.*
11. *Robinson, F. E., Wilson, J. L., Yu, M.W., Fasenko, G.M. and Hardin, R. T. (1993).*The relationship between body weight and reproductive efficiency in meat-type chicken. *Poultry. Sci. 72:912–922.*
12. *Samara M.H., Robinson, K.R and Smith, M.O. (1996).* Interaction of feeding time and temperature and their relationship to performance of the broiler breeder hen. *Poultry Science 75: 34-41.*
13. *Simon, J. (1973).* Influence d'un temps d'alimentation limite a deux heures par jour chez la poulette et chez la pondeuse de souche ponte. 4th European Poultry conference, London 203-201.
14. *Spradley, J. M., Freeman, M. E., Wilson, J. L. and Davis, A. J. (2008).*The influence of a twice-a-day feeding regimen after photostimulation on the

© 2014 Global Journals Inc. (US)

reproductive performance of broiler breeder hens. *Poultry. Sci. 87:561–568.*

15. *Sun, J. M., Richards, M. P., Rosebrough, R. W., Ashwell, C. M., McMurtry, J.P., and Coon, C.N.* (2006). The relationship of body composition, feed intake, and metabolic hormones for broiler breederfemales. *Poultry. Sci. 85:1173–11*

© 2014 Global Journals Inc. (US)

Global Journals Inc. (US) Guidelines Handbook 2014

www.GlobalJournals.org

FELLOWS

FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (FARSS)

Global Journals Incorporate (USA) is accredited by Open Association of Research Society (OARS), U.S.A and in turn, awards "FARSS" title to individuals. The 'FARSS' title is accorded to a selected professional after the approval of the Editor-in-Chief/Editorial Board Members/Dean.



- The "FARSS" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSS or William Walldroff, M.S., FARSS.

FARSS accrediting is an honor. It authenticates your research activities. After recognition as FARSB, you can add 'FARSS' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, and Visiting Card etc.

The following benefits can be availed by you only for next three years from the date of certification:

FARSS designated members are entitled to avail a 40% discount while publishing their research papers (of a single author) with Global Journals Incorporation (USA), if the same is accepted by Editorial Board/Peer Reviewers. If you are a main author or co-author in case of multiple authors, you will be entitled to avail discount of 10%.

Once FARSB title is accorded, the Fellow is authorized to organize a symposium/seminar/conference on behalf of Global Journal Incorporation (USA). The Fellow can also participate in conference/seminar/symposium organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent.





You may join as member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer. In addition, it is also desirable that you should organize seminar/symposium/conference at least once.

We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



© Copyright by Global Journals Inc.(US) | Guidelines Handbook

The FARSS can go through standards of OARS. You can also play vital role if you have any suggestions so that proper amendment can take place to improve the same for the benefit of entire research community.



As FARSS, you will be given a renowned, secure and free professional email address with 100 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.



The FARSS will be eligible for a free application of standardization of their researches. Standardization of research will be subject to acceptability within stipulated norms as the next step after publishing in a journal. We shall depute a team of specialized research professionals who will render their services for elevating your researches to next higher level, which is worldwide open standardization.



The FARSS member can apply for grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.
Once you are designated as FARSS, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria. After certification of all your credentials by OARS, they will be published on your Fellow Profile link on website https://associationofresearch.org which will be helpful to upgrade the dignity.

The FARSS members can avail the benefits of free research podcasting in Global Research Radio with their research documents. After publishing the work, (including published elsewhere worldwide with proper authorization) you can upload your research paper with your recorded voice or you can utilize chargeable services of our professional RJs to record your paper in their voice on request.



The FARSS member also entitled to get the benefits of free research podcasting of their research documents through video clips. We can also streamline your conference videos and display your slides/ online slides and online research video clips at reasonable charges, on request.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook



The FARSS is eligible to earn from sales proceeds of his/her researches/reference/review Books or literature, while publishing with Global Journals. The FARSS can decide whether he/she would like to publish his/her research in a closed manner. In this case, whenever readers purchase that individual research paper for reading, maximum 60% of its profit earned as royalty by Global Journals, will be credited to his/her bank account. The entire entitled amount will be credited to his/her bank account exceeding limit of minimum fixed balance. There is no minimum time limit for collection. The FARSS member can decide its price and we can help in making the right decision.

The FARSS member is eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get remuneration of 15% of author fees, taken from the author of a respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account.



MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (MARSS)

The ' MARSS ' title is accorded to a selected professional after the approval of the Editor-in-Chief / Editorial Board Members/Dean.



The "MARSS" is a dignified ornament which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., MARSS or William Walldroff, M.S., MARSS.

MARSS accrediting is an honor. It authenticates your research activities. After becoming MARSS, you can add 'MARSS' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, Visiting Card and Name Plate etc.

The following benefitscan be availed by you only for next three years from the date of certification.



MARSS designated members are entitled to avail a 25% discount while publishing their research papers (of a single author) in Global Journals Inc., if the same is accepted by our Editorial Board and Peer Reviewers. If you are a main author or co-author of a group of authors, you will get discount of 10%.

As MARSS, you will be given a renowned, secure and free professional email address with 30 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.



© Copyright by Global Journals Inc.(US) | Guidelines Handbook



We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

The MARSS member can apply for approval, grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.





Once you are designated as MARSS, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria.

It is mandatory to read all terms and conditions carefully.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

AUXILIARY MEMBERSHIPS

Institutional Fellow of Global Journals Incorporation (USA)-OARS (USA)

Global Journals Incorporation (USA) is accredited by Open Association of Research Society, U.S.A (OARS) and in turn, affiliates research institutions as "Institutional Fellow of Open Association of Research Society" (IFOARS).



The "FARSC" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSC or William Walldroff, M.S., FARSC.

The IFOARS institution is entitled to form a Board comprised of one Chairperson and three to five board members preferably from different streams. The Board will be recognized as "Institutional Board of Open Association of Research Society"-(IBOARS).

The Institute will be entitled to following benefits:

The IBOARS can initially review research papers of their institute and recommend them to publish with respective journal of Global Journals. It can also review the papers of other institutions after obtaining our consent. The second review will be done by peer reviewer of Global Journals Incorporation (USA) The Board is at liberty to appoint a peer reviewer with the approval of chairperson after consulting us.

The author fees of such paper may be waived off up to 40%.

The Global Journals Incorporation (USA) at its discretion can also refer double blind peer reviewed paper at their end to the board for the verification and to get recommendation for final stage of acceptance of publication.





The IBOARS can organize symposium/seminar/conference in their country on behalf of Global Journals Incorporation (USA)-OARS (USA). The terms and conditions can be discussed separately.

The Board can also play vital role by exploring and giving valuable suggestions regarding the Standards of "Open Association of Research Society, U.S.A (OARS)" so that proper amendment can take place for the benefit of entire research community. We shall provide details of particular standard only on receipt of request from the Board.



The board members can also join us as Individual Fellow with 40% discount on total fees applicable to Individual Fellow. They will be entitled to avail all the benefits as declared. Please visit Individual Fellow-sub menu of GlobalJournals.org to have more relevant details.

© Copyright by Global Journals Inc.(US) | Guidelines Handbook

We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



After nomination of your institution as "Institutional Fellow" and constantly functioning successfully for one year, we can consider giving recognition to your institute to function as Regional/Zonal office on our behalf.
The board can also take up the additional allied activities for betterment after our consultation.

The following entitlements are applicable to individual Fellows:



Open Association of Research Society, U.S.A (OARS) By-laws states that an individual Fellow may use the designations as applicable, or the corresponding initials. The Credentials of individual Fellow and Associate designations signify that the individual has gained knowledge of the fundamental concepts. One is magnanimous and proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice.

Open Association of Research Society (US)/ Global Journals Incorporation (USA), as described in Corporate Statements, are educational, research publishing and professional membership organizations. Achieving our individual Fellow or Associate status is based mainly on meeting stated educational research requirements.



Disbursement of 40% Royalty earned through Global Journals : Researcher = 50%, Peer Reviewer = 37.50%, Institution = 12.50% E.g. Out of 40%, the 20% benefit should be passed on to researcher, 15 % benefit towards remuneration should be given to a reviewer and remaining 5% is to be retained by the institution.

We shall provide print version of 12 issues of any three journals [as per your requirement] out of our 38 journals worth $ 2376 USD.

Other:

The individual Fellow and Associate designations accredited by Open Association of Research Society (US) credentials signify guarantees following achievements:

- The professional accredited with Fellow honor, is entitled to various benefits viz. name, fame, honor, regular flow of income, secured bright future, social status etc.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

- In addition to above, if one is single author, then entitled to 40% discount on publishing research paper and can get 10%discount if one is co-author or main author among group of authors.
- The Fellow can organize symposium/seminar/conference on behalf of Global Journals Incorporation (USA) and he/she can also attend the same organized by other institutes on behalf of Global Journals.
- The Fellow can become member of Editorial Board Member after completing 3yrs.
- The Fellow can earn 60% of sales proceeds from the sale of reference/review books/literature/publishing of research paper.
- Fellow can also join as paid peer reviewer and earn 15% remuneration of author charges and can also get an opportunity to join as member of the Editorial Board of Global Journals Incorporation (USA)
- • This individual has learned the basic methods of applying those concepts and techniques to common challenging situations. This individual has further demonstrated an in–depth understanding of the application of suitable techniques to a particular area of research practice.

Note :

"

- In future, if the board feels the necessity to change any board member, the same can be done with the consent of the chairperson along with anyone board member without our approval.
- In case, the chairperson needs to be replaced then consent of 2/3rd board members are required and they are also required to jointly pass the resolution copy of which should be sent to us. In such case, it will be compulsory to obtain our approval before replacement.
- In case of "Difference of Opinion [if any]" among the Board members, our decision will be final and binding to everyone.

"

Process of submission of Research Paper

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (*.DOC,*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission.Online Submission: There are three ways to submit your paper:

(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.

(II) Choose corresponding Journal.

(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.

(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.

(C) If these two are not conveninet , and then email the paper directly to dean@globaljournals.org.

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Preferred Author Guidelines

MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11'"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

You can use your own standard format also.
Author Guidelines:

1. General,

2. Ethical Guidelines,

3. Submission of Manuscripts,

4. Manuscript's Category,

5. Structure and Format of Manuscript,

6. After Acceptance.

1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global

Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.

2) Drafting the paper and revising it critically regarding important academic content.

3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.

Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.



To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

5.STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also.Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

Papers: These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

(a)Title should be relevant and commensurate with the theme of the paper.

(b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.

(c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.

(d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.

(e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.

(f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;

(g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.

(h) Brief Acknowledgements.

(i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.

The Editorial Board reserves the right to make literary corrections and to make suggestions to improve briefness.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

Format

Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than 1.4 × 10-3 m3, or 4 mm somewhat than 4 × 10-3 m. Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

Abstract, used in Original Papers and Reviews:

Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art.A few tips for deciding as strategically as possible about keyword search:

- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

Acknowledgements: Please make these as concise as possible.

References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

Tables, Figures and Figure Legends

Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.

Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.

Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.

Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.

6. AFTER ACCEPTANCE

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

6.1 Proof Corrections

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

www.adobe.com/products/acrobat/readstep2.html. This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at dean@globaljournals.org within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

6.3 Author Services

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

6.4 Author Material Archive Policy

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

6.5 Offprint and Extra Copies

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: editor@globaljournals.org .



Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

1. Choosing the topic: In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

2. Evaluators are human: First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

3. Think Like Evaluators: If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

4. Make blueprints of paper: The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

5. Ask your Guides: If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

6. Use of computer is recommended: As you are doing research in the field of Computer Science, then this point is quite obvious.

7. Use right software: Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

8. Use the Internet for help: An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

9. Use and get big pictures: Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

10. Bookmarks are useful: When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

11. Revise what you wrote: When you write anything, always read it, summarize it and then finalize it.

12. Make all efforts: Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

13. Have backups: When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

14. Produce good diagrams of your own: Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

15. Use of direct quotes: When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

16. Use proper verb tense: Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

17. Never use online paper: If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

18. Pick a good study spot: To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

19. Know what you know: Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

20. Use good quality grammar: Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

21. Arrangement of information: Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

22. Never start in last minute: Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

23. Multitasking in research is not good: Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

24. Never copy others' work: Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

25. Take proper rest and food: No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

26. Go for seminars: Attend seminars if the topic is relevant to your research area. Utilize all your resources.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

27. Refresh your mind after intervals: Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

28. Make colleagues: Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

29. Think technically: Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

30. Think and then print: When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

31. Adding unnecessary information: Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

32. Never oversimplify everything: To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

33. Report concluded results: Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

34. After conclusion: Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium though which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

Informal Guidelines of Research Paper Writing

Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.

Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

General style:

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

· Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

· Use standard writing style including articles ("a", "the," etc.)

· Keep on paying attention on the research topic of the paper

· Use paragraphs to split each significant point (excluding for the abstract)

· Align the primary line of each section

· Present your points in sound order

· Use present tense to report well accepted

· Use past tense to describe specific results

· Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives

· Shun use of extra pictures - include only those figures essential to presenting results

Title Page:

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript--must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for briefness. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

Approach:

- Single section, and succinct
- As a outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an conceptual must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.

- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

Procedures (Methods and Materials):

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

Methods:

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

Approach:

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

What to keep away from

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

Results:

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently.You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

Segment Draft and Final Research Paper: You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptive of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.

- Do not give permission to anyone else to "PROOFREAD" your manuscript.

- Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Criterion for Grading a Research Paper (Compilation) by Global Journals Inc. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

Topics	Grades		
	A-B	**C-D**	**E-F**
Abstract	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form Above 200 words	No specific data with ambiguous information Above 250 words
Introduction	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
Methods and Procedures	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
Result	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
Discussion	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
References	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring

INDEX

A

Actinomycetes · 4
Adlibtium · 5
Algal · 14
Angulyagra · 20

C

Chroococcus Alga · 12
Cordillera · 2, 18, 19, 20, 26
Cyanobacteria · 12
Cysteine · 13, 16

E

Enteropathic · 4
Eukaryotes · 12
Extinction · 23

H

Hatchability · 29

L

Lactoferrin · 13, 16
Lactoperoxidase · 13

M

Microflora · 8

P

Paddy · 18, 20, 21, 23

R

Regimen · 2, 28, 29, 30, I
Relics · 12

S

Sulawesi · 11

U

Unialgal · 13



Global Journal of Science Frontier Research

Visit us on the Web at www.GlobalJournals.org | www.JournalofScience.org
or email us at helpdesk@globaljournals.org

ISSN 9755896



© Global Journals