Online ISSN : 2249-4626
Print ISSN : 0975-5896
DOI : 10.17406/GJSFR

GLOBAL JOURNAL

OF SCIENCE FRONTIER RESEARCH: F

Mathematics and Decision Science



White Noise Analysis

Gauss Legendre two Point Rule

Highlights

Visualizing Finite Field

Theory of Hypercomplex Systems

Discovering Thoughts, Inventing Future

VOLUME 16 ISSUE 1 VERSION 1.0

© 2001-2016 by Global Journal of Science Frontier Research, USA

Volume 16 Issue 2 was originally issued as F1

Global Journal of Science Frontier Research: F
Mathematics & Decision Sciences

Volume 16 Issue 2
Originally issued as F1

Global Journal of Science Frontier Research: F

Mathematics & Decision Sciences

Volume 16 Issue 1 (Ver. 1.0)

Open Association of Research Society

© Global Journal of Science Frontier Research. 2016.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Science Frontier Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Science Frontier Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Science Frontier Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website http://globaljournals.us/terms-and-condition/menu-id-1463/

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089
License No.: 42125/022010/1186
Registration No.: 430374
Import-Export Code: 1109007027
Employer Identification Number (EIN):
USA Tax ID: 98-0673427

Global Journals Inc.

*(A Delaware USA Incorporation with "Good Standing"; **Reg. Number: 0423089**)*

Sponsors: Open Association of Research Society
Open Scientific Standards

Publisher's Headquarters office

Global Journals® Headquarters
945th Concord Streets,
Framingham Massachusetts Pin: 01701,
United States of America
USA Toll Free: +001-888-839-7392
USA Toll Free Fax: +001-888-839-7392

Offset Typesetting

Global Journals Incorporated
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,
Pin: CR9 2ER, United Kingdom

Packaging & Continental Dispatching

Global Journals
E-3130 Sudama Nagar, Near Gopur Square,
Indore, M.P., Pin:452009, India

Find a correspondence nodal officer near you

To find nodal officer of your country, please email us at *local@globaljournals.org*

eContacts

Press Inquiries: *press@globaljournals.org*
Investor Inquiries: *investors@globaljournals.org*
Technical Support: *technology@globaljournals.org*
Media & Releases: *media@globaljournals.org*

Pricing (Including by Air Parcel Charges):

For Authors:
22 USD (B/W) & 50 USD (Color)
Yearly Subscription (Personal & Institutional):
200 USD (B/W) & 250 USD (Color)

Integrated Editorial Board
(Computer Science, Engineering, Medical, Management, Natural Science, Social Science)

John A. Hamilton,"Drew" Jr.,
Ph.D., Professor, Management
Computer Science and Software Engineering
Director, Information Assurance Laboratory
Auburn University

Dr. Henry Hexmoor
IEEE senior member since 2004
Ph.D. Computer Science, University at Buffalo
Department of Computer Science
Southern Illinois University at Carbondale

Dr. Osman Balci, Professor
Department of Computer Science
Virginia Tech, Virginia University
Ph.D. and M.S.Syracuse University, Syracuse, New York
M.S. and B.S. Bogazici University, Istanbul, Turkey

Yogita Bajpai
M.Sc. (Computer Science), FICCT
U.S.A.Email:
yogita@computerresearch.org

Dr. T. David A. Forbes
Associate Professor and Range Nutritionist
Ph.D. Edinburgh University - Animal Nutrition
M.S. Aberdeen University - Animal Nutrition
B.A. University of Dublin- Zoology

Dr. Wenying Feng
Professor, Department of Computing & Information Systems
Department of Mathematics
Trent University, Peterborough,
ON Canada K9J 7B8

Dr. Thomas Wischgoll
Computer Science and Engineering,
Wright State University, Dayton, Ohio
B.S., M.S., Ph.D.
(University of Kaiserslautern)

Dr. Abdurrahman Arslanyilmaz
Computer Science & Information Systems Department
Youngstown State University
Ph.D., Texas A&M University
University of Missouri, Columbia
Gazi University, Turkey

Dr. Xiaohong He
Professor of International Business
University of Quinnipiac
BS, Jilin Institute of Technology; MA, MS, PhD,. (University of Texas-Dallas)

Burcin Becerik-Gerber
University of Southern California
Ph.D. in Civil Engineering
DDes from Harvard University
M.S. from University of California, Berkeley & Istanbul University

Dr. Bart Lambrecht
Director of Research in Accounting and FinanceProfessor of Finance
Lancaster University Management School
BA (Antwerp); MPhil, MA, PhD (Cambridge)

Dr. Carlos García Pont
Associate Professor of Marketing
IESE Business School, University of Navarra
Doctor of Philosophy (Management), Massachusetts Institute of Technology (MIT)
Master in Business Administration, IESE, University of Navarra
Degree in Industrial Engineering, Universitat Politècnica de Catalunya

Dr. Fotini Labropulu
Mathematics - Luther College
University of ReginaPh.D., M.Sc. in Mathematics
B.A. (Honors) in Mathematics
University of Windso

Dr. Lynn Lim
Reader in Business and Marketing
Roehampton University, London
BCom, PGDip, MBA (Distinction), PhD, FHEA

Dr. Mihaly Mezei
ASSOCIATE PROFESSOR
Department of Structural and Chemical Biology, Mount Sinai School of Medical Center
Ph.D., Etvs Lornd University
Postdoctoral Training,
New York University

Dr. Söhnke M. Bartram
Department of Accounting and FinanceLancaster University Management SchoolPh.D. (WHU Koblenz)
MBA/BBA (University of Saarbrücken)

Dr. Miguel Angel Ariño
Professor of Decision Sciences
IESE Business School
Barcelona, Spain (Universidad de Navarra)
CEIBS (China Europe International Business School).
Beijing, Shanghai and Shenzhen
Ph.D. in Mathematics
University of Barcelona
BA in Mathematics (Licenciatura)
University of Barcelona

Philip G. Moscoso
Technology and Operations Management
IESE Business School, University of Navarra
Ph.D in Industrial Engineering and Management, ETH Zurich
M.Sc. in Chemical Engineering, ETH Zurich

Dr. Sanjay Dixit, M.D.
Director, EP Laboratories, Philadelphia VA Medical Center
Cardiovascular Medicine - Cardiac Arrhythmia
Univ of Penn School of Medicine

Dr. Han-Xiang Deng
MD., Ph.D
Associate Professor and Research Department Division of Neuromuscular Medicine
Davee Department of Neurology and Clinical Neuroscience Northwestern University Feinberg School of Medicine

Dr. Pina C. Sanelli
Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic Radiology
M.D., State University of New York at Buffalo, School of Medicine and Biomedical Sciences

Dr. Roberto Sanchez
Associate Professor
Department of Structural and Chemical Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University

Dr. Wen-Yih Sun
Professor of Earth and Atmospheric Sciences Purdue University Director National Center for Typhoon and Flooding Research, Taiwan
University Chair Professor
Department of Atmospheric Sciences, National Central University, Chung-Li, TaiwanUniversity Chair Professor
Institute of Environmental Engineering, National Chiao Tung University, Hsin-chu, Taiwan.Ph.D., MS The University of Chicago, Geophysical Sciences
BS National Taiwan University, Atmospheric Sciences
Associate Professor of Radiology

Dr. Michael R. Rudnick
M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and Hypertension Division (PMC)
Penn Medicine, University of Pennsylvania
Presbyterian Medical Center, Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of Internal Medicine

Dr. Bassey Benjamin Esu
B.Sc. Marketing; MBA Marketing; Ph.D Marketing
Lecturer, Department of Marketing, University of Calabar
Tourism Consultant, Cross River State Tourism Development Department
Co-ordinator , Sustainable Tourism Initiative, Calabar, Nigeria

Dr. Aziz M. Barbar, Ph.D.
IEEE Senior Member
Chairperson, Department of Computer Science
AUST - American University of Science & Technology
Alfred Naccash Avenue – Ashrafieh

President Editor (HON.)

Dr. George Perry, (Neuroscientist)
Dean and Professor, College of Sciences
Denham Harman Research Award (American Aging Association)
ISI Highly Cited Researcher, Iberoamerican Molecular Biology Organization
AAAS Fellow, Correspondent Member of Spanish Royal Academy of Sciences
University of Texas at San Antonio
Postdoctoral Fellow (Department of Cell Biology)
Baylor College of Medicine
Houston, Texas, United States

Chief Author (HON.)

Dr. R.K. Dixit
M.Sc., Ph.D., FICCT
Chief Author, India
Email: authorind@computerresearch.org

Dean & Editor-in-Chief (HON.)

Vivek Dubey(HON.)
MS (Industrial Engineering),
MS (Mechanical Engineering)
University of Wisconsin, FICCT
Editor-in-Chief, USA
editorusa@computerresearch.org

Sangita Dixit
M.Sc., FICCT
Dean & Chancellor (Asia Pacific)
deanind@computerresearch.org

Suyash Dixit
(B.E., Computer Science Engineering), FICCTT
President, Web Administration and
Development , CEO at IOSRD
COO at GAOR & OSS

Er. Suyog Dixit
(M. Tech), BE (HONS. in CSE), FICCT
SAP Certified Consultant
CEO at IOSRD, GAOR & OSS
Technical Dean, Global Journals Inc. (US)
Website: www.suyogdixit.com
Email: suyog@suyogdixit.com

Pritesh Rajvaidya
(MS) Computer Science Department
California State University
BE (Computer Science), FICCT
Technical Dean, USA
Email: pritesh@computerresearch.org

Luis Galárraga
J!Research Project Leader
Saarbrücken, Germany

Contents of the Issue

i. Copyright Notice
ii. Editorial Board Members
iii. Chief Author and Dean
iv. Contents of the Issue

1. Lyapunov-Type Inequality for Fractional order Difference Equations. *1-9*
2. A Construction of Non-Gaussian White Noise Analysis using the theory of Hypercomplex Systems. *11-25*
3. Visualizing Finite Field of Order p^2 through Matrices. *27-30*
4. Mixed Quadrature Rule for Double Integrals of Simpsons $\frac{1}{3}rd$ and Gauss Legendre Two Point Rule in Two Variables. *31-37*

v. Fellows
vi. Auxiliary Memberships
vii. Process of Submission of Research Paper
viii. Preferred Author Guidelines
ix. Index

Global Journal of Science Frontier Research: F
Mathematics and Decision Sciences
Volume 16 Issue 1 Version 1.0 Year 2016
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Lyapunov-Type Inequality for Fractional order Difference Equations

By Meihong Cui, Jingxu Xin, Xianmei Huang & Chengmin Houx
Yanbian University, China

Abstract- In this paper, we obtain some Lyapunov-type inequalities for a class of fractional order difference equations with homogeneous boundary value conditions, the results of this paper are new and generalize some early results in the literature.

Keywords: lyapunov-type inequalities, fractional order difference equations, boundary value problem.

GJSFR-F Classification : MSC 2010: 37B25



Strictly as per the compliance and regulations of :

© 2016. Meihong Cui, Jingxu Xin, Xianmei Huang & Chengmin Houx. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Lyapunov-Type Inequality for Fractional order Difference Equations

Meihong Cui [α], Jingxu Xin [σ], Xianmei Huang [ρ] & Chengmin Houx [ω]

Abstract- In this paper, we obtain some Lyapunov-type inequalities for a class of fractional order difference equations with homogeneous boundary value conditions, the results of this paper are new and generalize some early results in the literature.

Keywords: *lyapunov-type inequalities, fractional order difference equations, boundary value problem.*

I. Introduction

Lyapunov inequality plays an important role both in the study of various properties of solutions of differential and difference equations such as oscillation theory, disconjugacy and eigenvalue problems and in the application in many directions of mathematics research areas. There have been many proofs and generalizations as well as improvements in this topic. A thorough literature review of continuous and discrete Lyapunov-type inequalities and their applications can be found in the survey paper [1] by Cheng. For other related references, we refer the reader to the few papers [2-9]. In 1983, Cheng [10] first obtained the following Lyapunov inequality:

$$\mathfrak{F}(b-a)\sum_{n=a}^{b-2} q(n) \geq 4, \tag{1.1}$$

where $a, b \in \mathbf{Z}, q(n)$ is a non-negative function defined on the set $\{a, a+1, \cdots, b\}$ and

$$\mathfrak{F}(m) = \begin{cases} \frac{m^2-1}{m}, & \text{if } m-1 \text{ is even}, \\ m, & \text{if } m-1 \text{ is odd}, \end{cases} \tag{1.2}$$

if the second-order difference equation

$$\Delta^2 x(n) + q(n)x(n+1) = 0, \tag{1.3}$$

has a real solution $x(n)$ such that

$$x(a) = x(b) = 0, x(n) \not\equiv 0, n \in \mathbf{Z}[a, b], \tag{1.4}$$

where and in the sequel, $a, b \in \mathbf{N}, \mathbf{Z}[a, b] = \{a, a+1, a+2, ..., b-1, b\}; c, d \in \mathbf{R}, d = c + N, N \in \mathbf{N}, [c, d]_{\mathbf{N}_c} = \{c, c+1, c+2, ..., d-1, d\}$. The constant 4 in (1.1) cannot be replaced by a larger number, either. For more discrete Lyapunov-type inequalities, we refer the reader to Cheng [11-

Author α σ ρ ω: *Department of Mathematics, Yanbian University, Yanji 133002, P. R. China. e-mail: cmhou@foxmail.com*

© 2016 Global Journals Inc. (US)

13], Clark and Hinton [14,15], Guseinov and Kaymakcalan [16], Lin and Yang [17] and Zhang and Tang [18].

In 2012, Zhang and Tang [19] considered the following even order difference equation

$$\Delta^{2k}x(n) + (-1)^{k-1}q(n)x(n+1) = 0, \tag{1.5}$$

under the following boundary conditions:

$$\Delta^{2i}x(a) = \Delta^{2i}x(b) = 0, i = 0, 1, ..., k-1; x(n) \not\equiv 0, n \in \mathbf{Z}[a, b]. \tag{1.6}$$

They obtained the following main result:

Theorem A. *Assume that $k \in \mathbf{N}$ and $q(n)$ is a real-valued function on $\mathbf{Z}$. If (1.5) has a solution $x(n) \not\equiv 0$ satisfying the boundary value conditions (1.6), then*

$$\sum_{n=a}^{b-1}[|q(n)|(n-a+1)(b-n-1)] \geq \frac{2^{3(k-1)}}{(b-a)^{2k-3}}. \tag{1.7}$$

In this paper, we will consider the following fractional order difference equation

$$\Delta^{\nu}_{\nu-2k+a}x(n) + (-1)^{k-1}q(n)x(n+\nu-2k+1) = 0, \tag{1.8}$$

where $k \in \mathbf{N}, n \in \mathbf{Z}, \nu \in \mathbf{R}^+$ satisfy $2k-1 < \nu \leq 2k$ and $q(n)$ is a real-valued function defined on $\mathbf{Z}$. In this work, we establish two discrete Lyapunov-type inequalities for (1.8) under the following boundary conditions:

$$x(a+\nu-2k) = x(b+\nu-2k) = 0, \tag{1.9}$$

$$\Delta^{\nu-2i}_{a+\nu-2k}x(n)|_{n=a} = \Delta^{\nu-2i}_{a+\nu-2k}x(n)|_{n=b} = 0, i = 1, ..., k-1; x(n+\nu-2k) \not\equiv 0, n \in \mathbf{Z}[a, b]. \tag{1.10}$$

When $\nu = 2, k = 1$, (1.8)(1.9) reduce to (1.3)(1.4). When $\nu = 2k$, (1.8)(1.9) (1.10) reduce to (1.5)(1.6).

II. Preliminaries

In this section, we collect some basic definitions and lemmas for manipulating discrete fractional operators. These can be found in the reference [20].

For any integer β, let $\mathbf{N}_\beta = \{\beta, \beta+1, \beta+2, ...\}$. We define $t^{\underline{\nu}} := \frac{\Gamma(t+1)}{\Gamma(t+1-\nu)}$, for any t and ν for which the right-hand side is defined. We also appeal to the convention that if $t+1-\nu$ is a pole of the Gamma function and $t+1$ is not a pole, then $t^{\underline{\nu}} = 0$.

Deffinition 2.1. The ν-th fractional sum of f for $\nu > 0$ is defined by

$$\Delta^{-\nu}_a f(t) = \frac{1}{\Gamma(\nu)}\sum_{s=a}^{t-\nu}(t-s-1)^{\underline{\nu-1}}f(s),$$

for $t \in \mathbf{N}_{a+\nu}$. Also, we define the trivial sum by $\Delta^{-0}_a f(t) := f(t)$, for $t \in N_a$. We also define the ν-th fractional difference for $\nu > 0$ by $\Delta^{\nu}_a f(t) := \Delta^N \Delta^{\nu-N}_a f(t)$,where $t \in \mathbf{N}_{a+N-\nu}$ and $N \in \mathbf{N}$ is chosen so that $0 \leq N-1 < \nu \leq N$.

Lemma 2.2. *Let $f : \mathbf{N}_a \longrightarrow \mathbf{R}$ be given and suppose $k \in \mathbf{N}_0$ and $\nu > 0$. Then for $t \in \mathbf{N}_{a+\nu}$,*

$$\Delta^{-\nu}_a \Delta^k f(t) = \Delta^{k-\nu}_a f(t) - \sum_{j=0}^{k-1}\frac{\Delta^j f(a)}{\Gamma(\nu-k+j+1)}(t-a)^{\underline{\nu-k+j}}.$$

© 2016 Global Journals Inc. (US)

Ref
19. Q. Zhang, X.H. Tang, *Lyapunov-type inequalities for even order difference equations*, Appl. Math. Lett. 25 (2012) 1830-1834.

Moreover, if $\mu > 0$ *with* $M-1 < \mu \leq M$, *then for* $t \in \mathbf{N}_{a+M-\mu+\nu}$,

$$\Delta_{a+M-\mu}^{-\nu}\Delta_a^{\mu} f(t) = \Delta_a^{\mu-\nu} f(t) - \sum_{j=0}^{M-1} \frac{\Delta_a^{j-M+\mu} f(a+M-\mu)}{\Gamma(\nu-M+j+1)}(t-a-M+\mu)^{\underline{\nu-M+j}}.$$

III. The Main Results

Notes

Theorem 3.1. *Let* $\nu \in (1,2]$. *Assume that* $x(n+\nu-2)$ *is a real valued function on* $\mathbf{Z}[a,b]$ *satisfy (1.8)(1.9) and* $x(n+\nu-2) \not\equiv 0$ *for* $n \in \mathbf{Z}[a,b]$. *Then,*

$$\widetilde{\mathfrak{F}}(b-a)\sum_{n=a}^{b-2}|q(n)| \geq 1, \tag{3.1}$$

where $a, b \in \mathbf{Z}$ *and*

$$\widetilde{\mathfrak{F}}(m) = \frac{1}{\Gamma(\nu)}\begin{cases} \frac{(\frac{m+1}{2}+\nu-2)^{\underline{\nu-1}}(\frac{m-1}{2}+\nu-2)^{\underline{\nu-1}}}{(m+\nu-2)^{\underline{\nu-1}}}, & \text{if } m-1 \text{ is even,} \\ \frac{((\frac{m}{2}+\nu-2)^{\underline{\nu-1}})^2}{(m+\nu-2)^{\underline{\nu-1}}}, & \text{if } m-1 \text{ is odd.} \end{cases} \tag{3.2}$$

Proof. By Lemma 2.2, we have

$$x(n) = -\Delta_a^{-\nu} q(n)x(n+\nu-1) + c_1(n-a)^{\underline{\nu-1}} + c_2(n-a)^{\underline{\nu-2}}.$$

Since $x(a+\nu-2) = x(b+\nu-2) = 0$, we get

$$x(n) = \sum_{s=a}^{b-2} G(n,s)q(s)x(s+\nu-1) = -\sum_{s=a}^{b-2} G(n,s)\Delta_{a+\nu-2}^{\nu} x(s), \tag{3.3}$$

where

$$G(n,s) = \frac{1}{\Gamma(\nu)}\begin{cases} \frac{(b+\nu-s-3)^{\underline{\nu-1}}(n-a)^{\underline{\nu-1}}}{(b+\nu-a-2)^{\underline{\nu-1}}} - (n-s-1)^{\underline{\nu-1}}, & a \leq s < n-\nu+1 \leq b-2, \\ \frac{(b+\nu-s-3)^{\underline{\nu-1}}(n-a)^{\underline{\nu-1}}}{(b+\nu-a-2)^{\underline{\nu-1}}}, & a \leq n-\nu+1 \leq s \leq b-2. \end{cases} \tag{3.4}$$

One can see that $\Delta_s G(n,s) < 0$, for $a \leq n-\nu+1 \leq s \leq b-2$ and $\Delta_s G(n,s) > 0$, for $a \leq s < n-\nu+1 \leq b-2$. Indeed, for $a \leq s < n-\nu+1 \leq b-2$, we have $\Delta_s G(n,s) = -(\nu-1)\frac{(b+\nu-s-4)^{\underline{\nu-2}}(n-a)^{\underline{\nu-1}}}{(b+\nu-a-2)^{\underline{\nu-1}}} + (\nu-1)(n-s-2)^{\underline{\nu-2}}$.

Thus, $\Delta_s G(n,s) > 0$, if and only if

$$\frac{(n-s-2)^{\underline{\nu-2}}(b+\nu-a-2)^{\underline{\nu-1}}}{(n-a)^{\underline{\nu-1}}(b+\nu-s-4)^{\underline{\nu-2}}} > 1.$$

The inequality follows from the fact that $t^{\underline{\alpha}}$ is increasing and $t^{\underline{-\alpha}}$ is decreasing if $0 < \alpha \leq 1$. Note that $G(n,b-2) > 0$ and

$$\begin{aligned} G(n,a) &= \frac{1}{\Gamma(\nu)}\left(\frac{(b+\nu-a-3)^{\underline{\nu-1}}(n-a)^{\underline{\nu-1}}}{(b+\nu-a-2)^{\underline{\nu-1}}} - (n-a-1)^{\underline{\nu-1}}\right) \\ &= \frac{1}{\Gamma(\nu)}\frac{\Gamma(n-a)}{\Gamma(n-a-\nu+1)}\left(\frac{(b-a-1)(n-a)}{(b-a+\nu-2)(n-a+1-\nu)} - 1\right) \\ &= \frac{1}{\Gamma(\nu)}\frac{\Gamma(n-a)}{\Gamma(n-a-\nu+1)}\frac{(\nu-1)(b-n+\nu-2)}{(b-a+\nu-2)(n-a+1-\nu)} > 0, \end{aligned}$$

© 2016 Global Journals Inc. (US)

we can get

$$G(n,s) > 0, \text{ for } (n,s) \in [a+\nu-1, b+\nu-3]_{\mathbf{N}_{a+\nu-1}} \times \mathbf{Z}[a, b-2].$$

Thus

$$\max_{s\in\mathbf{Z}[a,b-2]} G(n,s) = G(n, n-\nu+1), n \in [a+\nu-1, b+\nu-3]_{\mathbf{N}_{a+\nu-1}}.$$

$$|x(n)| \leq \frac{(b+2\nu-n-4)^{\underline{\nu-1}}(n-a)^{\underline{\nu-1}}}{(b+\nu-a-2)^{\underline{\nu-1}}\Gamma(\nu)} \sum_{s=a}^{b-2} |\Delta^{\nu}_{a+\nu-2} x(s)|, n \in [a+\nu-1, b+\nu-3]_{\mathbf{N}_{a+\nu-1}}$$

and

$$|x(n+\nu-1)| \leq \frac{(b+\nu-n-3)^{\underline{\nu-1}}(n+\nu-a-1)^{\underline{\nu-1}}}{(b+\nu-a-2)^{\underline{\nu-1}}\Gamma(\nu)} \sum_{s=a}^{b-2} |\Delta^{\nu}_{a+\nu-2} x(s)|, n \in \mathbf{Z}[a, b-2]. \quad (3.5)$$

Next, we denote $g(n) = (b+\nu-n-3)^{\underline{\nu-1}}(n+\nu-a-1)^{\underline{\nu-1}}$, then by computing, we have

$$\Delta g(n) = \frac{\Gamma(b+\nu-n-3)\Gamma(n+\nu-a)}{\Gamma(b-n-1)\Gamma(n-a+2)}(\nu-1)(b-2n+a-3).$$

If $b-a-1$ is even, let $\Delta g(n) = 0$, then $n = \frac{b-a-1}{2} + a - 1$.

$$\max_{n\in\mathbf{Z}[a,b-2]} g(n) = g(\frac{b-a-1}{2} + a - 1) = (\frac{b-a+1+\nu-2}{2})^{\underline{\nu-1}}(\frac{b-a-1+\nu-2}{2})^{\underline{\nu-1}}. \quad (3.6)$$

If $b-a-1$ is odd, we obtain $\Delta g(n) > 0$, for $n < \frac{b-a-1}{2} + a - 1$; $\Delta g(n) < 0$, for $n > \frac{b-a-1}{2} + a - 1$. Therefore,

$$\max_{n\in\mathbf{Z}[a,b-2]} g(n) = \max\{g(\frac{b-a}{2} + a - 1), g(\frac{b-a}{2} + a - 2)\}.$$

Since

$$g(\frac{b-a}{2} + a - 1) = g(\frac{b+a}{2} - 1) = ((\frac{b-a}{2} + \nu - 2)^{\underline{\nu-1}})^2,$$

$$g(\frac{b-a}{2} + a - 2) = g(\frac{b+a}{2} - 2) = (\frac{b-a}{2} + \nu - 3)^{\underline{\nu-1}}(\frac{b-a}{2} + \nu - 1)^{\underline{\nu-1}},$$

$$\begin{aligned} g(\frac{b-a}{2} + a - 1) - g(\frac{b-a}{2} + a - 2) &= ((x+\nu-2)^{\underline{\nu-1}})^2 - (x+\nu-3)^{\underline{\nu-1}}(x+\nu-1)^{\underline{\nu-1}} \\ &= \frac{(x+\nu-2)\Gamma^2(x+\nu-2)}{x(x-1)^2\Gamma(x-1)}(\nu-1) > 0, \end{aligned}$$

where, $x = \frac{b-a}{2}$, we see that

$$\max_{n\in\mathbf{Z}[a,b-2]} g(n) = g(\frac{b-a}{2} + a - 1) = \big((\frac{b-a}{2} + \nu - 2)^{\underline{\nu-1}}\big)^2. \quad (3.7)$$

Note that (3.5) and by (3.6)(3.7), we obtain

$$|x(n+\nu-1)| \leq \widetilde{\mathfrak{F}}(b-a) \sum_{s=a}^{b-2} |\Delta^{\nu}_{a} x(s)| = \widetilde{\mathfrak{F}}(b-a) \sum_{s=a}^{b-2} |q(s)||x(s+\nu-1)|.$$

Notes

© 2016 Global Journals Inc. (US)

Furthermore,

$$\sum_{n=a}^{b-2}|q(n)||x(n+\nu-1)| \leq \widetilde{\mathfrak{F}}(b-a)\sum_{s=a}^{b-2}|q(n)|\sum_{s=a}^{b-2}|q(s)||x(s+\nu-1)|,$$

it shows that (3.1) holds.

Remark 3.2. The inequality (3.1) is sharp. Indeed, since

Notes

$$\begin{aligned}
\Delta\widetilde{\mathfrak{F}}(m) &= \frac{1}{\Gamma(\nu)}\left(\frac{((\frac{m+1}{2}+\nu-2)^{\underline{\nu-1}})^2}{(m+\nu-1)^{\underline{\nu-1}}}-\frac{(\frac{m+1}{2}+\nu-2)^{\underline{\nu-1}}(\frac{m-1}{2}+\nu-2)^{\underline{\nu-1}}}{(m+\nu-2)^{\underline{\nu-1}}}\right)\\
&= \frac{(\frac{m+1}{2}+\nu-2)^{\underline{\nu-1}}}{\Gamma(\nu)(m+\nu-2)^{\underline{\nu-1}}(m+\nu-1)^{\underline{\nu-1}}}\\
&\quad\times\left((\frac{m+1}{2}+\nu-2)^{\underline{\nu-1}}(m+\nu-2)^{\underline{\nu-1}}-(\frac{m-1}{2}+\nu-2)^{\underline{\nu-1}}(m+\nu-1)^{\underline{\nu-1}}\right)\\
&= \frac{(\frac{m+1}{2}+\nu-2)^{\underline{\nu-1}}}{\Gamma(\nu)(m+\nu-2)^{\underline{\nu-1}}(m+\nu-1)^{\underline{\nu-1}}}\\
&\quad\times\left(\frac{\Gamma(\frac{m+1}{2}+\nu-1)\Gamma(m+\nu-1)}{\Gamma(\frac{m+1}{2})\Gamma(m)}-\frac{\Gamma(\frac{m-1}{2}+\nu-1)\Gamma(m+\nu)}{\Gamma(\frac{m-1}{2})\Gamma(m+1)}\right)\\
&= \frac{(\frac{m+1}{2}+\nu-2)^{\underline{\nu-1}}}{\Gamma(\nu)(m+\nu-2)^{\underline{\nu-1}}(m+\nu-1)^{\underline{\nu-1}}}\frac{\Gamma(\frac{m-1}{2}+\nu-1)\Gamma(m+\nu-1)}{\Gamma(\frac{m-1}{2})\Gamma(m)}(\frac{2(\nu-1)}{m-1}-\frac{\nu-1}{m})>0
\end{aligned}$$

for $m-1$ is even,

$$\begin{aligned}
\Delta\widetilde{\mathfrak{F}}(m) &= \frac{1}{\Gamma(\nu)}\left(\frac{(\frac{m+2}{2}+\nu-2)^{\underline{\nu-1}}(\frac{m}{2}+\nu-2)^{\underline{\nu-1}}}{(m+\nu-1)^{\underline{\nu-1}}}-\frac{((\frac{m}{2}+\nu-2)^{\underline{\nu-1}})^2}{(m+\nu-2)^{\underline{\nu-1}}}\right)\\
&= \frac{(\frac{m}{2}+\nu-2)^{\underline{\nu-1}}}{\Gamma(\nu)(m+\nu-2)^{\underline{\nu-1}}(m+\nu-1)^{\underline{\nu-1}}}\\
&\quad\times\left((\frac{m+2}{2}+\nu-2)^{\underline{\nu-1}}(m+\nu-2)^{\underline{\nu-1}}-(\frac{m}{2}+\nu-2)^{\underline{\nu-1}}(m+\nu-1)^{\underline{\nu-1}}\right)\\
&= \frac{(\frac{m}{2}+\nu-2)^{\underline{\nu-1}}}{\Gamma(\nu)(m+\nu-2)^{\underline{\nu-1}}(m+\nu-1)^{\underline{\nu-1}}}\left(\frac{\Gamma(\frac{m+2}{2}+\nu-1)\Gamma(m+\nu-1)}{\Gamma(\frac{m+2}{2})\Gamma(m)}-\frac{\Gamma(\frac{m}{2}+\nu-1)\Gamma(m+\nu)}{\Gamma(\frac{m}{2})\Gamma(m+1)}\right)\\
&= \frac{(\frac{m+1}{2}+\nu-2)^{\underline{\nu-1}}}{\Gamma(\nu)(m+\nu-2)^{\underline{\nu-1}}(m+\nu-1)^{\underline{\nu-1}}}\frac{\Gamma(\frac{m}{2}+\nu-1)\Gamma(m+\nu-1)}{\Gamma(\frac{m}{2})\Gamma(m)}\frac{\nu-1}{m}>0
\end{aligned}$$

for $m-1$ is odd. We see that $\widetilde{\mathfrak{F}}$ is a strictly increasing function of m. If

$$\widetilde{\mathfrak{F}}(b-a)\sum_{n=a}^{b-2}|q(n)|<1,$$

then (1.8) cannot have a nontrivial solution $x(n+\nu-1)$ defined on the set $\mathbf{Z}[c,d]$ which satisfies $x(c+\nu-2)=x(d+\nu-2)=0$, where $a-1\leq c-1<d+1\leq b+1, \nu\in(1,2]$. Otherwise,

$$\sum_{n=a}^{b-2}|q(n)|\geq\sum_{n=c}^{d-2}|q(n)|\geq\frac{1}{\widetilde{\mathfrak{F}}(d-c)}\geq\frac{1}{\widetilde{\mathfrak{F}}(b-a)}>\sum_{n=a}^{b-2}|q(n)|$$

which is a contradiction.

© 2016 Global Journals Inc. (US)

Remark 3.3. In the Theorem 3.1, if $\nu = 2$ and $q(n)$ is a non-negative function defined on the set $\mathbf{Z}[a, b]$, then $\widetilde{\mathfrak{F}}(m) = \mathfrak{F}(m)$, i.e., (1.1) holds. Therefore Theorem 3.1 contains the result of reference [10].

Theorem 3.4. *Let $\nu \in (1, 2]$. Assume that $x(n + \nu - 2)$ is a real valued function on $\mathbf{Z}[a, b]$ satisfy (1.8)(1.9) and $x(n + \nu - 2) \not\equiv 0$ for $\mathbf{Z}[a, b]$. Then,*

$$|x(n+\nu-1)| \leq \frac{(b+\nu-n-3)^{\underline{\nu-1}}(n+\nu-a-1)^{\underline{\nu-1}}}{(b+\nu-a-2)^{\underline{\nu-1}}\Gamma(\nu)} \sum_{s=a}^{b-2} |\Delta^{\nu}_{a+\nu-2} x(s)|, \tag{3.8}$$

$$\sum_{n=a}^{b-2} |x(n+\nu-1)| \leq \sum_{n=a}^{b-2} \frac{(b+\nu-n-3)^{\underline{\nu-1}})(n+\nu-a-1)^{\underline{\nu-1}}}{(b+\nu-a-2)^{\underline{\nu-1}}\Gamma(\nu)} \sum_{s=a}^{b-2} |\Delta^{\nu}_{a+\nu-2} x(s)|, \tag{3.9}$$

and

$$\sum_{n=a}^{b-2} |x(n+\nu-1)|^2 \leq \sum_{n=a}^{b-2} \sum_{s=a}^{b-2} [G(n+\nu-1, s)]^2 \sum_{s=a}^{b-2} |\Delta^{\nu}_{a+\nu-2} x(s)|^2. \tag{3.10}$$

Proof. By (3.5), we can immediately get (3.8) and (3.9). Using Cauchy inequality and by (3.3), we may easily see that (3.10) holds.

Corollary 3.5. *Let $\nu = 2$. Assume that $x(n)$ is a real valued function on $\mathbf{Z}[a, b]$ satisfy (1.3)(1.4) and $x(n) \not\equiv 0$ for $\mathbf{Z}[a, b]$. Then,*

$$|x(n)| \leq \frac{(b-n)(n-a)}{(b-a)} \sum_{s=a}^{b-2} |\Delta^2 x(s)|, \tag{3.11}$$

$$\sum_{n=a}^{b-2} |x(n)| \leq \sum_{n=a}^{b-2} \sum_{s=a}^{b-2} G(n, s) |\Delta^2 x(s)| \leq \frac{(b-a)^2}{8} \sum_{s=a}^{b-2} |\Delta^2 x(s)|. \tag{3.12}$$

Proof.

Since

$$G(n, s) = \frac{1}{b-a} \begin{cases} (s+1-a)(b-n), & a \leq s < n-1 \leq b-2, \\ (b-s-1)(n-a), & a \leq n-1 \leq s \leq b-2, \end{cases}$$

it follows from above form that

$$|x(n)| \leq \sum_{s=a}^{b-2} G(n, s) |\Delta^2 x(s)| \leq \frac{(n-a)(b-n)}{b-a} \sum_{s=a}^{b-2} |\Delta^2 x(s)|.$$

Furthermore

$$\begin{aligned} \sum_{n=a}^{b-2} |x(n)| &\leq \sum_{n=a}^{b-2} \sum_{s=a}^{b-2} G(n, s) |\Delta^2 x(s)| \\ &\leq \sum_{s=a}^{b-2} \sum_{n=a}^{b-1} G(n, s) |\Delta^2 x(s)| \\ &\leq \sum_{s=a}^{b-2} \Big(\sum_{n=a}^{s+1} G(n, s) + \sum_{n=s+2}^{b-1} G(n, s) \Big) |\Delta^2 x(s)| \\ &= \frac{1}{b-a} \sum_{s=a}^{b-2} \big((b-s-1)(1+2+\cdots+(s-a+1)) \end{aligned}$$

© 2016 Global Journals Inc. (US)

Ref
10. S. S. Cheng, *A discrete analogue of the inequality of Lyapunov*, Hokkaido Math. J. 12 (1983) 105-112.

$$
\begin{aligned}
&+(s+1-a)(1+2+\cdots+(b-s-2)))|\Delta^2 x(s)| \\
= \;& \frac{1}{b-a}\sum_{s=a}^{b-2}((b-s-1)(1+2+\cdots+s-a) \\
&+(s+1-a)(1+2+\cdots+b-s-1))|\Delta^2 x(s)| \\
= \;& \frac{1}{2}\sum_{s=a}^{b-2}(b-s-1)(s+1-a)|\Delta^2 x(s)| \\
\leq \;& \frac{(b-a)^2}{8}\sum_{s=a}^{b-2}|\Delta^2 x(s)|.
\end{aligned}
$$

Theorem 3.6. *Let $\nu \in (2k-1, 2k]$. Assume that $x(n+\nu-2)$ is a real valued function on $\mathbf{Z}[a,b]$ satisfy (1.8)-(1.10) and $x(n+\nu-2) \not\equiv 0$ for $n \in \mathbf{Z}[a,b]$. Then,*

$$
\sum_{n=a}^{b-2}|q(n)|(b+\nu-2k-n-1)^{\underline{\nu-2k+1}}(n+\nu-2k+1-a)^{\underline{\nu-2k+1}} \geq (\frac{2^{3(k-1)}}{(b-a)^2})^{k-1}(b+\nu-2k-a)^{\underline{\nu-2k+1}}. \tag{3.13}
$$

Proof. Assume that $x(n+\nu-2)$ is a real valued function on $\mathbf{Z}[a,b]$ satisfy (1.8)-(1.10) and $x(n+\nu-2) \not\equiv 0$ for $n \in \mathbf{Z}[a,b]$. Since conditions (1.10), by Corollary 3.3, we have

$$
\begin{aligned}
\sum_{n=a}^{b-2}|\Delta_{\nu-2k+a}^{\nu-2(k-1)}x(s)| &\leq \frac{(b-a)^2}{8}\sum_{n=a}^{b-2}|\Delta_{\nu-2k+a}^{\nu-2(k-1)+2}x(s)| \\
&\leq (\frac{(b-a)^2}{8})^2\sum_{n=a}^{b-2}|\Delta_{\nu-2k+a}^{\nu-2(k-3)}x(s)| \\
&\leq \cdots \\
&\leq (\frac{(b-a)^2}{8})^{k-1}\sum_{n=a}^{b-2}|\Delta_{\nu-2k+a}^{\nu}x(s)|. \qquad (3.14)
\end{aligned}
$$

Denote $\beta = \nu - 2(k-1)$, then $\beta \in (1,2], \Delta_{\nu-2k+a}^{\nu-2(k-1)}x(n) = \Delta_{\beta-2+a}^{\beta}x(n)$. By Theorem 3.4, we have

$$
|x(n+\beta-1)| \leq \frac{(b+\beta-s-3)^{\underline{\beta-1}}(n+\beta-a-1)^{\underline{\beta-1}}}{(b+\beta-a-2)^{\underline{\beta-1}}}\sum_{s=a}^{b-2}|\Delta_{a+\beta-2}^{\beta}x(s)|,
$$

combine (3.14), we get

$$
\begin{aligned}
|x(n+\nu-2k+1)| &\leq (\frac{(b-a)^2}{8})^{k-1}\frac{(b+\beta-s-3)^{\underline{\beta-1}}(n+\beta-a-1)^{\underline{\beta-1}}}{(b+\beta-a-2)^{\underline{\beta-1}}}\sum_{s=a}^{b-2}|\Delta_{a+\nu-2k}^{\nu}x(s)| \\
&= (\frac{(b-a)^2}{8})^{k-1}\frac{(b+\nu-2k-n-1)^{\underline{\nu-2k+1}}(n+\nu-2k-a+1)^{\underline{\nu-2k+1}}}{(b+\nu-2k-a)^{\underline{\nu-2k+1}}} \\
&\quad \times\sum_{s=a}^{b-2}|q(s)x(s+\nu-2k+1)|.
\end{aligned}
$$

Global Journal of Science Frontier Research (F) Volume XVI Issue I Version I Year 2016

© 2016 Global Journals Inc. (US)

$$\sum_{n=a}^{b-2} q(n)|x(n+\nu-2k+1)| \leq \sum_{n=a}^{b-2} q(n)\left(\frac{(b-a)^2}{8}\right)^{k-1} \frac{(b+\nu-2k-n-1)^{\underline{\nu-2k+1}}(n+\nu-2k-a+1)^{\underline{\nu-2k+1}}}{(b+\nu-2k-a)^{\underline{\nu-2k+1}}}$$

$$\times \sum_{s=a}^{b-2} |q(s)x(s+\nu-2k+1)|.$$

it follows that (3.13) holds.

Corollary 3.7. *Let $\nu = 2k$. Assume that $x(n)$ is a real valued function on $\mathbf{Z}[a,b]$ satisfy (1.8)-(1.10) and $x(n) \not\equiv 0$ for $\mathbf{Z}[a,b]$. Then,*

$$\sum_{n=a}^{b-2} |q(n)|(b-n-1)(n+1-a) \geq \frac{2^{3(k-1)}}{(b-a)^{2k-3}}.$$

Remark 3.8. The Corollary 3.7 is the Theorem A. Therefore our results contains the main results of reference [19].

References Références Referencias

1. S.S. Cheng, *Lyapunov inequalities for differential and difference equations*, Fasc. Math. 23(1991) 25-41.
2. A.M. Liapunov, *Problme gnral de la stabilit du mouvement*, Ann. Fac. Sci. Univ. Toulouse 2 (1907) 203-407.
3. D. Cakmak, *Lyapunov-type integral inequalities for certain higher order differential equations,* Appl. Math. Comput. 216 (2010) 368-373.
4. X. He, X.H. Tang, *Lyapunov-type inequalities for even order differential equations,* Commun Pure Appl. Anal. 11 (2) (2012) 465-473.
5. L.Q. Jiang, Z. Zhou, *Lyapunov inequality for linear Hamiltonian systems on time scales,* J. Math. Anal. Appl. 310 (2005) 579-593.
6. X. Wang, *Stability criteria for linear periodic Hamiltonian systems,* J. Math. Anal. Appl. 367 (2010) 329-336.
7. X. Yang, *On Liapunov-type inequality for certain higher-order differential equations,* Appl. Math. Comput. 134 (2-3) (2003) 307-317.
8. X. Yang, *On inequalities of Lyapunov type,* Appl. Math. Comput. 134 (2-3) (2003) 293-300.
9. X. Yang, K. Lo, *Lyapunov-type inequality for a class of even-order differential equations,* Appl. Math. Comput. 215 (2010) 3884-3890.
10. S. S. Cheng, *A discrete analogue of the inequality of Lyapunov,* Hokkaido Math. J. 12 (1983) 105-112.
11. S.S. Cheng, L.Y. Hsieh, *On discrete analogue of Lyapunovs inequality: addendum,* Tamkang J. Math. 20 (1989) 333-340.
12. S.S. Cheng, *A sharp condition for the ground state of difference equation,* Appl. Anal. 34 (1989) 105-109.
13. S.S. Cheng, L.Y. Hsieh, D. Chao, *Discrete Lyapunov inequality conditions for partial difference equations,* Hokkaido Math. J. 19 (1990) 229-239.
14. S. Clark, D.B. Hinton, *Discrete Lyapunov inequalities for linear Hamiltonian systems,* Math. Inequal. Appl. 1 (1998) 201-209.
15. S. Clark, D.B. Hinton, *Discrete Lyapunov inequalities,* Dynam. Systems Appl. 8 (1999) 369-380.
16. G.Sh. Guseinov, B. Kaymakcalan, *Lyapunov inequalities for discrete linear Hamiltonian systems,* Comput. Math. Appl. 45 (2003) 1399-1416.

Ref

19. Q. Zhang, X.H. Tang, *Lyapunov-type inequalities for even order difference equations,* Appl. Math. Lett. 25 (2012) 1830-1834.

© 2016 Global Journals Inc. (US)

17. S.H. Lin, G.S. Yang, *On discrete analogue of Lyapunovs inequality,* Tamkang J. Math. 20(1989) 169-186.
18. Q. Zhang, X.H. Tang, *Lyapunov inequalities and stability for discrete linear Hamiltonian system,* Appl. Math. Comput. 218 (2011) 574-582.
19. Q. Zhang, X.H. Tang, *Lyapunov-type inequalities for even order difference equations,* Appl. Math. Lett. 25 (2012) 1830-1834.
20. M.T. Holm, *The Theory of Discrete Fractional Calculus: Development and Application,* DigitalCommons@University of Nebraska-Lincoln, 2011.
21. M. Unal, D. Cakmak, A. Tiryaki, *A discrete analogue of Lyapunov-type inequalities for nonlinear systems,* Comput. Math. Appl. 55 (2008) 2631-2642.

Notes

© 2016 Global Journals Inc. (US)

This page is intentionally left blank

© 2016 Global Journals Inc. (US)

Global Journal of Science Frontier Research: F
Mathematics and Decision Sciences
Volume 16 Issue 1 Version 1.0 Year 2016
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

A Construction of Non-Gaussian White Noise Analysis using the Theory of Hypercomplex Systems

By A. S. Okb El Bab, Hossam A. Ghany & M. Zakarya
Al-Azhar University, Egypt

Abstract- In this paper, we present a generalization of white noise analysis to the case of non-Gaussian measures. For this purpose, we use a biorthogonal approach in which instead of the exponentials the characters of commutative hypercomplex systems are employed. Moreover, we construct the elements of Wick calculus in a non-Gaussian setting.

Keywords: *white noise analysis, biorthogonal approach, hypercomplex systems, wick calculus.*

GJSFR-F Classification : *MSC 2010: 30G35*



Strictly as per the compliance and regulations of :

© 2016. A. S. Okb El Bab, Hossam A. Ghany & M. Zakarya. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

eJournal
physical existance Print Journal
epaper
open standard
3d virtual journal

25. T. Hida, *Analysis of Brownian functionals*, in Carleton Math. Lecture Notes **13**, Carleton Univ., Ottawa 1975.

A Construction of Non-Gaussian White Noise Analysis using the Theory of Hypercomplex Systems

A. S. Okb El Bab [α], Hossam A. Ghany [σ] & M. Zakarya [ρ]

Abstract- In this paper, we present a generalization of white noise analysis to the case of non-Gaussian measures. For this purpose, we use a biorthogonal approach in which instead of the exponentials the characters of commutative hypercomplex systems are employed. Moreover, we construct the elements of Wick calculus in a non-Gaussian setting.

Keywords: white noise analysis, biorthogonal approach, hypercomplex systems, wick calculus.

I. Introduction

White noise analysis appeared in 1975 after the work of T. Hida [25] and then has been actively developed in a series of articles and books. For a sufficiently full bibliography see [26]. As mentioned in [4], Gaussian white noise analysis can be thought of as a theory of generalized functions of infinitely many variables whose pairing with test functions from special spaces is given via the integration by the Gaussian measure. It is well known that there are several approaches to the construction of such theory of generalized functions: the Berezansky-Samoilenko approach [3] and the Hida approach [25]. In the Berezansky-Samoilenko approach, the spaces of test and generalized functions are constructed as infinite tensor products of one-dimensional spaces. The Hida approach consists in a construction of some rigging of a Fock space with subsequent application of the Wiener-Itô-Segal isomorphism to the spaces of this rigging. In most cases, investigations in white noise analysis and its generalizations are based on the Hida approach. So, the Hida approach is more convenient than Berezansky-Samoilenko approach. Recently, some authors as Okb El Bab, Zabel, Ghany and Hyder [15], Ghany [16], Ghany and Hyder [17–19], Ghany and Zakarya [20–22] and Ghany and Qurashi [23], studied some important subjects related to Gaussian white noise analysis. There exist many works investigated to white noise analysis development Works deal with the construction of spaces of test and generalized functions and operators acting in these spaces using the Wiener-Itô-Segal isomorphism and various riggings of the Fock space. For more details, see [4, 25, 28]. Works deal with the so-called Jacobi fields approach to a generalization of white noise analysis [6, 7]. In these works, the role of the Wiener-Itô-Segal isomorphism is

Author α: Department of Mathematics Faculty of Science Al-Azhar University, Naser City, Cairo, Egypt.
e-mail: ahmedokbelbab@yahoo.com
Author σ: Department of Mathematics, Faculty of Science Taif University, Taif, Saudi Arabia, Department of Mathematics, Helwan University, Cairo, Egypt. e-mail: h.abdelghany@yahoo.com
Author ρ: Department of Mathematics, Faculty of Science, Al-Azhar University, Assiut 71524, Egypt.
e-mail: mohammed zakarya@rocketmail.com

© 2016 Global Journals Inc. (US)

played as a unitary Fourier transform which is defined by the Jacobi field, i.e., by some family of commuting selfadjoint operators that act in the Fock space and have a Jacobi structure. Works use the biorthogonal approach to a generalization of white noise analysis. In this approach, one replaces the system of Hermite polynomials, that are orthogonal with respect to the Gaussian measure, with a certain biorthogonal system. The biorthogonal approach was inspired by [12], proposed in [1] and developed in [2,8,38]. Note that in [8] was first observed that the biorthogonal approach is deep connected with the theory of hypercomplex systems.

Hypercomplex systems date back to J. Delsarte and B. M. Levitan work during the 1930s and 1940s, but the substantial development had to wait till the 1990s when Berezansky and Kondratiev put hypercomplex systems in the right setting for harmonic analysis [9]. Recently, some authors as Okb El Bab, Zabel and Ghany [37, 40, 41] studied some important subjects in hypercomplex systems.

Generalized translation operators were first introduced by Delsarte [13] as an object that generalizes the idea of translation on a group. They were systematically studied by Levitan [31–35], for some classes of generalized translation operators, he obtained generalizations of harmonic analysis, the Lie theory, the theory of almost periodic functions, the theory of group representations, etc. In fact, each hypercomplex system can be associated with a family of generalized translation operators. This correspondence can be found in [5].

This paper focuses on the connection between white noise analysis and hypercomplex systems. Precisely, we produce a generalization of white noise analysis to the case of non-Gaussian measures. For this aim, we propose a biorthogonal approach in which instead of the exponentials the characters of commutative hypercomplex systems are used. Furthermore, we construct the elements of Wick calculus, namely, we introduce a new Wick product with respect to non-Gaussian measures, the associated Hermite transform and the characterization theorem for the constructed spaces of generalized functions. In Gaussian white noise analysis such calculus has found numerous applications, in particular, in fluid mechanics and financial mathematics, see e.g. [14, 27] for more details.

This paper is organized as follows: Section 2 is devoted to provide the basic topics of hypercomplex systems with locally compact basis. In Section 3, we construct spaces of test and generalized functions by means of generalized Delsarte characters. In Section 4, we introduce the elements of Wick calculus.

II. Hypercomplex Systems with Locally Compact Basis

Let Q be a complete separable locally compact metric space of points $p, q, r, ...; \mathcal{B}(Q)$ is the σ-algebra of Borel subsets, and $\mathcal{B}_0(Q)$ is the subring of $\mathcal{B}(Q)$, which consists of sets with compact closure. We shall consider the Borel measures; i.e., positive regular measures on $\mathcal{B}(Q)$, finite on compact sets. We denote by $C(Q)$ the space of continuous functions on $Q, C_b(Q), C_\infty(Q)$ and $C_0(Q)$ consists respectively of bounded, tending to zero at infinity and compactly supported functions from $C(Q)$.

A hypercomplex system with the basis Q is defined by its structure measure $c(A, B, r)(A, B \in B(Q); r \in Q)$. A structure measure $c(A, B, r)$ is a Borel measure in A (respectively B) if we fix B, r (respectively A, r) which satisfies the following properties:

(I) $\forall A, B \in \mathcal{B}_0(Q)$, the function $c(A, B, r) \in C_0(Q)$,

(II) $\forall A, B \in \mathcal{B}_0(Q)$ and $s, r \in Q$, the following associativity relation holds

Ref

13. J. Delsarte, Sur certaines transformation fonctionelles relative aux equations linearies aux derivees partiels du seconde ordre, *C. R. Acad. Sci., Ser.* **17** (1938), 178-182.

© 2016 Global Journals Inc. (US)

$$\int_Q c(A,B,r)d_r c(E_r,C,s) = \int_Q c(B,C,r)d_r c(A,E_r,s), \quad C \in \mathcal{B}(Q). \tag{2.1}$$

(III) The structure measure is said to be commutative if

$$c(A,B,r) = c(B,A,r), \quad (A,B \in \mathcal{B}_0(Q)). \tag{2.2}$$

A measure m is said to be a multiplicative measure if

$$\int_Q c(A,B,r)dm(r) = m(A)m(B); \quad A,B \in \mathcal{B}_0(Q). \tag{2.3}$$

(IV) We will suppose the existence of a multiplicative measure. Under certain restrictions imposed on the commutative structure measure, multiplicative measure exists. (See [30]).

Consider the space $L_1(Q,dm(x))$ of functions on Q with respect to the multiplicative measure m. For any $\Phi,\Psi \in L_1(Q,dm(x))$, the convolution

$$\begin{aligned}(\Phi * \Psi)(r) &= \int_Q \Phi(p)dp \int_Q \Psi(q)dqc(E_p,E_q,r) \\ &= \int_Q \int_Q \Phi(p)\Psi(q)c(p,q,r)dm(p)dm(q) \\ &= \int_Q \int_Q \Phi(p)\Psi(q)dm_r(p,q) \end{aligned} \tag{2.4}$$

is well defined. (See [5]).

The space $L_1(Q,dm(x))$ with the convolution (2.4) is a Banach algebra which is commutative if (**III**) holds. This Banach algebra is called the hypercomplex system with the basis Q.

It is obvious that $c(A,B,r) = (K_A * K_B)(r); A,B \in \mathcal{B}_0(Q)$ and K_A is the characteristic function of the set A.

A hypercomplex system may or may not have a unity. If a unity not included in $L_1(Q,dm(x))$, then it is convenient to join it formally to $L_1(Q,dm(x))$.

A non zero measurable and bounded almost everywhere function $Q \ni r \mapsto \chi(r) \in \mathbb{C}$ is said to be a character of the hypercomplex system $L_1(Q,dm(x))$ if $\forall A,B \in \mathcal{B}_0(Q)$

$$\int_Q c(A,B,r)\chi(r)dm(r) = \chi(A)\chi(B), \tag{2.5}$$

$$\int_C \chi(r)dm(r) = \chi(C), \quad C \in \mathcal{B}_0(Q). \tag{2.6}$$

(V) A hypercomplex system is said to be normal, if there exists an involution homomorphism $Q \ni r \mapsto r^* \in Q$ such that $m(A) = m(A^*)$ and $c(A,B,C) = c(C,B^*,A), c(A,B,C) = c(A^*,C,B), (A,B \in \mathcal{B}_0(Q))$, where

$$c(A,B,C) = \int_C c(A,B,r)dm(r) \tag{2.7}$$

Ref

30. A. A. Kalyuzhnyi, A theorem on the existence of multiplicative measure, *Ukr. Math. Zh.*, **35**(1983), 369-371.

© 2016 Global Journals Inc. (US)

(VI) A normal hypercomplex system possesses a basis unity if there exists a point $e \in Q$ such that $e^* = e$ and

$$c(A, B, e) = m(A^* \cap B), \quad A, B \in \mathcal{B}(Q). \tag{2.8}$$

We should remark that, for a normal hypercomplex system, the mapping

$$L_1(Q, dm(x)) \ni \Phi(r) \mapsto \Phi^*(r) \in L_1(Q, dm(x))$$

is an involution in the Banach algebra $L_1(Q, dm(x))$, the multiplicative measure is unique and the characters of such a system are continuous. (See [5]). A character χ of a normal hypercomplex system is said to be Hermitian if

$$\chi(r^*) = \overline{\chi(r)}, \quad (r \in Q), \tag{2.9}$$

and ordinary if it is bounded. The function $\chi(r) = 1, r \in Q$ is always a character; for all χ, $\chi(e) = 1$.

Denote the set of all bounded Hermitian characters by X_h, i.e.,

$$X_h = \{\chi \in C_b(Q) : \chi \neq 0, \int_Q c(A, B, r)\chi(r)dm(r) = \chi(A)\chi(B), \overline{\chi(r)} = \chi(r^*)\}. \tag{2.10}$$

Let $L_1(Q, dm(x))$ be a hypercomplex system with basis Q and let Ω be a space of complex valued functions on Q. Assume that an operator valued function $Q \ni p \mapsto L_p : \Omega \to \Omega$ is given such that the function $g(p) = (L_p f)(q)$ belongs to Ω for any $f \in \Omega$ and any fixed $q \in Q$. For such hypercomplex system it is possible to introduce generalized translation operators L_p:

$$Q \times Q \ni \langle p, q \rangle \mapsto (L_p\Phi)(q) \in \mathbb{C}, \quad \Phi \in C(Q), \tag{2.11}$$

where $(L_p\Phi)(q) = \Phi(q + p)$ in case of the usual translation on $Q = \mathbb{R}$. We will suppose, in addition, that this function is separately continuous. By using the operators L_p one can rewrite the convolution (2.4) as follows:

$$\big(\Phi * \Psi\big)(p) = \int_Q \Phi(q)\big(L_{q^*}\Psi\big)(p)\, dm(q) \quad \Phi, \Psi \in L_1(Q, dm(x)). \tag{2.12}$$

A generalized character of a hypercomplex system $L_1(Q, dm(x))$ is defined to be a function $\chi \in C(Q)$ for which

$$(L_p\chi)(q) = \chi(p)\chi(q), \qquad p, q \in Q. \tag{2.13}$$

Now, we give Some examples to illustrative the concept of hypercomplex systems

Example 2.1. Let $Q = G$ be commutative, locally compact group. with unite e. Consider G is groups algebra, i.e., a set $L_1(G, m)$ of functions defined on the groups G with respect to the measure m is denoted by Haar measure $dm(p)$. Then, we can define the involution with the form

$$G \ni p \longmapsto p^* = p^{-1}, \tag{2.14}$$

where

$$\big(L_p\Phi\big)(q) = \Phi(pq), \quad (p, q \in G), \tag{2.15}$$

Ref

5. Yu. M. Berezansky and A. A. Kalyuzhnyi, *Harmonic Analysis in Hypercomplex Systems*, Naukova Dumka,Kyiv, Ukrania, 1992. (in Russian; English transl.: Kluwer: Dordrecht-Boston-London, 1996).

© 2016 Global Journals Inc. (US)

then the convolution in (2.12) becomes in the form

$$\big(\Phi * \Psi\big)(p) = \int_Q \Phi(pq)\Psi(q^*)d(q), \tag{2.16}$$

and we can define the structure measure as the following

$$C\big(A, B, r\big) = m\big(A^{-1}r \cap B\big), \tag{2.17}$$

where $A \in \mathcal{B}(G), r \in G$. Thus, we obtain a commutative hypercomplex System $L_1(G, m)$ with basis unity e.

Example 2.2. Consider the sturm-Liouvvlle equation in the quadratic axis $Q = \mathbb{R}_+ = [0, \infty)$:

$$(Mu)(x) = -u'' + g(x)u = \lambda u, \tag{2.18}$$

where $g \in C(\mathbb{R}_+)$ is a non-negative and non-increasing potential. connect with this equation the following hyperbolic equation on $\mathbb{R}^2 : M_{(x)}u = M_{(y)}v$, i.e.,

$$-\frac{\partial^2 v}{\partial x^2} + g(x)v = -\frac{\partial^2 v}{\partial y^2} + g(y)v, \quad v = v(x, y), \tag{2.19}$$

where $x, y \in \mathbb{R}$, g is the even extension of the potential to the whole axis. Equation (2.19) gives rise to a generalized translation operator $L_{(x)}$, $x \in \mathbb{R}_+$: by definition

$$\big(L_{(x)}f\big)(y) = \frac{v\big(x, y\big)}{v(x)v(y)}, \quad x, y \in \mathbb{R}_+, \tag{2.20}$$

where $v(x, y)$ is the solution of the Cauchy problem for (2.19) with initial conditions $v(x, 0) = f(x)v(x)$, $\frac{\partial v}{\partial y}(x, 0) = 0$, and $v(x) = \phi(x, 0)$, where $\phi(x, \lambda)$ is the solution of the following problem on $\mathbb{R}_+ : M\phi = \lambda\phi$, $\phi(0, \lambda) = 1$, $\phi'(0, \lambda) = 0$ $\big($both functions $f(x)$ and $v(x)$ being evenly extended to $\mathbb{R}\big)$. Now, for any $x \in \mathbb{R}_+$, $x^* = x, dm(x) = v^2(x)d(x)$, and $\chi(x) = \phi\big(x, \lambda\big)$, where $\lambda \in \mathbb{C}$ is arbitrary, the convolution for hypercomplex system with basis $Q = \mathbb{R}_+$ is constructed by (2.12) or (2.4) with some structure measure c.

Example 2.3. Consider the special case Example 2.2, when $g = 0$. Then, define generalized translation operators as following

$$(L_{(x)}f)(y) = \frac{1}{2}\Big(f\big(|y - x|\big) + f\big(y + x\big)\Big), \quad x, y \in Q = \mathbb{R}_+, \tag{2.21}$$

and

$$\phi(x, \lambda) = \cos\Big(\sqrt{\lambda}x\Big), \quad v(x) = 1, \quad x \in \mathbb{R}_+, \ \lambda \in \mathbb{C}. \tag{2.22}$$

In fact, there are many examples of hypercomplex systems. For more details (see [5]).

III. Spaces of Test and Generalized Functions

The classical approach to white noise analysis [4, 25, 26] consists in the construction of some rigging of the Fock space $F(H)$ and in the consequent application to the spaces of this rigging the Wiener-Itô- Segal transform I. This operator transfers the space $F(H)$ to the space $L_2(\Omega', d\gamma(x))$ and the spaces of rigging to some spaces of test and generalized functions. The pairing between test and generalized functions is given by the scalar product in $L_2(\Omega', d\gamma(x))$. Here, Ω' is the dual space of a real nuclear space Ω and γ is a Gaussian measure on Ω'.

Ref

4. Yu. M. Berezansky and Yu. G. Kondratiev, *Spectral methods in infinite dimensional analysis*, vol. 1, 2, Kluwer, Dordrecht 1995.

© 2016 Global Journals Inc. (US)

In the model one-dimensional case the space $F(H)$ is equal to the space l_2 of complex-valued sequences $\varphi = (\varphi_0, \varphi_1, ...), \Omega' = \mathbb{R}$, and the Wiener-Itô- Segal transform has the form

$$l_2 \ni \varphi = (\varphi_0, \varphi_1, ...) \mapsto (I\varphi)(x) = \sum_{n=0}^{\infty} \varphi_n H_n(x) \ \in L_2(\mathbb{R}, d\gamma(x)), \ x \in \mathbb{R}, \tag{3.1}$$

where $\big(H_n(x)\big)_{n=0}^{\infty}$ is the sequence of orthonormal Hermite polynomials with respect to γ. The operator I transfers the rigging of l_2 to that of $L_2(\mathbb{R}, d\gamma(x))$, which gives a model of white noise analysis.

The generalization of white noise analysis proposed in [1], consists in the following. Let ρ be a probability measure on $\mathbb{R}$ (instead of the Gaussian measure γ) and let $(P_n(x), Q_n(x))_{n=0}^{\infty}$ $(x \in \mathbb{R})$ be the biorthogonal system of functions constructed in some canonical way. The biorthogonality means

$$\int_{\mathbb{R}} P_n(x) Q_m(x) d\rho(x) = \delta_{n,m}.\ n!, \quad , \quad m, n \in \mathbb{Z}_+. \tag{3.2}$$

Instead of the functions P_n, we take the Appell polynomials connected with the measure ρ, i.e. the coefficients of the power expansion with respect to $\lambda \in \mathbb{C}$

$$\widehat{\rho}(\lambda) = \int_{\mathbb{R}} e^{\lambda x} d\rho(x), \qquad \frac{e^{\lambda x}}{\widehat{\rho}\,(\lambda)} = \sum_{n=0}^{\infty} \frac{\lambda^n}{n!} P_n(x), \quad x \in \mathbb{R}. \tag{3.3}$$

The functions Q_n are defined by

$$Q_n(x) = \left(\left(\left(\frac{d}{dx} \right)^* \right)^n 1 \right)(x), \tag{3.4}$$

where the adjoint operator is considered in the $L_2(\mathbb{R}, d\rho(x))$ sense.

The operator I is defined now just as above, but with changing H_n by P_n. This operator transfers the rigging of l_2 to a rigging of $L_2(\mathbb{R}, d\rho(x))$, which gives some theory of generalized functions on $\mathbb{R}$ with pairing determinated by $L_2(\mathbb{R}, d\rho(x))$.

In this section, we suggest a generalization of the above mentioned construction in which instead of the function $e^{\lambda x}$ the character of some pretty arbitrary, commutative hypercomplex system is used. Such a generalization gives the possibility of constructing a lot of spaces of generalized functions connected with different examples of hypercomplex system.

Consider a subclass of the above hypercomplex systems for which the set of generalized characters is in one-to-one correspondence with the complex plane $\mathbb{C}$: $\chi \longleftrightarrow \lambda \in \mathbb{C}$; denote this character by $\chi(x, \lambda)$. We assume that $\chi(x, 0) = 1$ $(x \in Q)$, i.e., the unit character corresponds to $\lambda = 0$. We suppose the function $Q \times \mathbb{C} \ni \langle x, \lambda \rangle \mapsto \chi(x, \lambda) \in \mathbb{C}$ is continuous, and the function $\mathbb{C} \ni \lambda \mapsto \chi(x, \lambda) \in \mathbb{C}$ is an entire for each $x \in Q$. Thus, for each $x \in Q$ the following expansion holds

$$\chi(x, \lambda) = \sum_{n=0}^{\infty} \frac{\lambda^n}{n!} \chi_n(x), \quad \lambda \in \mathbb{C}, \tag{3.5}$$

where the coefficients $\chi_n \in C(Q)$ called the Delsarte characters [6]. It is possible, of course, to give a direct definition of the Delsarte characters if we will rewrite Eq.(2.13) in terms of χ_n.

Our purpose is now to construct the expansions of functions on Q using the Delsarte characters and to introduce the corresponding spaces of test and generalized functions.

Ref

6. Yu. M. Berezansky, A generalization of white noise analysis by means of theory of hypergroups, *Reports on Mathematical Physics* **38** (1996), 289-300.

© 2016 Global Journals Inc. (US)

Let ρ be a fixed Borel probability measure ($\rho(Q) = 1$), positive on open sets, which is not connected directly with the multiplicative measure m . We suppose that χ_n's belong to $L_2(Q, d\rho(x))$, satisfy the estimate

$$\exists C > 0 : \|\chi_n\|_{L_2(Q,d\rho(x))} = \left(\int_Q |\chi_n(x)|^2 d\rho(x) \right)^2 \leq C^n n!, \quad n \in \mathbb{Z}_+. \tag{3.6}$$

(explaining that always $\|\chi_n\|_{L_2(Q,d\rho(x))} = 1$), are linearly independent and the system $(\chi_n)_n^\infty$ is total in $L_2(Q, d\rho(x))$.

Note that for a large class of hypercomplex systems (the Taylor-Delsarte hypercomplex system) vectors χ_n are automatically linearly independent. If in the estimate (3.6) instead of $n!$ the quantity $(n!)^\epsilon$ with fixed ($\epsilon < 1$) stands or if supp ρ is a compact set, then also automatically the system $(\chi_n)_n^\infty$ is total. In general we will understand the above mentioned conditions as some conditions on the measure ρ.

Now, we use the simple constructions connected with Hilbert rigging [4].

Let us consider a rigging of a Hilbert space H_0 with positive and negative spaces H_+ and H_- as following

$$H_- \supseteq H_0 \supseteq H_+. \tag{3.7}$$

Let $V : H_- \longrightarrow H_+$ be the canonical isometry transferring H_- onto H_+ [4]. A biorthogonal basis $(p_n, q_n)_{n=0}^\infty$ in the space H_0 is to be understood as sequences $(p_n)_{n=0}^\infty \subset H_+$ and $(q_n = V^{-1}p_n)_{n=0}^\infty \subset H_-$, where the first one is an orthogonal basis in H_+ and therefore, the second one is an orthogonal basis in H_-.

Note that these systems of vectors p_n and q_n are biorthogonal (with respect to H_o):

$$(p_n, q_n)_{H_0} = \delta_{n,m} h_n, \quad h_n = \|p_n\|^2_{H_+} = \|q_n\|^2_{H_-}, \quad n, m \in \mathbb{Z}_+, \tag{3.8}$$

$$\forall \varphi \in H_+, \quad \varphi = \sum_{n=0}^\infty \varphi_n p_n, \quad \varphi_n = (\varphi, q_n)_{H_0} h_n^{-1}, \quad \sum_{n=0}^\infty |\varphi_n|^2 h_n = \|\varphi\|^2_{H_+} < \infty,$$

$$\forall \xi \in H_-, \quad \xi = \sum_{n=0}^\infty \xi_n q_n, \quad \xi_n = (\xi, p_n)_{H_0} h_n^{-1}, \quad \sum_{n=0}^\infty |\xi_n|^2 h_n = \|\xi\|^2_{H_-} < \infty,$$

$$(\xi, \varphi)_{H_0} = \sum_{n=0}^\infty \xi_n \overline{\varphi_n} h_n. \tag{3.9}$$

This definition is justified by the following.

Let $(p_n)_{n=0}^\infty$ be an arbitrary total sequence of vectors p_n of a Hilbert space H_0. It is easy to prove that such sequence $(h_n)_{n=0}^\infty$ of positive numbers h_n exists for which the set of test functions

$$H_+ = \left\{ \varphi = \sum_{n=0}^\infty \varphi_n \, p_n \mid \varphi_n \in \mathbb{C} : \|\varphi\|^2_{H_+} = \sum_{n=0}^\infty |\varphi_n|^2 \, h_n < \infty \right\}, \tag{3.10}$$

with the corresponding scalar product is the positive space with respect to H_0 and therefore the chain (3.7) exists. Note that it is necessary to assume in addition the fulfilment of the following necessary and sufficient condition on $(p_n)_{n=0}^\infty$: an arbitrary sequence $(\varphi^{(i)})_{i=0}^\infty$ of vectors $\varphi^{(i)} \in H_+$ with finite sequences of coordinates $\varphi_n^{(i)}$ which is fundamental in H_+ and converges to 0 in H_0 must converge to 0 in H_+. This condition will always be fulfilled in our case.

Ref

4. Yu. M. Berezansky and Yu. G. Kondratiev, *Spectral methods in infinite dimensional analysis*, vol. 1, 2, Kluwer, Dordrecht 1995.

© 2016 Global Journals Inc. (US)

Similarly for negative space H_- by replacement p_n by q_n, we have the set of generalized functions as follows

$$H_- = \left\{ \xi = \sum_{n=0}^{\infty} \xi_n \, q_n \mid \xi_n \in \mathbb{C} : \|\xi\|^2_{H_-} = \sum_{n=0}^{\infty} |\xi_n|^2 \, h_n < \infty \right\}. \tag{3.11}$$

Therefore, we can construct a rigging of the space $H_0 := L_2(Q, d\rho(x))$ of the form (3.7), (3.10) and (3.11) in which the Delsarte characters $\chi_n(x)$ will play the role of the vectors p_n. This result can be stated as follows:

Theorem 3.1. *There exists a quasinuclear rigging (3.7) such that* $H_0 = L_2(Q, d\rho(x))$,

$$H^{\chi}_{-1} \supseteq H_0 \supseteq H^{\chi}_1, \tag{3.12}$$

and the space $H^{\chi}_1 = H_+$ *is of the form (3.10), where*

$$p_n = \chi_n, \quad h_n = (n!)^2 K^n, \quad n \in \mathbb{Z}_+, \tag{3.13}$$

($K > 1$ *is a fixed sufficiently large number), and consists of continuous functions on* Q.

The system $(\chi_n, q^{\chi}_n)_{n=0}^{\infty}$, *where* $q^{\chi}_n = V^{-1}\chi_n$ *(*V *is connected with (3.12)), is a biorthogonal basis of the space* H_0. *For the vectors* q^{χ}_n, *the following representation holds:*

$$q^{\chi}_n = (n!)K^n \Theta_n, \quad \Theta_n = (D^+)^n \delta_e \in H^{\chi}_{-1}, \quad n \in \mathbb{Z}_+, \tag{3.14}$$

where Θ_n *are "Delsarte co-characters" connected with the hypercomplex system* $L_1(Q, d\rho(x))$, *and the measure* ρ. *Here* D *is a continuous operator acting on the space* H^{χ}_1 *and is defined by the expression*

$$D\chi_n = n\chi_{n-1}, \quad n \in \mathbb{N}, \quad D\chi_0 = 0, \tag{3.15}$$

$D^+ : H^{\chi}_{-1} \longrightarrow H^{\chi}_{-1}$ *is its dual operator with respect to the rigging (3.12). The space* H^{χ}_1 *consists of continuous functions, therefore the* δ*-function* δ_x, *lumped at the point* $x \in Q$, *exist as an element of the negative space* H^{χ}_{-1}.

It is essential to introduce the rigging of the space H_0 by means of projective and inductive limits of Hilbert spaces which are constructed by rules of type (3.10), (3.12) and (3.13). Namely, for every $q \in \mathbb{N}$ we introduce the Hilbert space of type (3.10):

$$H^{\chi}_q = \left\{ \varphi = \sum_{n=0}^{\infty} \varphi_n \chi_n \in H_0 : \|\varphi\|^2_{H^{\chi}_q} = \sum_{n=0}^{\infty} |\varphi_n|^2 (n!)^2 K^{qn} < \infty \right\}. \tag{3.16}$$

Then we have the nuclear rigging:

$$(\Psi^{\chi})' \supseteq H^{\chi}_{-q} \supseteq H_0 \supseteq H^{\chi}_q \supseteq \Psi^{\chi}, \tag{3.17}$$

$$\Psi^{\chi} = \operatorname{pr}\lim_{q \in \mathbb{N}} H^{\chi}_q = \bigcap_{q \in \mathbb{N}} H^{\chi}_q, \qquad (\Psi^{\chi})' = \operatorname{ind}\lim_{q \in \mathbb{N}} H^{\chi}_{-q} = \bigcup_{q \in \mathbb{N}} H^{\chi}_{-q},$$

$$H^{\chi}_{-q} = \left\{ \xi = \sum_{n=0}^{\infty} \xi_n q^{\chi}_n : \|\xi\|^2_{H^{\chi}_{-q}} = \sum_{n=0}^{\infty} |\xi_n|^2 (n!)^2 K^{-qn} < \infty \right\}, \tag{3.18}$$

© 2016 Global Journals Inc. (US)

with the action

$$(\xi,\varphi)_{H_0}=\sum_{n=0}^{\infty}\xi_n\overline{\varphi_n}(n!)^2K^{qn},\quad \varphi\in H_q^{\chi},\ \ \xi\in H_{-q}^{\chi}.$$

Note that due to the first equality in (3.8) for Delsarte characters and co-characters the following biorthogonality is true:

$$(\Theta_n,\chi_n)_{H_0}=\delta_{n,m}.n!,\quad n,m\in\mathbb{Z}_+. \tag{3.19}$$

To illustrate the above result, we give the following example

Example 3.1. Let $H_0=L_2(\mathbb{R},dx)$, with respect to the Lebesgue measure dx and ordinary convolution given in Eq.(2.12) the generalized character $\chi(x,\lambda)=e^{\lambda x}$ $(\lambda\in\mathbb{C})$ and $\chi_n(x)=x^n$ $(x\in\mathbb{R},\ n\in\mathbb{Z}_+)$. Therefore, the space (3.16) consists of entire functions $\varphi(x)$ and $\varphi_n(x)$ are the Taylor coefficients of $\varphi(x)$. Formula (3.8) gives their representation as the Fourier coefficients using the scalar product $(\xi,\varphi)_{H_0}$, $(\xi\in H_{-1}^{\chi},\ \varphi\in H_1^{\chi})$. The operator D now is equal d/dx.

IV. Elements of Wick Calculus

The Wick product was first introduced by Wick [39] and used as a tool to renormalize certain infinite quantities in quantum field theory. Later on, the Wick product was considered, in a stochastic setting, by Hida and Ikeda [24]. In [11], Dobroshin and Minlos were comprehensively treated this subject both in mathematical physics and probability theory. Currently, the Wick product provides a useful concept for various applications, for example, it is important in the study of stochastic ordinary and partial differential equations (see, e.g., [27]).

In this section, under the assumption that $\|\chi\|_{H_0}^2\le C^n$ for some $C>0$, we define a new Wick product, called χ-Wick product, on the space H_{-q}^{χ}, which is constructed in Section 3. Then, we give the definition of the χ-Hermite transform and apply it to establish a characterization theorem for the space H_{-q}^{χ}.

Definition 4.1. Let $\xi=\sum_{m=0}^{\infty}\xi_m q_m^{\chi}$, $\eta=\sum_{n=0}^{\infty}\eta_n q_n^{\chi}\in H_{-q}^{\chi}$ with $\xi_m,\eta_n\in\mathbb{C}$. The χ-Wick product of ξ, η, denoted by $\xi\diamond_{\chi}\eta$, is defined by the formula

$$\xi\diamond_{\chi}\eta=\sum_{m,n=0}^{\infty}\xi_m\eta_n q_{m+n}^{\chi}. \tag{4.1}$$

It is important to show that the spaces $H_{-q}^{\chi}, H_q^{\chi}$ are closed under χ-Wick product.

Lemma 4.1. If $\xi,\eta\in H_{-q}^{\chi}$ and $\varphi,\psi\in H_q^{\chi}$, *we have*

(i) $\xi\diamond_{\chi}\eta\in H_{-q}^{\chi}$,

(ii) $\varphi\diamond_{\chi}\quad\in H_q^{\chi}$.

Proof. If $\xi=\sum\limits_{m=0}^{\infty}\xi_m q_m^{\chi}$, $\eta=\sum\limits_{n=0}^{\infty}\eta_n q_n^{\chi}\in H_{-q}^{\chi}$, then for some $q_1\in\mathbb{N}$ we have

$$\sum_{m=0}^{\infty}|\xi_m|^2K^{-q_1m}<\infty\ \text{and}\ \sum_{n=0}^{\infty}|\eta_n|^2K^{-q_1n}<\infty. \tag{4.2}$$

Ref
39. G. C. Wick, The evaluation of the collinear matrix, *Phys. Rev.* **80** (1950), 268-272.

© 2016 Global Journals Inc. (US)

We note that

$$\xi \diamond_\chi \eta = \sum_{m,n=0}^{\infty} \xi_m \eta_n q_{m+n}^\chi = \sum_{l=0}^{\infty} \left(\sum_{m+n=l}^{\infty} \xi_m \eta_n \right) q_l^\chi = \sum_{l=0}^{\infty} \zeta_l q_l^\chi, \tag{4.3}$$

where $\zeta_l = \sum_{m+n=l}^{\infty} \xi_m \eta_n$. With $q = q_1 + p$ we have

$$\begin{aligned} \sum_{l=0}^{\infty} |\zeta_l|^2 K^{-ql} &= \sum_{l=0}^{\infty} \left| \sum_{m+n=l}^{\infty} \xi_m \eta_n \right|^2 K^{-q_1 l} K^{-pl} \\ &\leq \sum_{l=0}^{\infty} \left(\sum_{m+n=l}^{\infty} |\xi_m|^2 K^{-q_1 m} \right) \left(\sum_{m+n=l}^{\infty} |\eta_n|^2 K^{-q_1 n} \right) K^{-pl} \\ &\leq \left(\sum_{l=0}^{\infty} K^{-pl} \right) \left(\sum_{m=0}^{\infty} |\xi_m|^2 K^{-q_1 m} \right) \left(\sum_{n=0}^{\infty} |\eta_n|^2 K^{-q_1 n} \right) \\ &< \infty, \end{aligned} \tag{4.4}$$

which proves (i). The proof of (ii) is similar. ■

The following important algebraic properties of the χ-Wick product follow directly from Definition 4.1.

Lemma 4.2. *For each $\xi, \eta, \zeta \in H_{-q}^\chi$, we get*

(i) $\xi \diamond_\chi \eta = \eta \diamond_\chi \xi$ *(Commutative law),*

(ii) $\xi \diamond_\chi (\eta \diamond_\chi \zeta) = (\xi \diamond_\chi \eta) \diamond_\chi \zeta$ *(Associative law),*

(iii) $\xi \diamond_\chi (\eta + \zeta) = \xi \diamond_\chi \eta + \xi \diamond_\chi \zeta$ *(Distributive law).*

Remark 4.1. According to Lemmas 4.1 and 4.2, we can conclude that the spaces H_{-q}^χ and H_q^χ form topological algebras with respect to the χ-Wick product.

As shown in Lemmas 4.1 and 4.2, the χ-Wick product satisfies all the ordinary algebraic rules for multiplication. Therefore, one can carry out calculations in much the same way as with usual products. But, there are some problems when limit operations are involved. To treat these situations it is convenient to apply a transformation, called the χ-Hermite transform, which converts χ-Wick products into ordinary (complex) products and convergence in H_{-q}^χ into bounded, pointwise convergence in a certain neighborhood of 0 in $\mathbb{C}$. The original Hermite transform, which first appeared in Lindstrøm et al. [29], has been applied by the authors in many different connections. Now, we give the definition of the χ-Hermite transform and discuss its basic properties.

Definition 4.2. Let $\xi = \sum_{n=0}^{\infty} \xi_n q_n^\chi \in H_{-q}^\chi$ with $\xi_n \in \mathbb{C}$. Then, the χ-Hermite transform of ξ, denoted by $\mathcal{H}_\chi \xi$, is defined by

$$\mathcal{H}_\chi \xi(z) = \sum_{n=0}^{\infty} \xi_n z^n \in \mathbb{C} \quad \text{(when convergent)}. \tag{4.5}$$

In the following we define for $0 < M, q < \infty$ the neighborhoods $\mathbb{O}_q(M)$ of zero in $\mathbb{C}$ by

$$\mathbb{O}_q(M) = \left\{ z \in \mathbb{C} : \sum_{n=0}^{\infty} |z^n|^2 K^{qn} < M^2 \right\}. \tag{4.6}$$

Ref

29. T. Lindstrom, B. Oksendal and J. Uboe, Stochastic differential equations involving positive noise. In M. Barlow and N. Bingham (editors): *Stochastic Analysis.* Cambridge University Press: Cambridge, MA, (1991), 261-303.

© 2016 Global Journals Inc. (US)

Notes

It is easy to see that

$$q \leq p,\ N \leq M \Rightarrow \mathbb{O}_p(N) \subseteq \mathbb{O}_q(M). \tag{4.7}$$

Note that if $\xi = \sum_{n=0}^{\infty} \xi_n q_n^{\chi} \in H_{-q}^{\chi}, z \in \mathbb{O}_q(M)$ for some $0 < M, q < \infty$, we have the estimate

$$\begin{aligned} \sum_{n=0}^{\infty} |\xi_n||z^n| &= \sum_{n=0}^{\infty} |\xi_n||z^n| K^{-\frac{qn}{2}} K^{\frac{qn}{2}} \\ &\leq \left(\sum_{n=0}^{\infty} |\xi_n|^2 K^{-qn} \right)^{\frac{1}{2}} \left(\sum_{n=0}^{\infty} |z^n|^2 K^{qn} \right)^{\frac{1}{2}} \\ &< M \left(\sum_{n=0}^{\infty} |\xi_n|^2 K^{-qn} \right)^{\frac{1}{2}} \\ &< \infty. \end{aligned} \tag{4.8}$$

The conclusion above can be stated as follows:

Proposition 4.1. *If $\xi \in H_{-q}^{\chi}$, then $\mathcal{H}_{\chi}\xi$ converges for all $z \in \mathbb{O}_q(M)$ for all $q, M < \infty$.*

A useful property of the χ-Hermite transform is that it converts the χ-Wick product into ordinary (complex) product.

Proposition 4.2. *If $\xi, \eta \in H_{-q}^{\chi}$, then*

$$\mathcal{H}_{\chi}(\xi \diamond_{\chi} \eta)(z) = \mathcal{H}_{\chi}\xi(z).\mathcal{H}_{\chi}\eta(z). \tag{4.9}$$

for all z such that $\mathcal{H}_{\chi}\xi$ and $\mathcal{H}_{\chi}\eta$ exist.

Proof. The proof is an immediate consequence of Definitions 4.1 and 4.2. ■

Let $\xi = \sum_{n=0}^{\infty} \xi_n q_n^{\chi} \in H_{-q}^{\chi}$, with $\xi_n \in \mathbb{R}$. Then, the number $\xi_0 = \mathcal{H}_{\chi}\xi(0) \in \mathbb{R}$ is called the generalized expectation of ξ and is denoted by $\mathbb{E}(\xi)$. Suppose that $V \ni z \mapsto f(z) \in \mathbb{C}$ is an analytic function, where V is a neighborhood of $\mathbb{E}(\xi)$. Assume that the Taylor series of f around $\mathbb{E}(\xi)$ has coefficients in $\mathbb{R}$. Then, the χ-Wick version $f^{\diamond_{\chi}}$ of f is defined by

$$H_{-q}^{\chi} \ni \xi \mapsto f^{\diamond_{\chi}}(\xi) = \mathcal{H}^{-1}\big(f \circ \mathcal{H}_{\chi}(\xi)\big) \in H_{-q}^{\chi}. \tag{4.10}$$

Example 4.1. If the function $f : \mathbb{C} \to \mathbb{C}$ is entire, then $f^{\diamond_{\chi}}$ is defined for all $\xi \in H_{-q}^{\chi}$. For example, the χ-Wick exponential is defined by

$$\exp^{\diamond_{\chi}}(\xi) = \sum_{j=0}^{\infty} \frac{1}{j!} \xi^{\diamond_{\chi} n}. \tag{4.11}$$

Using χ-Hermite transform we see that χ-Wick exponential has the same algebraic properties as the usual exponential. For instance,

$$\exp^{\diamond_{\chi}}(\xi + \eta) = \exp^{\diamond_{\chi}}(\xi) \diamond_{\chi} \exp^{\diamond_{\chi}}(\eta),\ \xi, \eta \in H_{-q}^{\chi}. \tag{4.12}$$

From Proposition 4.1, we can deduce that χ-Hermite transform of any $\xi \in H_{-q}^{\chi}$ is a complex-valued analytic function on $\mathbb{O}_q(M)$ for all $q, M < \infty$. Moreover, the converse of this deduction is true, i.e., every complex-valued analytic function on $\mathbb{O}_q(M)$ (for some $q, M < \infty$) is the χ-Hermite transform of some element in H_{-q}^{χ}. To prove this, we need the following two auxiliary results.

© 2016 Global Journals Inc. (US)

Lemma 4.3. *Let $f(z) = \sum_{n=0}^{\infty} \eta_n z^n$ be an analytic function in $z \in \mathbb{C}$ such that there exists $M < \infty, C > 0$ and $\delta > 0$ such that $|f(z)| \leq M$ when $z \in \mathbb{O} := \{z \in \mathbb{C} : C|z| \leq \delta^2\}$. Then $|\eta_n z^n| \leq M$ for all $z \in \mathbb{O}$ and $n \in \mathbb{N}$.*

Proof. See [27], Lemma 2.6.10.

Proposition 4.3. *Let $f(z) = \sum_{n=0}^{\infty} \eta_n z^n$, $\eta_n \in \mathbb{C}$ be a formal power series in $z \in \mathbb{C}$. Suppose there exist $q, M < \infty$ and $\delta > 0$ such that $f(z)$ is convergent for $z \in \mathbb{O}_q(\delta)$ and $|f(z)| \leq M$ for all $z \in \mathbb{O}_q(\delta)$. Then*

$$\sum_{n=0}^{\infty} |\eta_n z^n| \leq MA(q) \quad \text{for all } z \in \mathbb{O}_{3q}(\delta) \tag{4.13}$$

where

$$A(q) := \sum_{n=0}^{\infty} K^{-qn} < \infty \quad (\text{Note that } K > 1). \tag{4.14}$$

Proof. It is evident that $z \in \mathbb{O}_{3q}\ \delta)$ implies $K^q z \in \mathbb{O}_q\ \delta)$. According to Lemma 4.3, we get

$$\begin{aligned} \sum_{n=0}^{\infty} |\eta_n z^n| &\leq \left(\sum_{n=0}^{\infty} |\eta_n|^2 |z^n|^2 K^{qn} \right)^{\frac{1}{2}} \left(\sum_{n=0}^{\infty} K^{-qn} \right)^{\frac{1}{2}} \\ &= \left(\sum_{n=0}^{\infty} |\eta_n|^2 |(K^q z)^n|^2 K^{-qn} \right)^{\frac{1}{2}} \left(\sum_{n=0}^{\infty} K^{-qn} \right)^{\frac{1}{2}} \\ &\leq M \sum_{n=0}^{\infty} K^{-qn}. \end{aligned} \tag{4.15}$$

Theorem 4.1. **(Characterization Theorem for H_{-q}^{χ})** *If $\xi = \sum_{n=0}^{\infty} \xi_n q_n^{\chi} \in H_{-q}^{\chi}$, where $\xi_n \in \mathbb{C}$, then there exist $q < \infty$ and $R_q < \infty$ such that*

$$|\mathcal{H}_{\chi}\xi(z)| \leq R_q \left(\sum_{n=0}^{\infty} |z^n|^2 K^{qn} \right)^{\frac{1}{2}} \quad \forall z \in \mathbb{C}. \tag{4.16}$$

In particular, $\mathcal{H}_{\chi}\xi$ is a bounded analytic function on $\mathbb{O}_q(M)$ for all $M < \infty$. Conversely, suppose $f(z) = \sum_{n=0}^{\infty} \eta_n z^n$ is a given analytic power series of $z \in \mathbb{C}$ with $\eta_n \in \mathbb{C}$ such that there exist $q < \infty$ and $\delta > 0$, such that $f(z)$ is absolutely convergent when $z \in \mathbb{O}_q(\delta)$ and

$$\sup_{z \in \mathbb{O}_q(\delta)} |f(z)| < \infty. \tag{4.17}$$

Then, there exists a unique $\eta \in H_{-q}^{\chi}$ such that $\mathcal{H}_{\chi}\eta = f$, namely

$$\eta = \sum_{n=0}^{\infty} \eta_n q_n^{\chi}. \tag{4.18}$$

Proof. For each $z \in \mathbb{C}$, we have

$$|\mathcal{H}_{\chi}\xi(z)| \leq \sum_{n=0}^{\infty} |\xi_n||z^n| \leq \left(\sum_{n=0}^{\infty} |\xi_n|^2 K^{-qn} \right)^{\frac{1}{2}} \left(\sum_{n=0}^{\infty} |z^n|^2 K^{qn} \right)^{\frac{1}{2}}. \tag{4.19}$$

Since $\xi \in H_{-q}^{\chi}$, we see that $R_q^2 := \sum_{n=0}^{\infty} |\xi_n|^2 K^{-qn} < \infty$ for all $q < \infty$.

Ref
27. H. Holden, B. Osendal, J. Uboe and T. Zhang, *Stochastic partial differential equations*, Springer Science+Business Media, LLC, (2010).

© 2016 Global Journals Inc. (US)

Conversely, Since $K > 1$, then $K^{-r} \in \mathbb{O}_r(\delta)$ for all $r < \infty$ and for some $\delta < \infty$. By virtue of Proposition 4.3, we have

$$\sum_{n=0}^{\infty} |\eta_n||z^n| \leq MA(q) \quad \text{for all } z \in \mathbb{O}_{3q}(\delta). \tag{4.20}$$

Hence, for $r \geq 3q$ and $z \in \mathbb{O}_{3q}(\delta)$, we get

$$\sum_{n=0}^{\infty} |\eta_n|^2 K^{-rn} \leq C \sum_{n=0}^{\infty} |\eta_n| K^{-rn} \leq C \sum_{n=0}^{\infty} |\eta_n||z^n| \leq CMA(q) < \infty. \tag{4.21}$$

where $C := \sup\{|\eta_n| : n \in \mathbb{N}\}$, and hence $\eta := \sum_{n=0}^{\infty} \eta_n q_n^{\chi} \in H_{-q}^{\chi}$, as claimed. ■

Notes

References Références Referencias

1. S. Albeverio, Yu. G. Kondratiev and L. Streit, How to generalize white noise analysis to non-Gaussian measures. Proc. Symp. *Dynamics of Complex and Irregular Systems* (Bielefeld, 1991); Bielefeld Encount. Math. Phys., VIII, World Sci. Publ.: River Edge, NJ, (1993), 120-130.
2. S. Albeverio, Yu. L. Daletsky, Yu. G. Kondratiev and L. Streit, Non-Gaussian infinitedimensional analysis, *J. Funct. Anal.* **138** (1996), 311-350.
3. Yu. M. Berezansky and Yu. S. Samoilenko, Nuclear spaces of functions of an infinite number of variables, *Ukrainian Math. J.* **25** (1973), 723-737.
4. Yu. M. Berezansky and Yu. G. Kondratiev, *Spectral methods in infinite dimensional analysis*, vol. 1, 2, Kluwer, Dordrecht 1995.
5. Yu. M. Berezansky and A. A. Kalyuzhnyi, *Harmonic Analysis in Hypercomplex Systems*, Naukova Dumka,Kyiv, Ukrania, 1992. (in Russian; English transl.: Kluwer: Dordrecht-Boston-London, 1996).
6. Yu. M. Berezansky, A generalization of white noise analysis by means of theory of hypergroups, *Reports on Mathematical Physics* **38** (1996), 289-300.
7. Yu. M. Berezansky, Spectral approach to white noise analysis.Proc. Symp. *Dynamics of Complex and Irregular Systems* (Bielefeld, 1991); Bielefeld Encount. Math. Phys., VIII, World Sci. Publ., River Edge, NJ, (1993), 131-140.
8. Yu. M. Berezansky and Yu. G. Kondratiev, Biorthogonal systems in hypergroups: an extension of non-Gaussian analysis, *Methods Funct. Anal. Topopology* **2** (1996), 1-50.
9. Yu. M. Berezansky and A. A. Kalyuzhnyi, Harmonic Analysis in Hypercomplex Systems, Naukova Dumka: Kiev, 1992 (in Russian; English transl.: Kluwer: Dordrecht-Boston-London, 1996).
10. W. R. Bloom and H. Heyer, *Harmonic Analysis of Probability Measures on Hypergroups*, de Gruyter, Berlin 1994.
11. R. L. Dobrushin and R. L. Minlos, Polynomials in linear random functions, *Russian Math. Surveys* **32** (1977), 71-127.
12. Yu. L. Daletsky, A biorthogonal analogy of the Hermite polynomials and the inversion of the Fourier transform with respect to a non-Gaussian measure, *Funct. Anal.Appl.* **25** (1991), 138-40.
13. J. Delsarte, Sur certaines transformation fonctionelles relative aux equations linearies aux derivees partiels du seconde ordre, *C. R. Acad. Sci., Ser.* **17** (1938), 178-182.
14. R. J. Elliott and J. van der Hoek, A general fractional white noise theory and applications to finance, *Math. Finance* **13** (2003), 301-330.

© 2016 Global Journals Inc. (US)

15. H. A. Ghany, A. S. Okb El Bab, A. M. Zabel, and A. Hyder, The fractional coupled KdV equations: exact solutions and white noise functional approach,*Chinese Physics B*, vol. **22**, no. 8, (2013) Article ID 080501.
16. H. A. Ghany, Exact solutions for stochastic generalized Hirota-Satsuma coupled KdV equations, *Chinese Journal of Physics*, vol. **49**, no. 4, (2011) pp. 926940.
17. H. A. Ghany and A. Hyder, Local and global well-posedness of stochastic Zakharov-Kuznetsov equation *Journal of Computational Analysis and Applications,* **15** (7), 1332-1343.
18. H. A. Ghany and A. Hyder, Exact solutions for the wicktype stochastic time-fractional KdV equations, *Kuwait Journal of Science*, vol. 41, no. 1,(2014) pp. 7584.
19. H. A. Ghany and A. Hyder, Abundant solutions of Wick-type stochastic fractional 2D KdV equations, *Chinese Physics B,*, vol. **23**, no. 6,(2014) Article ID 060503.
20. H. A. Ghany and M. Zakarya, Generalized Solutions of Wick-type Stochastic KdV-Burgers Equations Using Exp-function Method *International Review of Physics,* (IREPHY) **8**(2),(2014) 38-46.
21. H. A. Ghany and M. Zakarya, Exact Solutions for Wick-type Stochastic Coupled KdV Equations *Global Journal of Science Frontier Research: F Mathematics and Decision Sciences,* **14**(2014) 56-72.
22. H. A. Ghany and M. Zakarya, Exact Traveling Wave Solutions for Wick-Type Stochastic Schamel KdV Equation *Physics Research International,* (2014) p. 1-9
23. H. A. Ghany and M. A. Qurashi, Travelling SolitaryWave Solutions for Stochastic Kadomtsev-Petviashvili Equation. *Journal of Computational Analysis and Applications,* **21** (1), (2015) p.121-131.
24. T. Hida and N. Ikeda, Analysis on Hilbert space with reproducing kernel arising from multiple Wiener integral. *Proc. Fifth Berkeley Symp.Math. Stat. Probab. II* (1965), 117-143. Univ. California Press, Berkeley.
25. T. Hida, *Analysis of Brownian functionals*, in Carleton Math. Lecture Notes **13**, Carleton Univ., Ottawa 1975.
26. T. Hida, H-H. Kuo, J. Potthoff and L. Streit, *White Noise: An infinite Dimensional Calculus*, Kluwer, Dordrecht 1993.
27. H. Holden, B. Osendal, J. Uboe and T. Zhang, *Stochastic partial differential equations,* Springer Science+Business Media, LLC, (2010).
28. H-H. Kuo, White noise theory, *Handbook of Stochastic Analysis and Applications* D. Kannan and V. Lakshmikantham (eds.) (2002), 107-158.
29. T. Lindstrom, B. Oksendal and J. Uboe, Stochastic differential equations involving positive noise. In M. Barlow and N. Bingham (editors): *Stochastic Analysis*. Cambridge University Press: Cambridge, MA, (1991), 261-303.
30. A. A. Kalyuzhnyi, A theorem on the existence of multiplicative measure, *Ukr. Math. Zh.,* **35**(1983), 369-371.
31. B. M. Levitan, Generalization of translation operation and infinite-dimensional hypercomplex systems, *Mat. Sb.* **16** (1945), 259-280.
32. B. M. Levitan, Application of generalized translation operators to linear differential equations of the second order, *Usp. Mat. Nauk* **4** (1949), 3-112.
33. B. M. Levitan, Lie theorems for generalized translation operators, *Usp. Mat. Nauk* **16** (1961), 3-30.
34. B. M. Levitan, *Generalized translation operators and some their applications*, Nauka: Moscow, 1962.
35. B. M. Levitan, *Theory of generalized translation operators*, Nauka: Moscow, 1973.

Notes

© 2016 Global Journals Inc. (US)

36. G. L. Litvinov, On generalized translation operators and their representations, *Tr. Sem. Vekt. Anal. MGU* **18** (1978), 345-371.
37. A. S. Okb El Bab, A. M. Zabel, H. A. Ghany, Harmonic analysis in Hypercomplex systems, *International Journal of Pure and Applied Mathematics*, **80** (2012), 739-750.
38. G. F. Us, Dual Appel systems in Poissonian analysis. *Methods Funct. Anal. Topopology* (1995), 93-108.
39. G. C. Wick, The evaluation of the collinear matrix, *Phys. Rev.* **80** (1950), 268-272.
40. A. M. Zabel and Buthinah A. Bin Dehaish, Negative definite functions on hypercomplex systems, *KYUNGPOOK Math. Journal* **46** (2006), 285-295.
41. A. M. Zabel and Buthinah A. Bin Dehaish, Ljevy Khinchin formula on commutative hypercomplex system, *KYUNGPOOK Math. Journal* **48** (2008), 559-575.

Notes

© 2016 Global Journals Inc. (US)

This page is intentionally left blank

© 2016 Global Journals Inc. (US)

Global Journal of Science Frontier Research: F
Mathematics and Decision Sciences
Volume 16 Issue 1 Version 1.0 Year 2016
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Visualizing Finite Field of Order p^2 through Matrices

By S. K. Pandey
Sardar Patel University of Police, India

Abstract- In this article we consider finite field of order $p^2(p \neq 2)$. We provide matrix representation of finite field of order at most $p^2 = 121$. However this article provides a technique to yield matrix field of order p^2 for every prime $p > 2$.

Keywords: *finite field, matrix field, Galois field.*

GJSFR-F Classification : *MSC 2010: 12E20*



Strictly as per the compliance and regulations of :



© 2016. S. K. Pandey. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Visualizing Finite Field of Order p^2 through Matrices

S. K. Pandey

Abstract- In this article we consider finite field of order $p^2(p \neq 2)$. We provide matrix representation of finite field of order at most $p^2 = 121$. However this article provides a technique to yield matrix field of order p^2 for every prime $p > 2$.

Keywords: finite field, matrix field, Galois field.

MSCS2010: 12E20.

I. Introduction

There is a finite field of order p^n for every positive prime p and positive integer n. If F is a finite field of order p and $F[x]$ is the polynomial domain defined over F then one can construct a finite field $K = \frac{F[x]}{[f(x)]}$ of order p^n, where $f(x) \in F[x]$ is an irreducible polynomial of degree n and $[f(x)]$ is the ideal generated by $f(x)$. This field K has a subfield isomorphic with the field F. One may refer [1, 2] for further details. This is the usual method of construction of finite fields.

In this article we construct finite matrix fields of order p^2 without adopting the usual method of construction of finite fields. We give finite matrix field of order at most 121. However following the approach given in this article one can construct finite matrix field of order p^2 for each $p \neq 2$.

II. Finite Matrix Field of Order p^2

In [3] one can find different matrix representations of a finite field of order p. Let p be a prime number. Then $Z_p = \{0,1,2,3,4,5 \ldots p-1\}$ is a field under addition and multiplication modulo p. One matrix representation of Z_p as given in [3] is $F_p = \left\{ \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} : a \in Z_p \right\}$. Every field of characteristic p contains a copy of a field of order

Author: Department of Mathematics, Sardar Patel University of Police, Security and Criminal Justice Daijar, Jodhpur, Rajasthan., India.
e-mail: skpandey12@@gmail.com

© 2016 Global Journals Inc. (US)

p. All the fields given below are finite matrix field of order p^2. One can easily see that all these fields contain F_p.

We use very simple technique to obtain matrix fields of order p^2. It is seen that $F = \left\{ \begin{pmatrix} a & b \\ -b & a \end{pmatrix} : a,b \in R \right\}$ is isomorphic to the field C of complex numbers. Here R stands for the field of real numbers. We notice that if we replace R by Z_p then we obtain a finite matrix field of order p^2. It may be noted that this approach does not work for $p = 2$.

a) Finite Matrix Field Of Order 9

$$M_9 = \left\{ \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 2 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 2 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 2 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 1 \\ 2 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 2 \\ 1 & 2 \end{pmatrix} \right\}$$

This is a finite field of order 9 under addition and multiplication of matrices modulo 3. It is a field of characteristic 3 and it contains $F_3 = \left\{ \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} \right\}$.

b) Finite Matrix Field Of Order 25

$$M_{25} = \left\{ \begin{array}{l} \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}, \begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}, \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 4 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 4 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 1 \\ 4 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 4 \\ 1 & 2 \end{pmatrix}, \begin{pmatrix} 3 & 4 \\ 1 & 3 \end{pmatrix}, \begin{pmatrix} 3 & 1 \\ 4 & 3 \end{pmatrix}, \\ \begin{pmatrix} 4 & 1 \\ 4 & 4 \end{pmatrix}, \begin{pmatrix} 4 & 4 \\ 1 & 4 \end{pmatrix}, \begin{pmatrix} 0 & 4 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 4 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 2 \\ 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 3 \\ 2 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 3 \\ 2 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 2 \\ 3 & 0 \end{pmatrix}, \begin{pmatrix} 2 & 2 \\ 3 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 3 \\ 2 & 2 \end{pmatrix}, \begin{pmatrix} 3 & 3 \\ 2 & 3 \end{pmatrix}, \\ \begin{pmatrix} 3 & 2 \\ 3 & 3 \end{pmatrix}, \begin{pmatrix} 4 & 2 \\ 3 & 4 \end{pmatrix}, \begin{pmatrix} 4 & 3 \\ 2 & 4 \end{pmatrix} \end{array} \right\}$$

This is a finite field of order 25 under addition and multiplication of matrices modulo 5. One can see that it is a field of characteristic 5 and it contains

$$F_5 = \left\{ \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}, \begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}, \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix} \right\}.$$

c) Finite Matrix Field Of Order 49

$$M_{49} = \left\{ \begin{array}{l} \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}, \begin{pmatrix} 3 & 0 \\ 0 & 3 \end{pmatrix}, \begin{pmatrix} 4 & 0 \\ 0 & 4 \end{pmatrix}, \begin{pmatrix} 5 & 0 \\ 0 & 5 \end{pmatrix}, \begin{pmatrix} 6 & 0 \\ 0 & 6 \end{pmatrix}, \begin{pmatrix} 0 & 6 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 6 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 2 \\ 5 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 5 \\ 2 & 0 \end{pmatrix}, \\ \begin{pmatrix} 0 & 4 \\ 3 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 3 \\ 4 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 6 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 6 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 2 \\ 5 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 5 \\ 2 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 4 \\ 3 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 3 \\ 4 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 3 \\ 4 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 4 \\ 3 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 5 \\ 2 & 2 \end{pmatrix}, \\ \begin{pmatrix} 2 & 2 \\ 5 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 6 \\ 1 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 1 \\ 6 & 2 \end{pmatrix}, \begin{pmatrix} 3 & 4 \\ 3 & 3 \end{pmatrix}, \begin{pmatrix} 3 & 3 \\ 4 & 3 \end{pmatrix}, \begin{pmatrix} 3 & 5 \\ 2 & 3 \end{pmatrix}, \begin{pmatrix} 3 & 2 \\ 5 & 3 \end{pmatrix}, \begin{pmatrix} 3 & 1 \\ 6 & 3 \end{pmatrix}, \begin{pmatrix} 3 & 6 \\ 1 & 3 \end{pmatrix}, \begin{pmatrix} 4 & 4 \\ 3 & 4 \end{pmatrix}, \begin{pmatrix} 4 & 3 \\ 4 & 4 \end{pmatrix}, \\ \begin{pmatrix} 4 & 5 \\ 2 & 4 \end{pmatrix}, \begin{pmatrix} 4 & 2 \\ 5 & 4 \end{pmatrix}, \begin{pmatrix} 4 & 1 \\ 6 & 4 \end{pmatrix}, \begin{pmatrix} 4 & 6 \\ 1 & 4 \end{pmatrix}, \begin{pmatrix} 5 & 4 \\ 3 & 5 \end{pmatrix}, \begin{pmatrix} 5 & 3 \\ 4 & 5 \end{pmatrix}, \begin{pmatrix} 5 & 5 \\ 2 & 5 \end{pmatrix}, \begin{pmatrix} 5 & 2 \\ 5 & 5 \end{pmatrix}, \begin{pmatrix} 5 & 6 \\ 1 & 5 \end{pmatrix}, \begin{pmatrix} 5 & 1 \\ 6 & 5 \end{pmatrix}, \begin{pmatrix} 6 & 4 \\ 3 & 6 \end{pmatrix}, \\ \begin{pmatrix} 6 & 3 \\ 4 & 6 \end{pmatrix}, \begin{pmatrix} 6 & 5 \\ 2 & 6 \end{pmatrix}, \begin{pmatrix} 6 & 2 \\ 5 & 6 \end{pmatrix}, \begin{pmatrix} 6 & 1 \\ 6 & 6 \end{pmatrix}, \begin{pmatrix} 6 & 6 \\ 1 & 6 \end{pmatrix} \end{array} \right\}$$

Notes

© 2016 Global Journals Inc. (US)

This is a finite field of order 49 under addition and multiplication of matrices modulo 7. One can see that it is a field of characteristic 7 and it contains

$$F_7 = \left\{\begin{pmatrix}0&0\\0&0\end{pmatrix},\begin{pmatrix}1&0\\0&1\end{pmatrix},\begin{pmatrix}2&0\\0&2\end{pmatrix},\begin{pmatrix}3&0\\0&3\end{pmatrix},\begin{pmatrix}4&0\\0&4\end{pmatrix},\begin{pmatrix}5&0\\0&5\end{pmatrix},\begin{pmatrix}6&0\\0&6\end{pmatrix}\right\}.$$

Notes

d) Finite Matrix Field Of Order 121

$$M_{121}=\left\{\begin{pmatrix}0&0\\0&0\end{pmatrix},\begin{pmatrix}1&0\\0&1\end{pmatrix},\begin{pmatrix}2&0\\0&2\end{pmatrix},\begin{pmatrix}3&0\\0&3\end{pmatrix},\begin{pmatrix}4&0\\0&4\end{pmatrix},\begin{pmatrix}5&0\\0&5\end{pmatrix},\begin{pmatrix}6&0\\0&6\end{pmatrix},\begin{pmatrix}7&0\\0&7\end{pmatrix},\begin{pmatrix}8&0\\0&8\end{pmatrix},\begin{pmatrix}9&0\\0&9\end{pmatrix},\begin{pmatrix}10&0\\0&10\end{pmatrix},\right.$$
$$\begin{pmatrix}0&10\\1&0\end{pmatrix},\begin{pmatrix}0&1\\10&0\end{pmatrix},\begin{pmatrix}0&9\\2&0\end{pmatrix},\begin{pmatrix}0&2\\9&0\end{pmatrix},\begin{pmatrix}0&8\\3&0\end{pmatrix},\begin{pmatrix}0&3\\8&0\end{pmatrix},\begin{pmatrix}0&4\\7&0\end{pmatrix},\begin{pmatrix}0&7\\4&0\end{pmatrix},\begin{pmatrix}0&6\\5&0\end{pmatrix},\begin{pmatrix}0&5\\6&0\end{pmatrix},\begin{pmatrix}1&10\\1&1\end{pmatrix},$$
$$\begin{pmatrix}1&1\\10&1\end{pmatrix},\begin{pmatrix}1&2\\9&1\end{pmatrix},\begin{pmatrix}1&9\\2&1\end{pmatrix},\begin{pmatrix}1&8\\3&1\end{pmatrix},\begin{pmatrix}1&3\\8&1\end{pmatrix},\begin{pmatrix}1&4\\7&1\end{pmatrix},\begin{pmatrix}1&7\\4&1\end{pmatrix},\begin{pmatrix}1&6\\5&1\end{pmatrix},\begin{pmatrix}1&5\\6&1\end{pmatrix},\begin{pmatrix}2&10\\1&2\end{pmatrix},\begin{pmatrix}2&1\\10&2\end{pmatrix},$$
$$\begin{pmatrix}2&9\\2&2\end{pmatrix}\begin{pmatrix}2&2\\9&2\end{pmatrix},\begin{pmatrix}2&8\\3&2\end{pmatrix},\begin{pmatrix}2&3\\8&2\end{pmatrix},\begin{pmatrix}2&7\\4&2\end{pmatrix},\begin{pmatrix}2&4\\7&2\end{pmatrix},\begin{pmatrix}2&6\\5&2\end{pmatrix},\begin{pmatrix}2&5\\6&2\end{pmatrix},\begin{pmatrix}3&10\\1&3\end{pmatrix},\begin{pmatrix}3&1\\10&3\end{pmatrix},\begin{pmatrix}3&9\\2&3\end{pmatrix},$$
$$\begin{pmatrix}3&2\\9&3\end{pmatrix},\begin{pmatrix}3&8\\3&3\end{pmatrix},\begin{pmatrix}3&3\\8&3\end{pmatrix},\begin{pmatrix}3&7\\4&3\end{pmatrix},\begin{pmatrix}3&4\\7&3\end{pmatrix},\begin{pmatrix}3&6\\5&3\end{pmatrix},\begin{pmatrix}3&5\\6&3\end{pmatrix},\begin{pmatrix}4&10\\1&4\end{pmatrix},\begin{pmatrix}4&1\\10&4\end{pmatrix},\begin{pmatrix}4&2\\9&4\end{pmatrix},\begin{pmatrix}4&9\\2&4\end{pmatrix},$$
$$\begin{pmatrix}4&8\\3&4\end{pmatrix},\begin{pmatrix}4&3\\8&4\end{pmatrix},\begin{pmatrix}4&7\\4&4\end{pmatrix},\begin{pmatrix}4&4\\7&4\end{pmatrix},\begin{pmatrix}4&6\\5&4\end{pmatrix},\begin{pmatrix}4&5\\6&4\end{pmatrix},\begin{pmatrix}5&10\\1&5\end{pmatrix},\begin{pmatrix}5&1\\10&5\end{pmatrix},\begin{pmatrix}5&9\\2&5\end{pmatrix},\begin{pmatrix}5&2\\9&5\end{pmatrix},\begin{pmatrix}5&3\\8&5\end{pmatrix},$$
$$\begin{pmatrix}5&8\\3&5\end{pmatrix},\begin{pmatrix}5&7\\4&5\end{pmatrix},\begin{pmatrix}5&4\\7&5\end{pmatrix},\begin{pmatrix}5&6\\5&5\end{pmatrix},\begin{pmatrix}5&5\\6&5\end{pmatrix},\begin{pmatrix}6&10\\1&6\end{pmatrix},\begin{pmatrix}6&1\\10&6\end{pmatrix},\begin{pmatrix}6&9\\2&6\end{pmatrix},\begin{pmatrix}6&2\\9&6\end{pmatrix},\begin{pmatrix}6&8\\3&6\end{pmatrix},\begin{pmatrix}6&3\\8&6\end{pmatrix},$$
$$\begin{pmatrix}6&7\\4&6\end{pmatrix},\begin{pmatrix}6&4\\7&6\end{pmatrix},\begin{pmatrix}6&6\\5&6\end{pmatrix},\begin{pmatrix}6&5\\6&6\end{pmatrix},\begin{pmatrix}7&10\\1&7\end{pmatrix},\begin{pmatrix}7&1\\10&7\end{pmatrix},\begin{pmatrix}7&9\\2&7\end{pmatrix},\begin{pmatrix}7&2\\9&7\end{pmatrix},\begin{pmatrix}7&8\\3&7\end{pmatrix},\begin{pmatrix}7&3\\8&7\end{pmatrix},\begin{pmatrix}7&7\\4&7\end{pmatrix},$$
$$\begin{pmatrix}7&4\\7&7\end{pmatrix},\begin{pmatrix}7&6\\5&7\end{pmatrix},\begin{pmatrix}7&5\\6&7\end{pmatrix},\begin{pmatrix}8&10\\1&8\end{pmatrix},\begin{pmatrix}8&1\\10&8\end{pmatrix},\begin{pmatrix}8&9\\2&8\end{pmatrix},\begin{pmatrix}8&2\\9&8\end{pmatrix},\begin{pmatrix}8&8\\3&8\end{pmatrix},\begin{pmatrix}8&3\\8&8\end{pmatrix},\begin{pmatrix}8&7\\4&8\end{pmatrix},\begin{pmatrix}8&4\\7&8\end{pmatrix},$$
$$\begin{pmatrix}8&6\\5&8\end{pmatrix},\begin{pmatrix}8&5\\6&8\end{pmatrix},\begin{pmatrix}9&10\\1&9\end{pmatrix},\begin{pmatrix}9&1\\10&9\end{pmatrix},\begin{pmatrix}9&9\\2&9\end{pmatrix},\begin{pmatrix}9&2\\9&9\end{pmatrix},\begin{pmatrix}9&8\\3&9\end{pmatrix},\begin{pmatrix}9&3\\8&9\end{pmatrix},\begin{pmatrix}9&7\\4&9\end{pmatrix},\begin{pmatrix}9&4\\7&9\end{pmatrix},\begin{pmatrix}9&6\\5&9\end{pmatrix},$$
$$\begin{pmatrix}9&5\\6&9\end{pmatrix},\begin{pmatrix}10&10\\1&10\end{pmatrix},\begin{pmatrix}10&1\\10&10\end{pmatrix},\begin{pmatrix}10&9\\2&10\end{pmatrix},\begin{pmatrix}10&2\\9&10\end{pmatrix},\begin{pmatrix}10&8\\3&10\end{pmatrix},\begin{pmatrix}10&3\\8&10\end{pmatrix},\begin{pmatrix}10&4\\7&10\end{pmatrix},\begin{pmatrix}10&7\\4&10\end{pmatrix},$$
$$\left.\begin{pmatrix}10&6\\5&10\end{pmatrix},\begin{pmatrix}10&5\\6&10\end{pmatrix}\right\}$$

This is a finite field of order 121 under addition and multiplication of matrices modulo 11. It has characteristic 11 and it contains

$$F_{11}=\left\{\begin{pmatrix}0&0\\0&0\end{pmatrix},\begin{pmatrix}1&0\\0&1\end{pmatrix},\begin{pmatrix}2&0\\0&2\end{pmatrix},\begin{pmatrix}3&0\\0&3\end{pmatrix},\begin{pmatrix}4&0\\0&4\end{pmatrix},\begin{pmatrix}5&0\\0&5\end{pmatrix},\begin{pmatrix}6&0\\0&6\end{pmatrix},\begin{pmatrix}7&0\\0&7\end{pmatrix},\begin{pmatrix}8&0\\0&8\end{pmatrix},\begin{pmatrix}9&0\\0&9\end{pmatrix},\begin{pmatrix}10&0\\0&10\end{pmatrix}\right\}$$

Following same technique one can find finite matrix field of order p^2 for $p > 11$.

© 2016 Global Journals Inc. (US)

References Références Referencias

1. M. Artin, Algebra, Prentice Hall of India Private Limited, New Delhi, 2000.
2. T. W. Hungerford, Algebra, Springer-India, New Delhi 2005.
3. S. K. Pandey, Matrix Field of Finite and Infinite Order, Research Journal of Pure Algebra **5** (12), 214-216, 2015.

Notes

© 2016 Global Journals Inc. (US)

Global Journal of Science Frontier Research: F
Mathematics and Decision Sciences
Volume 16 Issue 1 Version 1.0 Year 2016
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Mixed Quadrature Rule for Double Integrals of Simpsons $\frac{1}{3}rd$ and Gauss Legendre Two Point Rule in Two Variables

By Saumya Ranjan Jena & Subash Chandra Mishra
KIIT University, India

Abstract- In this article we are concerned with mixed quadrature rule of higher degree of precision for double integrals with double variables. The rule is numerically tested taking some suitable texts and the error bound is determined.

Keywords: *mixed quadrature rule, maclaurin's theorem, error bound, degree of precision.*

GJSFR-F Classification : *MSC2010: 65D30, 65D32*



Strictly as per the compliance and regulations of :

© 2016. Saumya Ranjan Jena & Subash Chandra Mishra. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Ref

3. Jena, S.,Dash, P., (2014)"Approximaton of real definite integrals via hybrid quadrature domain" *IJSETR*, ,3(42),.

Mixed Quadrature Rule for Double Integrals of Simpsons $\frac{1}{3}rd$ and Gauss Legendre Two Point Rule in Two Variables

Saumya Ranjan Jena [α] & Subash Chandra Mishra [σ]

Abstract- In this article we are concerned with mixed quadrature rule of higher degree of precision for double integrals with double variables. The rule is numerically tested taking some suitable texts and the error bound is determined.

Keywords: *mixed quadrature rule, maclaurin's theorem, error bound, degree of precision.*

I. Introduction

A Mixed quadrature rule of higher degree of precision has been formed by [2], [3], [4], [5], [6], [7]. This rule is meant for single integral. In the same vein here we develop a mixed quadrature rule of degree of precision-7 for double integrals taking the convex combination of Simpson's $\frac{1}{3}$rd and Gauss-Legendre-2 point rule each of degree of precision 5. This paper is designed as follows. Section -2 contains formulation of quadrature of constituent rules and the corresponding errors in-2 variables. Section-3 is devoted to construction of mixed quadrature rule. The error analysis is done in section-4. In section-5 the rule is numerically verified by taking suitable examples. The conclusions are drawn in section-6.

II. Formulation of Quadrature Rule in Two Variables X and Y

For approximate evaluation of real definite integral

$$I(f)=\int_{-1}^{1}\int_{-1}^{1} f(x,y)dxdy \tag{2.1}$$

The Simpson's $\frac{1}{3}$ *rd rule is*

$$I(f)\cong R_{s\frac{1}{3}}(f)=\frac{1}{9}\begin{bmatrix}\{f(-1,-1)+4f(0,-1)+f(1,-1)\}\\+4\{f(-1,0)+4f(0,0)+f(1,0)\}\\+\{f(-1,1)+4f(0,1)+f(1,1)\}\end{bmatrix} \tag{2.1}$$

And the Gauss-Legendre's two point rule is

Author α σ: Department of Mathematics, School of Applied Sciences, KIIT University, Bhubaneswar, Odisha, India.
e-mails: saumyafma@kiit.ac.in, subashmishra2010@gmail.com

© 2016 Global Journals Inc. (US)

$$I(f) \cong R_{GL2}(f) = f\left(-\frac{1}{\sqrt{3}},-\frac{1}{\sqrt{3}}\right)+ f\left(-\frac{1}{\sqrt{3}},\frac{1}{\sqrt{3}}\right)+ f\left(\frac{1}{\sqrt{3}},-\frac{1}{\sqrt{3}}\right)+ f\left(\frac{1}{\sqrt{3}},\frac{1}{\sqrt{3}}\right) \tag{2.3}$$

where each rule of equation (2.2) and equation (2.3) is of precision 3.

Hence

$$I(f) = R_{s\frac{1}{3}}(f) + E_{s\frac{1}{3}}(f) \tag{2.4}$$

$$I(f) = R_{GL2}(f) + E_{GL2}(f) \tag{2.5}$$

Where $E_{S\frac{1}{3}}(f)$ and $E_{GL2}(f)$ are error in approximating the integrals $I(f)$ by equation (2.2) and equation (2.3) respectively.

Now assuming $f(x,y)$ to be sufficiently differentiable in $-1\le x\le 1, -1\le y\le 1$

We can write equation (2.1) using Maclaurin's expansion

$$I(f)=\int_{-1}^{1}\int_{-1}^{1}\left[\begin{array}{l} f(0,0)+\{x\ f_{1,0}(0,0)+y\ f_{0,1}(0,0)\} \\ +\frac{1}{2!}\{x^2\ f_{2,0}(0,0)+2xy\ f_{1,1}(0,0)+y^2\ f_{0,2}(0,0)\} \\ +\frac{1}{3!}\{x^3\ f_{3,0}(0,0)+3x^2y\ f_{2,1}(0,0)+3xy^2\ f_{1,2}(0,0)+y^3\ f_{0,3}(0,0)\} \\ +\frac{1}{4!}\left\{\begin{array}{l} x^4\ f_{4,0}(0,0)+4x^3y\ f_{3,1}(0,0)+6x^2y^2\ f_{2,2}(0,0) \\ +4xy^3\ f_{1,3}(0,0)+y^4\ f_{0,4}(0,0) \end{array}\right\} \\ +\frac{1}{5!}\left\{\begin{array}{l} x^5\ f_{5,0}(0,0)+5x^4y\ f_{4,1}(0,0)+10x^3y^2\ f_{3,2}(0,0) \\ +10x^2y^3\ f_{2,3}(0,0)+5xy^4\ f_{1,4}(0,0)+y^5\ f_{0,5}(0,0) \end{array}\right\} \\ +\frac{1}{6!}\left\{\begin{array}{l} x^6\ f_{6,0}(0,0)+6x^5y\ f_{5,1}(0,0)+15x^4y^2\ f_{4,2}(0,0) \\ +20x^3y^3\ f_{3,3}(0,0)+15x^2y^4\ f_{2,4}(0,0) \\ +6xy^5\ f_{1,5}(0,0)+y^6\ f_{0,6}(0,0) \end{array}\right\} \end{array}\right] dxdy \tag{2.6}$$

Integrating equation (2.6) we have

$$I(f) = 4f_{0,0}(0,0)+\frac{2}{3}\left[f_{2,0}(0,0)+ f_{0,2}(0,0)\right]+\frac{1}{30}\left[f_{4,0}(0,0)+ f_{0,4}(0,0)\right]$$
$$+\frac{1}{9}f_{2,2}(0,0)+\frac{1}{180}\left[f_{4,2}(0,0)+ f_{2,4}(0,0)\right]+\frac{4}{7!}\left[f_{6,0}(0,0)+ f_{0,6}(0,0)\right]+.... \tag{2.7}$$

Again using Maclaurin's expansion

$$f(-1,-1) = f_{0,0}-\left[f_{1,0}(0,0)+ f_{0,1}(0,0)\right]+\frac{1}{2!}\left[f_{2,0}(0,0)+2f_{1,1}(0,0)+ f_{0,2}(0,0)\right]$$
$$-\frac{1}{3!}\left[f_{3,0}(0,0)+3f_{2,1}(0,0)+3f_{1,2}(0,0)+ f_{0,3}(0,0)\right]$$
$$+\frac{1}{4!}\left[f_{4,0}(0,0)+4f_{3,1}(0,0)+6f_{2,2}(0,0)+4f_{1,3}(0,0)+ f_{0,4}(0,0)\right]$$

Notes

© 2016 Global Journals Inc. (US)

$$-\frac{1}{5!}\left[f_{5,0}(0,0)+5f_{4,1}(0,0)+10f_{3,2}(0,0)+10f_{2,3}(0,0)+5f_{1,4}(0,0)+f_{0,5}(0,0)\right]$$

$$+\frac{1}{6!}\left[f_{6,0}(0,0)+6f_{5,1}(0,0)+15f_{4,2}(0,0)+20f_{3,3}(0,0)+15f_{2,4}(0,0)+6f_{1,5}(0,0)+f_{0,6}(0,0)\right]$$

$$+\ldots\ldots\ldots\ldots \tag{2.8}$$

Notes

Using equation (2.8) and $f(0,-1), f(1,-1), f(-1,0), f(1,0), f(0,1), f(-1,1)$ *and* $f(1,1)$ in equation (2.2)

$$R_{S\frac{1}{3}}(f)=4f_{0,0}(0,0)+\frac{2}{3}\left\{f_{2,0}(0,0)+f_{0,2}(0,0)\right\}+\frac{1}{18}\left\{f_{4,0}(0,0)+f_{0,4}(0,0)\right\}$$

$$+\frac{1}{9}f_{2,2}(0,0)+\frac{1}{108}\left\{f_{4,2}(0,0)+f_{2,4}(0,0)\right\}+\frac{11}{9\times 6!}\left\{f_{6,0}(0,0)+f_{0,6}(0,0)\right\} \tag{2.9}$$

a) Error in Simpson's $\frac{1}{3}$ *rd rule:*

Using equation (2.6), equation (2.9) in equation (2.4) error associated with Simpson's $\frac{1}{3}rd$ rule is

$$E_{S\frac{1}{3}}(f)=I(f)-R_{S\frac{1}{3}}(f)$$

$$E_{S\frac{1}{3}}(f)=-\frac{1}{45}\left\{f_{4,0}(0,0)+f_{0,4}(0,0)\right\}-\frac{1}{270}\left\{f_{4,2}(0,0)+f_{2,4}(0,0)\right\}-$$

$$\frac{41}{63\times 6!}\left\{f_{6,0}(0,0)+f_{0,6}(0,0)\right\} \tag{2.10}$$

As the error contains at least fourth derivative of the integral function, it vanishes for all polynomials of degree ≤ 3 in x and y. Thus the formula becomes exact for all polynomials of degree up to 3 i.e. degree of precision of the formula is three.

Similarly using Maclaurin's expansion

We have

$$f\left(-\frac{1}{\sqrt{3}},-\frac{1}{\sqrt{3}}\right)$$

$$=f(0,0)-\frac{1}{\sqrt{3}}\left[f_{1,0}(0,0)+f_{0,1}(0,0)\right]$$

$$+\frac{1}{2!}\left[\left(-\frac{1}{\sqrt{3}}\right)^2 f_{2,0}(0,0)+2\left(-\frac{1}{\sqrt{3}}\right)^2 f_{1,1}(0,0)+\left(-\frac{1}{\sqrt{3}}\right)^2 f_{0,2}(0,0)\right]$$

$$+\frac{1}{3!}\left[\left(-\frac{1}{\sqrt{3}}\right)^3 f_{3,0}(0,0)+3\left(-\frac{1}{\sqrt{3}}\right)^3 f_{2,1}(0,0)+3\left(-\frac{1}{\sqrt{3}}\right)^3 f_{1,2}(0,0)+\left(-\frac{1}{\sqrt{3}}\right)^3 f_{0,3}(0,0)\right]$$

© 2016 Global Journals Inc. (US)

$$+\frac{1}{4!}\left[\begin{array}{l}\left(-\frac{1}{\sqrt{3}}\right)^4 f_{4,0}(0,0)+4\left(-\frac{1}{\sqrt{3}}\right)^4 f_{3,1}(0,0)+6\left(-\frac{1}{\sqrt{3}}\right)^4 f_{2,2}(0,0)\\ +4\left(-\frac{1}{\sqrt{3}}\right)^4 f_{1,3}(0,0)+\left(-\frac{1}{\sqrt{3}}\right)^4 f_{0,4}(0,0)\end{array}\right]$$

$$+\frac{1}{5!}\left[\begin{array}{l}\left(-\frac{1}{\sqrt{3}}\right)^5 f_{5,0}(0,0)+5\left(-\frac{1}{\sqrt{3}}\right)^5 f_{4,1}(0,0)+10\left(-\frac{1}{\sqrt{3}}\right)^5 f_{3,2}(0,0)\\ +10\left(-\frac{1}{\sqrt{3}}\right)^5 f_{2,3}(0,0)+5\left(-\frac{1}{\sqrt{3}}\right)^5 f_{1,4}(0,0)+\left(-\frac{1}{\sqrt{3}}\right)^5 f_{0,5}(0,0)\end{array}\right]$$

$$+\frac{1}{6!}\left[\begin{array}{l}\left(-\frac{1}{\sqrt{3}}\right)^6 f_{6,0}(0,0)+6\left(-\frac{1}{\sqrt{3}}\right)^6 f_{5,1}(0,0)+15\left(-\frac{1}{\sqrt{3}}\right)^6 f_{4,2}(0,0)\\ +20\left(-\frac{1}{\sqrt{3}}\right)^6 f_{3,3}(0,0)+15\left(-\frac{1}{\sqrt{3}}\right)^6 f_{2,4}(0,0)\\ +6\left(-\frac{1}{\sqrt{3}}\right)^6 f_{1,5}(0,0)++\left(-\frac{1}{\sqrt{3}}\right)^6 f_{0,6}(0,0)\end{array}\right]\ldots\ldots \quad (2.11)$$

Now $f\left(-\frac{1}{\sqrt{3}},-\frac{1}{\sqrt{3}}\right), f\left(\frac{1}{\sqrt{3}},-\frac{1}{\sqrt{3}}\right), f\left(\frac{1}{\sqrt{3}},\frac{1}{\sqrt{3}}\right)$, *can be found out same ways as* (2.10)

Now substituting the value of

$$f\left(-\frac{1}{\sqrt{3}},-\frac{1}{\sqrt{3}}\right), f\left(\frac{1}{\sqrt{3}},-\frac{1}{\sqrt{3}}\right), f\left(\frac{1}{\sqrt{3}},-\frac{1}{\sqrt{3}}\right), f\left(\frac{1}{\sqrt{3}},\frac{1}{\sqrt{3}}\right)$$

in equation (2.3)

$$R_{GL2}(f)=4f(0,0)+\frac{2}{3}\left[f_{2,0}(0,0)+f_{0,2}(0,0)\right]+\frac{1}{9\times 3!}\left[f_{4,0}(0,0)+f_{0,4}(0,0)\right]$$

$$+\frac{20}{9\times 6!}\left[f_{4,2}(0,0)+f_{2,4}(0,0)\right]+\frac{1}{9}f_{2,2}(0,0)+\frac{4}{27\times 6!}\left[f_{6,0}(0,0)+f_{0,6}(0,0)\right] \quad (2.12)$$

b) Error in Gauss-Legendre's two point rule

Now the error associated with the Gauss- Legendre 2-point rule is obtained substituting equation (2.6) and (2.11) in equation (2.5)

$$E_{GL2}(f)=I(f)-R_{GL2}(f)$$

$$=\frac{2}{135}\left[f_{4,0}(0,0)+f_{0,4}(0,0)\right]+\frac{1}{405}\left[f_{4,2}(0,0)+f_{2,4}(0,0)\right]+\frac{1}{1701}\left[f_{6,0}(0,0)+f_{0,6}(0,0)\right] \quad (2.13)$$

© 2016 Global Journals Inc. (US)

Notes

In this case also, the error contains at least fourth derivative of the integral function. Thus the degree of the precision is 3.

III. Mixed Quadrature Rule

Now multiplying (2) by equation (2.10) and (3) by equation (2.13) and adding them, we get

$$I(f)=\frac{1}{5}\left[2R_{S\frac{1}{3}}(f)+3R_{GL2}(f)\right]+\frac{1}{5}\left[2E_{S\frac{1}{3}}(f)+3E_{GL2}(f)\right] \tag{3.1}$$

Where

$$I_{mix}(f)=R_{S\frac{1}{3}GL2}(f)+E_{S\frac{1}{3}GL2}(f) \tag{3.2}$$

Where

$$R_{S\frac{1}{3}GL2}(f)=\frac{1}{5}\left[2R_{S\frac{1}{3}}(f)+3R_{GL2}(f)\right] \tag{3.3}$$

Where $R_{S\frac{1}{3}GL2}(f)$ and $E_{S\frac{1}{3}GL2}(f)$ are mixed quadrature rule and its error obtained by Simpson's $\frac{1}{3}rd$ and Gauss-Legendre's 2 point rule respectively.

The truncation error generated by this approximation is given by

$$E_{S\frac{1}{3}GL2}=\frac{1}{5}\left[2E_{S\frac{1}{3}}(f)+3R_{GL2}(f)\right]=-\frac{1}{189\times 5!}\left[f_{6,0}(0,0)+f_{0,6}(0,0)\right] \tag{3.4}$$

The rule (3.3) may be called as mixed quadrature rule integrate exactly all polynomial of degree ≤ 5 in x and y.

IV. Error Analysis

a) Theorem 4.1

Let $f(x,y)$ be sufficiently differentiable function in the closed interval $[-1,1]$. The bounds of truncatinal error $E_{S\frac{1}{3}GL2}(f)$ associated with the rule $R_{S\frac{1}{3}GL2}(f)$ is given by

$$\left|E_{S\frac{1}{3}GL2}(f)\right|\cong\frac{1}{189\times 5!}\left[f_{6,0}(0,0)+f_{0,6}(0,0)\right]$$

Proof:

The proof obviously follows from the equation (3.4)

Theorem 4.2

The bounds for the truncational error $\left|E_{S\frac{1}{3}GL2}(f)\right|\leq\frac{2M}{225}\left|\eta_2-\eta_1\right|$

Where $\eta_1,\eta_2\in[-1,1]$

Where $M=Max\left|f_{6,0}(0,0)+f_{0,6}(0,0)\right|,\ -1\leq x\leq 1,\ -1\leq y\leq 1$

© 2016 Global Journals Inc. (US)

Proof:
We have

$$E_{S\frac{1}{3}}(f) = -\frac{1}{45}\left|f_{4,0}(n_1) + f_{0,4}(n_1)\right|$$

$$E_{GL2}(f) = \frac{2}{135}\left|f_{4,0}(n_2) + f_{0,4}(n_2)\right|$$

Where

$$n_1, n_2 \in [-1,1]$$

Hence

$$E_{S\frac{1}{3}GL2}(f) = \frac{1}{5}\left|2E_{s\frac{1}{3}}(f) + 3E_{GL2}(f)\right|$$

$$= \frac{2}{225}\left[f_{4,0}(n_2,0) + f_{0,4}(0,n_2) - f_{4,0}(n_1,0) - f_{0,4}(0,n_1)\right]$$

$$= \frac{2}{225}\int_{n_1}^{n_2}\int_{n_1}^{n_2}\left[f_{5,0}(x,*) + f_{0,5}(*,y)\right]$$

Hence

$$\left|E_{S\frac{1}{3}GL2}(f)\right| \leq \frac{2M|n_1 - n_2|}{225}$$

Where

$$M = \max\left|f_{5,0}(x,*) + f_{0,5}(*,y)\right| \qquad -1 \leq x \leq 1, \quad -1 \leq y \leq 1$$

Which, gives only the truncation error bound on n_1, n_2 are closed to each other.

b) Corollary

The error bound for the truncated error $\left|E_{S\frac{1}{3}GL2}(f)\right| \leq \frac{4M}{225}$

When $|n_1 - n_2| \leq 2$ [1]

V. Numerical Verification

The exact value of the integrals

$$I_1 = \int_{-1}^{1}\int_{-1}^{1} e^{x+y}dxdy = 5.524391382167262$$

$$I_1 = \int_{-1}^{1}\int_{-1}^{1} e^{-\left(x^2+y^2\right)}dxdy = 2.230985141404134$$

$$I_3 = \int_{0}^{1}\int_{0}^{1} \frac{\sin^2(x+y)}{(x+y)}dxdy = 0.613260369981918$$

Notes

© 2016 Global Journals Inc. (US)

Notes

Table-1 : (Comparison of exact result with approximate result)

Exact	$R_{S\frac{1}{3}}(f)$	$R_{GL2}(f)$	$R_{S\frac{1}{3}GL2}(f)$	$E_{S\frac{1}{3}GL2}(f)$
I_1 =5.524391382167262	5.577469135802469	5.488065843621398	5.523827160493826	0.00056421673436
I_3 =2.230985141404134	2.259259259259260	2.024691358024692	2.238518518518519	0.007533377114385
I_3 =0.613260369981918	0.612749599379052	0.609491314229328	0.610769318605380	0.002491051376538

VI. Conclusion

From the Table -1, we find that

$$\left|E_{S\frac{1}{3}}(f)\right| \leq \left|E_{GL2}(f)\right| \qquad (i)$$

$$\left|E_{S\frac{1}{3}GL2}(f)\right| \leq \left|E_{S\frac{1}{3}}(f)\right| \qquad (ii)$$

From (i) and (ii)

$$\left|E_{S\frac{1}{3}GL2}(f)\right| \leq \left|E_{S\frac{1}{3}}(f)\right| \leq \left|E_{GL2}(f)\right|$$

It is evident that the mixed quadrature rule $R_{S\frac{1}{3}GL2}(f)$ of degree of precision 5 provides us better result than constituent rule $R_{S\frac{1}{3}}(f), R_{GL2}(f)$ each of degree of precision three. Hence, the mixed quadrature rule is more efficient and numerically better convergent then exact result.

References Références Referencias

1. Conte ,S. Boor, C.De. (1980), "*Elementary Numerical Analysis*", Tata Mac-Graw Hill.
2. Jena,S.R. and Dash,R.B., (2009) "Mixed quadrature of real definite integral over triangles" *PAJM*, ,3, pp.119-124.
3. Jena, S.,Dash, P., (2014)"Approximaton of real definite integrals via hybrid quadrature domain" *IJSETR*, ,3(42),.
4. Kreyszig, E.,(2012), *"Advanced Engg.Mathematics,"* Wiley India.
5. Ma . J. Rokhlin, V.,Wandzura S., (1996)" *Generalised* Gaussian quadrature rules for systems for systems of arbitrary functions", *SIAM Journal of Numerical Analysis* ; 33(3) 971-996.
6. Das R. N. and Pradhan G., (1996) "A mixed quadrature rule for approximate evaluation of real definite integral". *Int. J. Math. Educ. Sci.Technol*, , 27(2), 279-283.
7. Jena, S.R. and Dash, R.B., (2011) "Study of approximate value of real definite integral by mixed quadrature rule obtained from Richardson extrapolation", *IJCSM* 3(1),pp.47-53.

© 2016 Global Journals Inc. (US)

Global Journals Inc. (US) Guidelines Handbook 2016

www.GlobalJournals.org

FELLOWS

FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (FARSS)

Global Journals Incorporate (USA) is accredited by Open Association of Research Society (OARS), U.S.A and in turn, awards "FARSS" title to individuals. The 'FARSS' title is accorded to a selected professional after the approval of the Editor-in-Chief/Editorial Board Members/Dean.



- The "FARSS" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSS or William Walldroff, M.S., FARSS.

FARSS accrediting is an honor. It authenticates your research activities. After recognition as FARSB, you can add 'FARSS' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, and Visiting Card etc.

The following benefits can be availed by you only for next three years from the date of certification:

FARSS designated members are entitled to avail a 40% discount while publishing their research papers (of a single author) with Global Journals Incorporation (USA), if the same is accepted by Editorial Board/Peer Reviewers. If you are a main author or co-author in case of multiple authors, you will be entitled to avail discount of 10%.

Once FARSB title is accorded, the Fellow is authorized to organize a symposium/seminar/conference on behalf of Global Journal Incorporation (USA). The Fellow can also participate in conference/seminar/symposium organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent.





You may join as member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer. In addition, it is also desirable that you should organize seminar/symposium/conference at least once.

We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



© Copyright by Global Journals Inc.(US) | Guidelines Handbook

The FARSS can go through standards of OARS. You can also play vital role if you have any suggestions so that proper amendment can take place to improve the same for the benefit of entire research community.



As FARSS, you will be given a renowned, secure and free professional email address with 100 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.



The FARSS will be eligible for a free application of standardization of their researches. Standardization of research will be subject to acceptability within stipulated norms as the next step after publishing in a journal. We shall depute a team of specialized research professionals who will render their services for elevating your researches to next higher level, which is worldwide open standardization.



The FARSS member can apply for grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.
Once you are designated as FARSS, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria. After certification of all your credentials by OARS, they will be published on your Fellow Profile link on website https://associationofresearch.org which will be helpful to upgrade the dignity.

The FARSS members can avail the benefits of free research podcasting in Global Research Radio with their research documents. After publishing the work, (including published elsewhere worldwide with proper authorization) you can upload your research paper with your recorded voice or you can utilize chargeable services of our professional RJs to record your paper in their voice on request.



The FARSS member also entitled to get the benefits of free research podcasting of their research documents through video clips. We can also streamline your conference videos and display your slides/ online slides and online research video clips at reasonable charges, on request.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook



The FARSS is eligible to earn from sales proceeds of his/her researches/reference/review Books or literature, while publishing with Global Journals. The FARSS can decide whether he/she would like to publish his/her research in a closed manner. In this case, whenever readers purchase that individual research paper for reading, maximum 60% of its profit earned as royalty by Global Journals, will be credited to his/her bank account. The entire entitled amount will be credited to his/her bank account exceeding limit of minimum fixed balance. There is no minimum time limit for collection. The FARSS member can decide its price and we can help in making the right decision.

The FARSS member is eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get remuneration of 15% of author fees, taken from the author of a respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account.



MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (MARSS)

The ' MARSS ' title is accorded to a selected professional after the approval of the Editor-in-Chief / Editorial Board Members/Dean.



The "MARSS" is a dignified ornament which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., MARSS or William Walldroff, M.S., MARSS.

MARSS accrediting is an honor. It authenticates your research activities. After becoming MARSS, you can add 'MARSS' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, Visiting Card and Name Plate etc.

The following benefitscan be availed by you only for next three years from the date of certification.



MARSS designated members are entitled to avail a 25% discount while publishing their research papers (of a single author) in Global Journals Inc., if the same is accepted by our Editorial Board and Peer Reviewers. If you are a main author or co-author of a group of authors, you will get discount of 10%.

As MARSS, you will be given a renowned, secure and free professional email address with 30 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.



© Copyright by Global Journals Inc.(US) | Guidelines Handbook



We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

The MARSS member can apply for approval, grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.





Once you are designated as MARSS, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria.

It is mandatory to read all terms and conditions carefully.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

AUXILIARY MEMBERSHIPS

Institutional Fellow of Global Journals Incorporation (USA)-OARS (USA)

Global Journals Incorporation (USA) is accredited by Open Association of Research Society, U.S.A (OARS) and in turn, affiliates research institutions as "Institutional Fellow of Open Association of Research Society" (IFOARS).



The "FARSC" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSC or William Walldroff, M.S., FARSC.

The IFOARS institution is entitled to form a Board comprised of one Chairperson and three to five board members preferably from different streams. The Board will be recognized as "Institutional Board of Open Association of Research Society"-(IBOARS).

The Institute will be entitled to following benefits:

The IBOARS can initially review research papers of their institute and recommend them to publish with respective journal of Global Journals. It can also review the papers of other institutions after obtaining our consent. The second review will be done by peer reviewer of Global Journals Incorporation (USA) The Board is at liberty to appoint a peer reviewer with the approval of chairperson after consulting us.

The author fees of such paper may be waived off up to 40%.

The Global Journals Incorporation (USA) at its discretion can also refer double blind peer reviewed paper at their end to the board for the verification and to get recommendation for final stage of acceptance of publication.





The IBOARS can organize symposium/seminar/conference in their country on behalf of Global Journals Incorporation (USA)-OARS (USA). The terms and conditions can be discussed separately.

The Board can also play vital role by exploring and giving valuable suggestions regarding the Standards of "Open Association of Research Society, U.S.A (OARS)" so that proper amendment can take place for the benefit of entire research community. We shall provide details of particular standard only on receipt of request from the Board.



The board members can also join us as Individual Fellow with 40% discount on total fees applicable to Individual Fellow. They will be entitled to avail all the benefits as declared. Please visit Individual Fellow-sub menu of GlobalJournals.org to have more relevant details.

© Copyright by Global Journals Inc.(US) | Guidelines Handbook

We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



After nomination of your institution as "Institutional Fellow" and constantly functioning successfully for one year, we can consider giving recognition to your institute to function as Regional/Zonal office on our behalf.
The board can also take up the additional allied activities for betterment after our consultation.

The following entitlements are applicable to individual Fellows:



Open Association of Research Society, U.S.A (OARS) By-laws states that an individual Fellow may use the designations as applicable, or the corresponding initials. The Credentials of individual Fellow and Associate designations signify that the individual has gained knowledge of the fundamental concepts. One is magnanimous and proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice.

Open Association of Research Society (US)/ Global Journals Incorporation (USA), as described in Corporate Statements, are educational, research publishing and professional membership organizations. Achieving our individual Fellow or Associate status is based mainly on meeting stated educational research requirements.



Disbursement of 40% Royalty earned through Global Journals : Researcher = 50%, Peer Reviewer = 37.50%, Institution = 12.50% E.g. Out of 40%, the 20% benefit should be passed on to researcher, 15 % benefit towards remuneration should be given to a reviewer and remaining 5% is to be retained by the institution.

We shall provide print version of 12 issues of any three journals [as per your requirement] out of our 38 journals worth $ 2376 USD.

Other:

The individual Fellow and Associate designations accredited by Open Association of Research Society (US) credentials signify guarantees following achievements:

- The professional accredited with Fellow honor, is entitled to various benefits viz. name, fame, honor, regular flow of income, secured bright future, social status etc.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

- In addition to above, if one is single author, then entitled to 40% discount on publishing research paper and can get 10%discount if one is co-author or main author among group of authors.
- The Fellow can organize symposium/seminar/conference on behalf of Global Journals Incorporation (USA) and he/she can also attend the same organized by other institutes on behalf of Global Journals.
- The Fellow can become member of Editorial Board Member after completing 3yrs.
- The Fellow can earn 60% of sales proceeds from the sale of reference/review books/literature/publishing of research paper.
- Fellow can also join as paid peer reviewer and earn 15% remuneration of author charges and can also get an opportunity to join as member of the Editorial Board of Global Journals Incorporation (USA)
- • This individual has learned the basic methods of applying those concepts and techniques to common challenging situations. This individual has further demonstrated an in–depth understanding of the application of suitable techniques to a particular area of research practice.

Note :

"

- In future, if the board feels the necessity to change any board member, the same can be done with the consent of the chairperson along with anyone board member without our approval.
- In case, the chairperson needs to be replaced then consent of 2/3rd board members are required and they are also required to jointly pass the resolution copy of which should be sent to us. In such case, it will be compulsory to obtain our approval before replacement.
- In case of "Difference of Opinion [if any]" among the Board members, our decision will be final and binding to everyone.

"

Process of submission of Research Paper

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (*.DOC,*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission.Online Submission: There are three ways to submit your paper:

(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.

(II) Choose corresponding Journal.

(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.

(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.

(C) If these two are not conveninet , and then email the paper directly to dean@globaljournals.org.

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Preferred Author Guidelines

MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11'"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

You can use your own standard format also.
Author Guidelines:

1. General,

2. Ethical Guidelines,

3. Submission of Manuscripts,

4. Manuscript's Category,

5. Structure and Format of Manuscript,

6. After Acceptance.

1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global

Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.

2) Drafting the paper and revising it critically regarding important academic content.

3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.

Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.



To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

5.STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also.Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

Papers: These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

(a)Title should be relevant and commensurate with the theme of the paper.

(b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.

(c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.

(d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.

(e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.

(f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;

(g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.

(h) Brief Acknowledgements.

(i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.

The Editorial Board reserves the right to make literary corrections and to make suggestions to improve briefness.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

Format

Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than 1.4 × 10-3 m3, or 4 mm somewhat than 4 × 10-3 m. Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

Abstract, used in Original Papers and Reviews:

Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art.A few tips for deciding as strategically as possible about keyword search:

- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

Acknowledgements: Please make these as concise as possible.

References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

Tables, Figures and Figure Legends

Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.

Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.

Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.

Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.

6. AFTER ACCEPTANCE

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

6.1 Proof Corrections

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

www.adobe.com/products/acrobat/readstep2.html. This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at dean@globaljournals.org within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

6.3 Author Services

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

6.4 Author Material Archive Policy

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

6.5 Offprint and Extra Copies

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: editor@globaljournals.org .



Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

1. Choosing the topic: In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

2. Evaluators are human: First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

3. Think Like Evaluators: If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

4. Make blueprints of paper: The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

5. Ask your Guides: If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

6. Use of computer is recommended: As you are doing research in the field of Computer Science, then this point is quite obvious.

7. Use right software: Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

8. Use the Internet for help: An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

9. Use and get big pictures: Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

10. Bookmarks are useful: When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

11. Revise what you wrote: When you write anything, always read it, summarize it and then finalize it.

12. Make all efforts: Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

13. Have backups: When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

14. Produce good diagrams of your own: Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

15. Use of direct quotes: When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

16. Use proper verb tense: Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

17. Never use online paper: If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

18. Pick a good study spot: To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

19. Know what you know: Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

20. Use good quality grammar: Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

21. Arrangement of information: Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

22. Never start in last minute: Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

23. Multitasking in research is not good: Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

24. Never copy others' work: Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

25. Take proper rest and food: No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

26. Go for seminars: Attend seminars if the topic is relevant to your research area. Utilize all your resources.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

27. Refresh your mind after intervals: Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

28. Make colleagues: Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

29. Think technically: Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

30. Think and then print: When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

31. Adding unnecessary information: Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

32. Never oversimplify everything: To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

33. Report concluded results: Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

34. After conclusion: Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium though which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

Informal Guidelines of Research Paper Writing

Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.

Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

General style:

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

· Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

· Use standard writing style including articles ("a", "the," etc.)

· Keep on paying attention on the research topic of the paper

· Use paragraphs to split each significant point (excluding for the abstract)

· Align the primary line of each section

· Present your points in sound order

· Use present tense to report well accepted

· Use past tense to describe specific results

· Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives

· Shun use of extra pictures - include only those figures essential to presenting results

Title Page:

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript--must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for briefness. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

Approach:

- Single section, and succinct
- As a outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an conceptual must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.

- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

Procedures (Methods and Materials):

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

Methods:

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

Approach:

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

What to keep away from

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

Results:

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently.You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

Segment Draft and Final Research Paper: You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptive of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.

- Do not give permission to anyone else to "PROOFREAD" your manuscript.

- Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Criterion for Grading a Research Paper (Compilation) by Global Journals Inc. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

Topics	Grades		
	A-B	**C-D**	**E-F**
Abstract	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form Above 200 words	No specific data with ambiguous information Above 250 words
Introduction	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
Methods and Procedures	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
Result	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
Discussion	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
References	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring

INDEX

A

Analogue · 11, 13

B

Berezansky · 15, 17, 32
Biorthogonal · 15, 17, 22, 24, 26, 31

D

Decoloratus · 1, 5, 7, 9

F
F
Fonctionelles · 31

G

Guseinov · 3, 12
Gaussian · 15, 17, 21, 22, 31,

H

Hermitian · 20
Hilbert · 24, 26, 34
Hinton · 3, 11, 12

K
K
Kaymakcalan · 3, 12
Kondratiev · 17, 31

L

Legendre · 43, 44, 49, 50
Lyapunov · 1, 2, 3, 11, 12, 13

M

Maclaurin's · 43

N

Naukova · 31

S

Samoilenko · 15, 32
Stabilit · 11



Global Journal of Science Frontier Research

Visit us on the Web at www.GlobalJournals.org | www.JournalofScience.org
or email us at helpdesk@globaljournals.org

ISSN 9755896



© Global Journals