Online ISSN : 2249-4626
Print ISSN : 0975-5896
DOI : 10.17406/GJSFR

Global Journal

of Science Frontier Research: A

Physics and Space Science



Highlights

- New Concept of Confinement
- Planckeon Origin of Dark Energy
- EPR Correlation in Cosmology
- Analytical Determination of Subsurface

Discovering Thoughts, Inventing Future

VOLUME 16 ISSUE 3 VERSION 1.0

© 2001-2016 by Global Journal of Science Frontier Research, USA

Volume 16 Issue 15 was originally issued as A3

Global Journal of Science Frontier Research: A
Physics & Space Science

Volume 16 Issue 15
Originally issued as A3

Global Journal of Science Frontier Research: A
Physics & Space Science

Volume 16 Issue 3 (Ver. 1.0)

Open Association of Research Society

© Global Journal of Science Frontier Research. 2016.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Science Frontier Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Science Frontier Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Science Frontier Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website http://globaljournals.us/terms-and-condition/menu-id-1463/

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089
License No.: 42125/022010/1186
Registration No.: 430374
Import-Export Code: 1109007027
Employer Identification Number (EIN):
USA Tax ID: 98-0673427

Global Journals Inc.

*(A Delaware USA Incorporation with "Good Standing"; **Reg. Number: 0423089**)*

Sponsors: Open Association of Research Society
Open Scientific Standards

Publisher's Headquarters office

Global Journals® Headquarters
945th Concord Streets,
Framingham Massachusetts Pin: 01701,
United States of America
USA Toll Free: +001-888-839-7392
USA Toll Free Fax: +001-888-839-7392

Offset Typesetting

Global Journals Incorporated
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,
Pin: CR9 2ER, United Kingdom

Packaging & Continental Dispatching

Global Journals
E-3130 Sudama Nagar, Near Gopur Square,
Indore, M.P., Pin:452009, India

Find a correspondence nodal officer near you

To find nodal officer of your country, please email us at *local@globaljournals.org*

eContacts

Press Inquiries: *press@globaljournals.org*
Investor Inquiries: *investors@globaljournals.org*
Technical Support: *technology@globaljournals.org*
Media & Releases: *media@globaljournals.org*

Pricing (Including by Air Parcel Charges):

For Authors:
22 USD (B/W) & 50 USD (Color)
Yearly Subscription (Personal & Institutional):
200 USD (B/W) & 250 USD (Color)

Integrated Editorial Board
(Computer Science, Engineering, Medical, Management, Natural Science, Social Science)

John A. Hamilton,"Drew" Jr.,
Ph.D., Professor, Management
Computer Science and Software Engineering
Director, Information Assurance Laboratory
Auburn University

Dr. Henry Hexmoor
IEEE senior member since 2004
Ph.D. Computer Science, University at Buffalo
Department of Computer Science
Southern Illinois University at Carbondale

Dr. Osman Balci, Professor
Department of Computer Science
Virginia Tech, Virginia University
Ph.D.and M.S.Syracuse University, Syracuse, New York
M.S. and B.S. Bogazici University, Istanbul, Turkey

Yogita Bajpai
M.Sc. (Computer Science), FICCT
U.S.A.Email:
yogita@computerresearch.org

Dr. T. David A. Forbes
Associate Professor and Range Nutritionist
Ph.D. Edinburgh University - Animal Nutrition
M.S. Aberdeen University - Animal Nutrition
B.A. University of Dublin- Zoology

Dr. Wenying Feng
Professor, Department of Computing & Information Systems
Department of Mathematics
Trent University, Peterborough,
ON Canada K9J 7B8

Dr. Thomas Wischgoll
Computer Science and Engineering,
Wright State University, Dayton, Ohio
B.S., M.S., Ph.D.
(University of Kaiserslautern)

Dr. Abdurrahman Arslanyilmaz
Computer Science & Information Systems Department
Youngstown State University
Ph.D., Texas A&M University
University of Missouri, Columbia
Gazi University, Turkey

Dr. Xiaohong He
Professor of International Business
University of Quinnipiac
BS, Jilin Institute of Technology; MA, MS, PhD,. (University of Texas-Dallas)

Burcin Becerik-Gerber
University of Southern California
Ph.D. in Civil Engineering
DDes from Harvard University
M.S. from University of California, Berkeley & Istanbul University

Dr. Bart Lambrecht
Director of Research in Accounting and Finance Professor of Finance
Lancaster University Management School
BA (Antwerp); MPhil, MA, PhD (Cambridge)

Dr. Carlos García Pont
Associate Professor of Marketing
IESE Business School, University of Navarra
Doctor of Philosophy (Management), Massachusetts Institute of Technology (MIT)
Master in Business Administration, IESE, University of Navarra
Degree in Industrial Engineering, Universitat Politècnica de Catalunya

Dr. Fotini Labropulu
Mathematics - Luther College
University of Regina Ph.D., M.Sc. in Mathematics
B.A. (Honors) in Mathematics
University of Windso

Dr. Lynn Lim
Reader in Business and Marketing
Roehampton University, London
BCom, PGDip, MBA (Distinction), PhD, FHEA

Dr. Mihaly Mezei
ASSOCIATE PROFESSOR
Department of Structural and Chemical Biology, Mount Sinai School of Medical Center
Ph.D., Etvs Lornd University
Postdoctoral Training,
New York University

Dr. Söhnke M. Bartram
Department of Accounting and Finance Lancaster University Management School Ph.D. (WHU Koblenz)
MBA/BBA (University of Saarbrücken)

Dr. Miguel Angel Ariño
Professor of Decision Sciences
IESE Business School
Barcelona, Spain (Universidad de Navarra)
CEIBS (China Europe International Business School).
Beijing, Shanghai and Shenzhen
Ph.D. in Mathematics
University of Barcelona
BA in Mathematics (Licenciatura)
University of Barcelona

Philip G. Moscoso
Technology and Operations Management
IESE Business School, University of Navarra
Ph.D in Industrial Engineering and Management, ETH Zurich
M.Sc. in Chemical Engineering, ETH Zurich

Dr. Sanjay Dixit, M.D.
Director, EP Laboratories, Philadelphia VA Medical Center
Cardiovascular Medicine - Cardiac Arrhythmia
Univ of Penn School of Medicine

Dr. Han-Xiang Deng
MD., Ph.D
Associate Professor and Research Department Division of Neuromuscular Medicine
Davee Department of Neurology and Clinical Neuroscience Northwestern University Feinberg School of Medicine

Dr. Pina C. Sanelli
Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic Radiology
M.D., State University of New York at Buffalo,School of Medicine and Biomedical Sciences

Dr. Roberto Sanchez
Associate Professor
Department of Structural and Chemical Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University

Dr. Wen-Yih Sun
Professor of Earth and Atmospheric Sciences Purdue University Director National Center for Typhoon and Flooding Research, Taiwan
University Chair Professor
Department of Atmospheric Sciences, National Central University, Chung-Li, TaiwanUniversity Chair Professor
Institute of Environmental Engineering, National Chiao Tung University, Hsin-chu, Taiwan.Ph.D., MS The University of Chicago, Geophysical Sciences
BS National Taiwan University, Atmospheric Sciences
Associate Professor of Radiology

Dr. Michael R. Rudnick
M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and Hypertension Division (PMC)
Penn Medicine, University of Pennsylvania
Presbyterian Medical Center, Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of Internal Medicine

Dr. Bassey Benjamin Esu
B.Sc. Marketing; MBA Marketing; Ph.D Marketing
Lecturer, Department of Marketing, University of Calabar
Tourism Consultant, Cross River State Tourism Development Department
Co-ordinator, Sustainable Tourism Initiative, Calabar, Nigeria

Dr. Aziz M. Barbar, Ph.D.
IEEE Senior Member
Chairperson, Department of Computer Science
AUST - American University of Science & Technology
Alfred Naccash Avenue – Ashrafieh

President Editor (HON.)

Dr. George Perry, (Neuroscientist)
Dean and Professor, College of Sciences
Denham Harman Research Award (American Aging Association)
ISI Highly Cited Researcher, Iberoamerican Molecular Biology Organization
AAAS Fellow, Correspondent Member of Spanish Royal Academy of Sciences
University of Texas at San Antonio
Postdoctoral Fellow (Department of Cell Biology)
Baylor College of Medicine
Houston, Texas, United States

Chief Author (HON.)

Dr. R.K. Dixit
M.Sc., Ph.D., FICCT
Chief Author, India
Email: authorind@computerresearch.org

Dean & Editor-in-Chief (HON.)

Vivek Dubey(HON.)
MS (Industrial Engineering),
MS (Mechanical Engineering)
University of Wisconsin, FICCT
Editor-in-Chief, USA
editorusa@computerresearch.org

Sangita Dixit
M.Sc., FICCT
Dean & Chancellor (Asia Pacific)
deanind@computerresearch.org

Suyash Dixit
(B.E., Computer Science Engineering), FICCTT
President, Web Administration and
Development , CEO at IOSRD
COO at GAOR & OSS

Er. Suyog Dixit
(M. Tech), BE (HONS. in CSE), FICCT
SAP Certified Consultant
CEO at IOSRD, GAOR & OSS
Technical Dean, Global Journals Inc. (US)
Website: www.suyogdixit.com
Email: suyog@suyogdixit.com

Pritesh Rajvaidya
(MS) Computer Science Department
California State University
BE (Computer Science), FICCT
Technical Dean, USA
Email: pritesh@computerresearch.org

Luis Galárraga
J!Research Project Leader
Saarbrücken, Germany

Contents of the Issue

i. Copyright Notice
ii. Editorial Board Members
iii. Chief Author and Dean
iv. Contents of the Issue

1. Dynamic Interaction: A New Concept of Confinement. *1-9*
2. Bi-Directional EPR Correlation in Cosmology and Planckeon Origin of Dark Energy. *11-13*
3. The Dance of the Spinning Top. *15-27*
4. Analytical Determination of Subsurface Temperature using Two Layers Model in Part of Niger Delta Sedimentary Basin, Nigeria. *29-34*

v. Fellows
vi. Auxiliary Memberships
vii. Process of Submission of Research Paper
viii. Preferred Author Guidelines
ix. Index

Global Journal of Science Frontier Research: a
Physics and Space Science
Volume 16 Issue 3 Version 1.0 Year 2016
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Dynamic Interaction: A New Concept of Confinement

By Gabriel Barceló

Abstract- We propose new dynamic hypotheses to enhance our understanding of the behavior of the plasma in the reactor. In doing so, we put forward a profound revision of classical dynamics. After over thirty years studying rotational dynamics, we propose a new theory of dynamic interactions to better interpret nature in rotation. This new theory has been tested experimentally returning positive results, including by third parties.

Plasma rotation is an essential factor in the analysis of the turbulent transport of momentum in axisymmetric systems. In magnetic confinement fusion systems, the plasma circulates in the container at a constant movement, which we could define as rotation with respect to its walls. Notwithstanding, it has been shown that the plasma in the reactor can initiate spontaneous circular movement or rotation, without the need for any external dynamic momentum input. The theoretical development of this behavior is still under study, and the origin of this spontaneous intrinsic rotation is still unclear.

GJSFR-A Classification : FOR Code: 010204



Strictly as per the compliance and regulations of :

© 2016. Gabriel Barceló. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Dynamic Interaction: A New Concept of Confinement

Gabriel Barceló

Abstract- We propose new dynamic hypotheses to enhance our understanding of the behavior of the plasma in the reactor. In doing so, we put forward a profound revision of classical dynamics. After over thirty years studying rotational dynamics, we propose a new theory of dynamic interactions to better interpret nature in rotation. This new theory has been tested experimentally returning positive results, including by third parties.

Plasma rotation is an essential factor in the analysis of the turbulent transport of momentum in axisymmetric systems. In magnetic confinement fusion systems, the plasma circulates in the container at a constant movement, which we could define as rotation with respect to its walls. Notwithstanding, it has been shown that the plasma in the reactor can initiate spontaneous circular movement or rotation, without the need for any external dynamic momentum input. The theoretical development of this behavior is still under study, and the origin of this spontaneous intrinsic rotation is still unclear.

We suggest the exploring of a new type of dynamic confinement based on the Theory of Dynamic Interactions (TDI) and one that is compatible with magnetic confinement. Applying this criterion we are proposing would enable a twin physical-theoretical principle to isolate plasma and try to minimize its turbulence.

We suggest that these new dynamic hypotheses, which we hold applicable to particle systems accelerated by rotation, be used in the interpretation and design of fusion reactors. We believe that this proposal could, in addition to magnetic confinement, achieve confinement by simultaneous and compatible dynamic interaction.

I. Magnetic Confinement

On magneto hydrodynamic, the study of nonlinear physics associated with tokamak plasma turbulence and transport, allows us to infer the behavior of the reactor. For this goal, it is possible to use Gyrokinetics plasma turbulence simulation. The rotational nature of astrophysical plasmas has been also taken into account in its development [12], the goal is to have a reactor with a continuous process, which is what is required for its industrial development.

Current developments in our knowledge concerning the magnetic confinement of plasma for fusion defines a toroidal design of magnetic field force lines, which in successive toroidal transits travel over a surface topologically equivalent, in geometrical terms, to a torus. Indeed, this is the basis of the JET tokamak design and that of later reactors.[1]

On these reactors, magnetic confinement consists of containing material in a state of plasma inside an enclosure in which, in turn, a magnetic field has been created. The particles constituting the plasma do not enter into contact with the inner walls of the enclosure.

This method is based on the *Lorentz force*, which acts on any charged particle in motion in a magnetic field, by virtue of which it experiences a force perpendicular to the magnetic field vector and the displacement vector.

Fusion reactions are caused in this plasma that release energy, while at the same time a quantity of movement and angular momentum is moved or "rotated" and transported.

it has been confirmed that circulation in the experimental reactors is more turbulent than that which is supposed in stellar dynamic systems and, as we have pointed out above, this reduction of fluid uniformity can cause undesired losses in the system's power of confinement, or even, the possible cavitation of the physical container.

The speed of the plasma is an important physical field in the study of current laboratory systems, and particularly, in the improvement of its material isolation. The rotation profile can exercise considerable influence in the confinement time of the magnetically confined fluid. Profiles with a sharp radial velocity variation can cause a reduction of the turbulent transport and, therefore, could lead to improved confinement.

The study of plasma micro turbulence caused by the electromagnetic interactions of its particles is one of the current goals in an attempt to improve the performance of these reactors. Understanding this micro turbulence in plasma would enable its control and thus have a direct favorable impact on the building of more efficient fusion reactors.[2]

Author: Advanced Dynamics CB, Madrid, Spain.
e-mail: gestor@advanceddynamics.net

[1] Barceló, G.: Dynamic Interaction Confinement. World Journal of Nuclear Science and Technology Vol.4 No.4, October 29, 2014, DOI: 10.4236/wjnst.2014.44031.

[2] Barceló, G.: *Dynamic Interaction Confinement.* World Journal of Nuclear Science and Technology Vol.4 No.4, October 29, 2014DOI: 10.4236/wjnst.2014.44031.

© 2016 Global Journals Inc. (US)

II. Gyrokinetics Theory and Simulations

The Gyrokinetics theory is also applied to determine the behavior of the plasma in the to kamak fusion reactors.

Nonlinear Gyrokinetics equations play a fundamental role in our understanding of the long-time behavior of strongly magnetized plasmas. The foundations of modern nonlinear Gyrokinetics theory are based on three pillars:

1) *A Gyrokinetics Vlasov equation written in terms of a gyro center Hamiltonian,*
2) *With quadratic low-frequency ponderomotive like terms, a set of Gyrokinetics Maxwell Poisson-Ampère equations written in terms of the gyro center Vlasov distribution that contain low-frequency polarization Poisson and magnetization Ampère terms, and*
3) *An exact energy conservation law for the Gyrokinetics Vlasov-Maxwell equations that includes all the relevant linear and nonlinear coupling terms. The foundations of nonlinear Gyrokinetics theory are reviewed with an emphasis on rigorous application of Lagrangian and Hamiltonian Lie-transform perturbation methods in the variational derivation of nonlinear Gyrokinetics Vlasov-Maxwell equations. The physical motivations and applications of the nonlinear Gyrokinetics equations that describe the turbulent evolution of low-frequency electromagnetic fluctuations in a non-uniform magnetized plasmas with arbitrary magnetic geometry are discussed.*[3]

The nonlinear Gyrokinetics system of equations for the arbitrary fluctuation frequencies are also derived from the generalized Lagrangian formulation and the variational principle. They are recommended for simulations and analytic applications in laboratory.

The Lagrangian formulation of the Gyrokinetics theory is generalized in order to describe the particles' dynamics as well as the self-consistent behavior of the electromagnetic fields. The Gyrokinetics equation for the particle distribution function and the Gyrokinetics Maxwell's equations for the electromagnetic fields are both derived from the variational principle for the Lagrangian consisting of the parts of particles, fields, and their interaction. In this generalized Lagrangian formulation, the energy conservation property for the total nonlinear Gyrokinetics system of equations is directly shown from the Noether's theorem. This formulation can be utilized in order to derive the nonlinear Gyrokinetics system of equations and the rigorously conserved total energy for fluctuations with arbitrary frequencies.[4]

From these formulations it is possible to perform computer simulations: *Simulations conducted for numerous situations, and their comparison, It has been confirmed that the existing Gyrokinetics models are sufficiently valid for shorter times than the real time scales of the evolution of the reactor's dynamic profiles, thus it is thought that the effect of turbulence on the electric field and plasma velocity provide incorrect values for the electric field. Consequently more precise mathematical models are needed to represent the real behaviour of plasma at a longer time scale, called the transport scale.*[5]

And also simulations for specific reactor designs: *Simulations of toroidal angular momentum transport have been carried out using global toroidal Gyrokinetics particlein-cell code. The significant redistribution of toroidal momentum is observed, driven by the ion temperature gradient turbulence with adiabatic electrons, resulting in a peaked momentum profile in the central region of the radial domain. Cases with rigid and sheared plasma rotation are considered. Diffusive and off-diagonal pinch like fluxes are identified. Toroidal momentum diffusivity is calculated by subtracting pinch contribution from the total momentum flux, and compared to quasilinear estimates. It is found that the ratio of momentum to heat conductivity is smaller than unity even after subtracting pinch contribution when wave-particle resonance energy is larger than thermal energy.*[6]

III. Transport Physics

New concepts of physical transport toroidal plasmas are studied. The transport of plasma in the reactor is generally externally provided by the momentum input from neutral beam injection in the current generation of tokamaks, which causes its rotation.

The transport characteristics of non-diffusion and non-locality of transport has been recognized to be important in determining radial structures of density, rotation, and temperature, as well as non-linearity in the flux-gradient relation. The non-diffusive term of momentum transport appears as a 'spontaneous rotation and intrinsic torque', while the non-diffusive term of particle transport appears as a 'particle pinch and particle exhaust'. The sign and magnitude of these non-diffusive terms have been found to be sensitive to the turbulence state, which causes reversal phenomena. In the momentum transport, the spontaneous flow reversal from the co- (parallel to plasma current) direction to the counter- (anti-parallel to plasma current) direction and

[3] Brizard, Alain J. and Hahm, T. S.: Foundations of Nonlinear Gyrokinetic Theory. Reviews of Modern Physics, Volume 79, April–June 2007. DOI: 10.1103/RevModPhys.79.421

[4] Sugama, H.: *Gyrokinetic Field Theory*; National Institute for Fusion Science (NIFS) Aug. 1999. NIFS-609 Nagoya, Japan.

[5] Barceló, G.: *Dynamic Interaction Confinement.* World Journal of Nuclear Science and Technology Vol.4 No.4, October 29, 2014DOI: 10.4236/wjnst.2014.44031.

[6] I. Holoda and Z. Lin: Gyrokinetic particle simulations of toroidal momentum transport Department of Physics and Astronomy, University of California, Irvine, California 92697, USA, © 2008, American Institute of Physics. DOI: 10.1063/1.2977769.

© 2016 Global Journals Inc. (US)

vice versa have been commonly observed even when the plasma parameter changes slightly. In the particle transport, especially in the impurity transport, reversals of convective radial flux from inward to outward are also observed in toroidal plasmas as phenomena of density peaking/flattening and impurity accumulation/exhaust. (…)

The non-diffusive term of momentum transport appears as a 'spontaneous rotation and intrinsic torque', while the non-diffusive term of particle transport appears as a 'particle pinch and particle exhaust'. [7]

The Toroidal angular momentum (TAM) is a conserved quantity.[8] *Plasma rotation plays important role in turbulence stabilization and in suppression of resistive wall modes, therefore affecting tokamak performance. In present day machines, rotation is usually driven by external sources, such as neutral beam injection; however, it might be unavailable in a future devices such as ITER, due to the large machine size and high densities. Possible solution of this problem could be generation of spontaneous intrinsic rotation and on-axis peaking of momentum profile by inward pinch like flux. Thus, it becomes crucial to understand the nature of intrinsic rotation phenomena, together with momentum transport mechanisms.*[9]

Poloidal and toroidal rotation has been recognized to play an important role in heat transport and magneto hydrodynamic (MHD) stability in tokamaks and helical systems. It is well known that the E × B shear due to poloidal and toroidal flow suppresses turbulence in the plasma and contributes to the improvement of heat and particle transport, while toroidal rotation helps one to stabilize MHD instabilities such as resistive wall modes and neoclassical tearing mode. Therefore, understanding the role of momentum transport in determining plasma rotation is crucial in toroidal discharges, both in tokamaks and helical systems.[10]

IV. Heuristic Description of *Intrinsic Toroidal Rotation*

Plasma rotation is an essential factor in this analysis of the turbulent transport of momentum. In magnetic confinement fusion systems such as on the ITER[11], the plasma circulates in the container at a constant movement, which we could define as rotation with respect to its walls. An intrinsic spontaneous toroidal rotation, with no external momentum input, have been observed on tokamaks: *it has been shown*[12] *that the plasma in the reactor can initiate* ***spontaneous circular movement or rotation****, without the need for any external dynamic momentum input. The theoretical development of this behaviour is still under study.*[13]

According to Parra: *Experimental observations have shown that tokamak plasmas rotate spontaneously without momentum input. This intrinsic rotation has been the object of recent work because of its relevance for ITER, where the projected momentum input from neutral beams is small, and the rotation is expected to be mostly intrinsic. The origin of the intrinsic rotation is still unclear.* (…)

The intrinsic rotation profile depends on the density and temperature profiles and on the up-down asymmetry.[14]

Changes in turbulence accompany the changes in spontaneous intrinsic rotation and they are studied in several successive radial regions, using a variety of diagnostics.

There has been some theoretical work in turbulent transport of momentum using Gyrokinetics simulations[15] [16] [17] [18] [19] *and different mechanisms have been proposed as candidates to explain intrinsic rotation*[20].

[7] K Ida: *New concepts of transport physics in toroidal plasmas.* Published 20 March 2015, IOP Publishing Ltd. Plasma Physics and Controlled Fusion, Volume 57, N°4.

[8] G.M. Staebler et al, BAPS 46 (2001) p221-LP1 17.

[9] I. Holoda and Z. Lin: *Gyrokinetic particle simulations of toroidal momentum transport* Department of Physics and Astronomy, University of California, Irvine, California 92697, USA, © 2008, American Institute of Physics. DOI: 10.1063/1.2977769.

[10] K. Ida and J.E. Rice: *Rotation and momentum transport in tokamaks and helical systems.* Published 10 March 2014, 2014 IAEA, Vienna. Nuclear Fusion, Volume 54, Number 4. doi:10.1088/0029-5515/54/4/045001

[11] Ikeda, K. *Progress in the ITER physics basis*. Nuclear Fusion, 47(6) (2007). http://stacks.iop.org/0029-5515/47/i=6/a=E01.

[12] Rice, J. E. et al. (2007) *Nuclear Fusion,* 47 1618.

[13] Parra, F. I., Barnes, M., Calvo, I. and Catto, P. J. *Intrinsic rotation with gyrokinetic models.* Physics of Plasmas, 19(5), 056116 (2012). DOI: 10.1063/ 1.3699186. http://link.aip.org/link/?PHP/19/056116/1.

[14] Parra, F. I., Barnes, M., Peierls, R. and Catto, P. J. *Sources of intrinsic rotation in the low flow ordering.* Centre for Theoretical Physics, University of Oxford, Oxford, UK. 2012, Plasma Science and Fusion Center, Massachusetts Institute of Technology.

[15] Peeters, A. G., Angioni, C. and Strinzi, D. (2007) *Physical Review Letters,* 98 265003.

[16] Waltz, R. E., Staebler, G. M., Candy, J. and Hinton, F. L. (2007) *Physics of Plasmas,* 14 122507 and Waltz, R. E., Staebler, G. M., Candy, J. and Hinton, F. L. (2009) 16 079902. Roach, C. M. et al. (2009) *Plasma Physics and Controlled Fusion,* 51 124020.

[17] Camenen, Y., Peeters, A. G., Angioni, C., Casson, F. J., Hornsby, W. A., Snodin, A. P. and Strintzi, D. (2009) *Physical Review Letters,* 102 125001.

[18] Casson, F. J., Peeters, A. G., Camenen, Y., Angioni, C., Hornsby, W. A., Snodin, A. P., Strintzi, D. and Szepesi, G. (2009) *Physics of Plasmas,* 16 092303 and Casson, F. J., Peeters, A. G., Angioni, C., Camenen, Y., Hornsby, W. A., Snodin, A. P. and Szepesi, G. (2010) *Physics of Plasmas,* 17 102305.

[19] Highcock, E. G., Barnes, M., Schekochihin, A. A., Parra, F. I., Roach, C. M. and Cowley, S. C. (2010) *Physical Review Letters,* 105 215003 and Barnes, M., Parra, F. I., Highcock, E. G., Schekochihin, A. A., Cowley, S. C. and Roach, C. M. (2011) submitted to *Physical Review Letters*.

[20] Wang, L. and Diamond, P. H. *Gyrokinetic Theory of Turbulent Acceleration of Parallel Rotation in Tokamak Plasmas.* Physical Review Letters, 110, 265006 – Published on 27 June 2013. CEEE, Huazhong University of Science and Technology, Wuhan, Hubei 430074, China. WCI Center for Fusion Theory, NFRI, Gwahangno 113, Yusung-gu, Daejeon 305-333, Korea and CMTFO and CASS, University of California, San Diego, La Jolla, California 92093-0424, USA.

© 2016 Global Journals Inc. (US)

The presence of an external magnetic field introduces a source of plasma anisotropy. Variations or fluctuations in the thermodynamic fields are much greater perpendicularly to the field than parallel to it. (…)

The ion toroidal rotation in a tokamak consists of an E×B flow due to the radial electric field and a diamagnetic flow due to the radial pressure gradient… the momentum pinch for the rotation generated by the radial pressure gradient is calculated and is compared with the Coriolis pinch. This distinction is important for subsonic flows or the flow in the pedestal where the two types of flows are similar in size and opposite in direction. At the edge, the different pinches due to the opposite rotations[21] can result in intrinsic momentum transport that gives significant rotation peaking.[22]

We assume that heating induced spontaneous toroidal rotation and other consequences of anomalous momentum transport, it has been observed toroidal rotation during heating, without external torque.

Parametric scaling of the intrinsic (spontaneous, with no external momentum input) toroidal rotation observed on a large number of tokamaks have been combined with an eye towards revealing the underlying mechanism(s) and extrapolation to future devices. The intrinsic rotation velocity has been found to increase with plasma stored energy or pressure in JET, Alcator C-Mod, Tore Supra, DIII-D, JT-60U and TCV, and to decrease with increasing plasma current in some of these cases. Use of dimensionless parameters has led to a roughly unified scaling with MA ∝ βN, although a variety of Mach numbers works fairly well; scaling of the intrinsic rotation velocity with normalized gyro-radius or collisionality show no correlation. Whether this suggests the predominant role of MHD phenomena such as ballooning transport over turbulent processes in driving the rotation remains an open question. For an ITER discharge with βN = 2.6, an intrinsic rotation Alfven Mach number of MA sime 0.02 may be expected from the above deduced scaling, possibly high enough to stabilize resistive wall modes without external momentum input.[23]

V. Phenomenology

Intrinsic spontaneous toroidal rotation driven by pressure and ion temperature gradients has been long predicted, studied and analyzed.

Spontaneous toroidal rotation, self-generated in the absence of an external momentum input, exhibits a rich phenomenology. In L-mode plasmas, the rotation varies in a complicated fashion with electron density, magnetic configuration and plasma current and is predominantly in the counter-current direction. The rotation depends sensitively on the balance between the upper and lower null and plays a crucial role in the H-mode power threshold. Rotation inversion between the counter- and co-current directions has been observed following small changes in the electron density and plasma current, with very distinct thresholds. In contrast, the intrinsic rotation in H-mode plasmas has a relatively simple parameter dependence, with the rotation velocity, proportional to the plasma stored energy normalized to the plasma current, and is nearly always directed co-current. In plasmas with internal transport barriers, formed either with off-axis ICRF heating or LHCD, the core rotation velocity increments in the counter-current direction as the barrier evolves.[24]

The experimental results of investigations of the mechanisms behind the momentum transport, and predictions of intrinsic rotation allow evolving in these investigations.

Regarding momentum transport, the radial flux of momentum has diffusive and nondiffusive (ND)terms, and experimental investigations of these are discussed. The magnitude of the diffusive term of momentum transport is expressed as a coefficient of viscous diffusivity. The ratio of the viscous diffusivity to the thermal diffusivity (Prandtl number) is one of the interesting parameters in plasma physics. It is typically close to unity, but sometimes can deviate significantly depending on the turbulent state. The ND terms have two categories: one is the so-called momentum pinch, whose magnitude is proportional to (or at least depends on) the velocity itself, and the other is an offdiagonal term in which the magnitude is proportional to (or at least depends on) the temperature or/and pressure gradient, independent of the velocity or its gradient. The former has no sign dependence; rotation due to the momentum pinch does not depend on the sign of the rotation itself, whether it is parallel to the plasma current (codirection) or anti-parallel to the plasma current (counter-direction). In contrast, the latter has a sign dependence; the rotation due to the off-diagonal residual term is either in the co- or counter-direction depending on the turbulence state, but not on the sign of the rotation itself. This residual term

[21] Lee, J. P., Parra, F. I. and Barnes, M. *Turbulent momentum pinch of diamagnetic flows in a tokamak.* arXiv.org physics arXiv:1301.4260. Plasma Science and Fusion Center, MIT, Cambridge, USA, 21 January 2013.

[22] Barceló, G.: *Dynamic Interaction Confinement.* World Journal of Nuclear Science and Technology Vol.4 No.4, October 29, 2014DOI: 10.4236/wjnst.2014.44031.

[23] J.E. Rice, A. Incecushman, J.S. de Grassie, L.-G. Eriksson, Y. Sakamoto, A. Scarabosio, A. Bortolon, K.H. Burrell, B.P. Duval, C. Fenzi-Bonizec: *Inter-machine comparison of intrinsic toroidal rotation in tokamaks.* Published 31 October 2007, IAEA, Vienna • Nuclear Fusion, Volume 47, Number 11

[24] J E Rice, A C Ince-Cushman, M L Reinke, Y Podpaly, M J Greenwald, B LaBombard and E S Marmar: Spontaneous core toroidal rotation in Alcator C-Mod L-mode, H-mode and ITB plasmas. Published 5 November 2008 • 2008 IOP Publishing Ltd • Plasma Physics and Controlled Fusion, Volume 50, Number 12

[25] K. Ida and J.E. Rice: Rotation and momentum transport in tokamaks and helical systems. Published 10 March 2014, 2014 IAEA, Vienna. Nuclear Fusion, Volume 54, Number 4. doi:10.1088/0029-5515/54/4/045001

© 2016 Global Journals Inc. (US)

can also act as a momentum source for intrinsic rotation. [25]

VI. Predictions for Intrinsic Rotation

It is important to make predictions for the intrinsic rotation for future reactors: *There are many unresolved problems regarding intrinsic rotation. One of the important issues is the sign of the rotation. As seen in the reversal phenomenon, the direction of the intrinsic rotation changes its sign, even with slight changes in plasma parameters, which cannot be expressed by a global parameter scaling. The other issue is that the intrinsic rotation is very sensitive to the change in turbulence type, and also to the confinement mode, such as LOC/SOC, L-mode/H-mode or L-mode/ITB. Although the magnitude of intrinsic rotation may be sufficient in ITER, the flip in rotation is of concern, so an understanding of rotation reversals is crucial. In order to achieve stable operation in ITER, a better understanding based on the physics mechanisms of intrinsic rotation is necessary.*

There are several physics issues which should be understood in order to predict the intrinsic rotation in future devices. The physics mechanism determining the direction and magnitude of intrinsic torque is still not well enough understood to give a precise prediction, although there are many experimental observations and simulations. Since the intrinsic torque is related to the symmetry breaking of turbulence, it should be correlated with the confinement regime, as well as to heat and particle transport. At the moment, the understanding of the connection among the particle pinch, intrinsic torque and non-local response of the heat transport is still phenomenological. The other remaining issue is how the angular momentum is released when the intrinsic rotation appears. It is not clear whether angular momentum is conserved by the kinetic momentum escaping from the last closed flux surface or by a change in the magnetic vector potential. [26]

There are several others outstanding issues, such as how the angular momentum is released when the intrinsic rotation appears.

Regression analysis has been performed on the intrinsic rotation data set:

Extrapolation to ITER predicts rotation velocities in some of the scatter may be due to the large variety in magnetic field ripple, error fields and/or wall conditioning in the various devices. Extrapolation to ITER predicts rotation velocities in excess of 300km/s. (…) *There are several physics issues which should be understood in order to predict the intrinsic rotation in future devices. The physics mechanism determining the direction and magnitude of intrinsic torque is still not well enough understood to give a precise prediction, although there are many experimental observations and simulations. Since the intrinsic torque is related to the symmetry breaking of regime, as well as to heat and particle transport.* [27]

Recently there has been widespread attention paid to rotation and momentum transport in tokamak plasmas. Of particular interest is spontaneous (intrinsic) toroidal rotation in plasmas without external momentum input. The strong co-current spontaneous rotation in enhanced confinement regimes, with ion thermal Mach numbers up to 0.3, may allow for resistive wall mode suppression in high-pressure ITER discharges, without requiring the use of neutral beam injection. Spontaneous rotation in L-mode discharges exhibits a complex dependence on plasma parameters and magnetic configuration compared to the relatively simple scaling of Alfven Mach number ($MA = V\Phi/CA$, where CA is the Alfven speed) $MA \sim \beta N$ observed in enhanced confinement plasmas. There is currently no comprehensive, quantitative explanation of this phenomenon. An accurate prediction of the expected rotation velocity profile from whatever neutral beam injection is available on ITER requires a detailed understanding of momentum transport. There have been extensive investigations into correlations between energy and momentum diffusivities, and whether there are systematic trends of the Prandtl number with plasma parameters. Of late, there has been vigorous theoretical activity regarding a possible momentum pinch that could help enhance the rotation in the plasma interior. There has been a renewed interest in poloidal rotation, especially in ITB discharges, which is generally found to be at odds with the predictions of neo-classical theory. (…)

Spontaneous rotation (in the absence of external momentum input) in low confinement mode (L- mode) Ohmic plasmas is often directed counter-current, and depends very sensitively on the magnetic configuration and in a complicated fashion on other parameters, such as the plasma current, electron density and ion temperature. While the study of spontaneous rotation in L-mode discharges is of interest in its own right (e.g. for its relation to the H-mode power threshold), these plasmas will not be considered here, since most ignition scenarii in future devices require H- mode confinement. The spontaneous toroidal rotation in H-mode (and in enhanced confinement regimes in Tore Supra) is generally in the co-current direction and has been observed on many devices and produced with a wide

[26] K. Ida and J.E. Rice: Rotation and momentum transport in tokamaks and helical systems. Published 10 March 2014, 2014 IAEA, Vienna. Nuclear Fusion, Volume 54, Number 4. doi:10.1088/0029-5515/54/4/045001.

[27] *K. Ida and J.E. Rice: Rotation and momentum transport in tokamaks and helical systems. Published 10 March 2014, 2014 IAEA, Vienna. Nuclear Fusion, Volume 54, Number 4. doi:10.1088/0029-5515/54/4/045001.*

© 2016 Global Journals Inc. (US)

variety of techniques (ICRF, Ohmic, ECH), demonstrating its fundamental nature. (…)

Spontaneous (in the absence of external momentum input) co-current toroidal rotation has been observed on many tokamaks in H-mode (and in enhanced confinement regimes in Tore Supra), produced by a variety of different methods: ICRH, ECH, Ohmic heating, ECH with lower hybrid heating and balanced neutral beam injection. A common feature of these observations is that the rotation velocity increases with the plasma pressure or stored energy. (…) *At present there is no comprehensive, quantitative explanation for spontaneous rotation. Some recent observations of poloidal rotation are inconsistent with the predictions of neo-classical theory. Momentum transport has been found to be closely coupled to energy transport, and with momentum diffusivities much higher than neo-classical values. These results are consistent with diffusive momentum transport driven by ITG turbulence. Evidence for an inward momentum pinch has also been observed, which could be due to the Coriolis drift effect or from symmetry breaking due to magnetic field curvature.*[28]

The theory of turbulent transport of toroidal momentum is discussed in the context of the phenomenon of spontaneous/intrinsic rotation. We review the basic phenomenology and survey the fundamental theoretical concepts. We then proceed to an in-depth discussion of the radial flux of toroidal momentum, with special emphasis on the off-diagonal elements, namely the residual stress (the portion independent of V) and the pinch. A simple model is developed which unifies these effects in a single framework and which recovers many of the features of the Rice scaling trends for intrinsic rotation. We also discuss extensions to finite beta and the effect of SOL boundary conditions. Several issues for future consideration are identified.[29]

Spontaneous spin-up in intrinsic toroidal and poloidal rotation of impurity ions (C and He) was observed at the edge of ohmic plasmas during L-H mode transitions in NSTX. The rotation was reduced again after the H-L back transition. The changes in rotation were accompanied by changes in plasma fluctuations and turbulence observed over the plasma minor radius from the far SOL too deep in the core using diagnostics such as fast reflectometry and Langmuir probes near the chamber walls, gas puff imaging (GPI), and microwave reflectometry and high-k scattering in the plasma core. Intrinsic rotation increases of 10s of km/s were observed using an edge rotation diagnostic and reflectometry showed the long wavelength correlation length in the plasma core to drop sharply at the L-H transition. GPI and a fast radial (midplane) scanning probe showed a strong decrease in plasma turbulence (…) at the plasma edge. The observed intrinsic velocity scalings are consistent with a variety of toroidal devices.[30]

VII. Dynamic Model for Confined Plasma

Against this background of study and constant advancement in unravelling the physical behaviour of tokamak-type nuclear reactors, we suggest a revision of the dynamic criteria being applied, given that we believe there may well be inappropriate interpretations in the principles of classical rotational dynamics at play.[31]

We propose a dynamic new model for confined plasma based on the theory of dynamics interactions (TDI), to allow, a natural dynamic confinement. We advise the study of rotational dynamics, given that it would afford us a better understanding of the dynamics of confined plasma by dynamic interaction, along with an improved reactor result.

In accordance with the theory of dynamics interactions, we suggest that each particle, instead of following a linear path, as would be expected from the classical mechanics equations of Newton-Euler, it would follow a closed path owing to the coupling of the velocities' fields that are caused on every point of each particle.

The issue can be summed up as follows: If we apply this dynamic theory to a tokamak reactor, every plasma particle with intrinsic rotation will follow a closed path like that of the boomerang when it is subjected to a non-coaxial momentum in addition to its intrinsic angular momentum. Consequently, there are different authors[32]

[28] *J E Rice: Spontaneous rotation and momentum transport in tokamak plasmas. Plasma Science and Fusion Center Massachusetts Institute of Technology Cambridge MA 02139 USA, 2007, U.S. Department of Energy, Grant DE DE-FC02- 99ER54512, and IOP Publishing Ltd. Journal of Physics: Conference Series, Volume 123, Number 1, 2008.*

[29] *P. H. Diamond and C. Mcdevitt and O. D. Gurcan and K. Miki and T. S. Hahm and W. Wang and G. Rewoldt and I. Holod and Z. Lin and V. Naulin and R. Singh: Physics of Non-Diffusive Turbulent Transport of Momentum and the Origins of Spontaneous Rotation in Tokamaks. The College of Information Sciences and Technology, 2015. The Pennsylvania State University.*

[30] Bush, C. E.; Bell, R. E.; Kubota, S.; Ahn, J.-W.; Maqueda, R. J.; Zweben, S. J.; Leblanc, B. P.; Lee, K. C.; Mazzucato, E.; Wilgen, J. B.; Fredrickson, E. D.; Raman, R.; Delgado-Aparicio, L.; Stutman, D.; Tritz, K.; Kaye, S. M.: Spontaneous Intrinsic Rotation and Changes in Turbulence in Ohmic H-modes on NSTX. American Physical Society, 50th Annual Meeting of the Division of Plasma Physics, November 17-21, 2008, abstract #NP6.091.

[31] Barceló, G.: Dynamic Interaction Confinement. World Journal of Nuclear Science and Technology Vol.4 No.4, October 29, 2014DOI: 10.4236/wjnst.2014.44031 http://www.scirp.org/journal/PaperInformation.aspx?paperID=51026& http://dx.doi.org/10.4236/wjnst.2014.44031

[32] Pérez, L. New Evidence on Rotational Dynamics. World Journal of Mechanics, Vol. 3, No. 3,2013, pp. 174-177, doi: 0.4236/wjm.2013.33016. http://www.scirp.org/ journal/wjm http://dx.doi.org/10.4236/wjm.2013.33016

© 2016 Global Journals Inc. (US)




Illustration with the image of the trajectory of the plasma according to the criteria of the TDI. (by Al. Pérez)

who share these same hypotheses[33] *with a view to understanding the dynamic behaviour of bodies or particles in those circumstances in which they have translational velocity, intrinsic rotation and are simultaneously subjected to a non-coaxial momentum with its own rotation.*[34] *Notwithstanding, in our hypothesis, based on a possible analogy, we are extrapolating the observed behaviour in macroscopic moving bodies to the reactor's plasma particles.*

In our opinion, this dynamic model also coincides with the behaviour of hurricanes on the Earth and also resembles atmospheric toroidal vortex rings.

It must be pointed out that in our investigations we have come to a rational deduction to the effect that in such circumstances, the kinetic translational energy can be transformed into kinetic rotational energy, and vice versa, and in general, energy transfers can occur at the core of particles endowed with intrinsic angular momentum: […] kinetic energy can be transferred, increasing its rotation velocity, its linear velocity or modifying its state of potential.[35]

We suggest that the spontaneous/intrinsic rotation can be explained by this dynamic theory: the plasma particles with intrinsic rotation, undergo to a not coaxial torque, initiated the orbiting movement, in the same way that spontaneously initiates a top. By the effect of the external torque, the travel speed of the particle changes its direction but not its magnitude, initiating a closed-orbiting movement, as expressed by the TID. The goal, in this case for better efficiency, it will match the radius of this spontaneous movement with the radius of the reactor. Thus we conceive the plasma as a whole on a closed orbit, inside the reactor, which defines the proposed dynamic inertial confinement, and justified by the TDI.

Thus, the plasma has a repetitive motion equivalent to the boomerang, returning to his place of origin. Both dynamic behaviors, the top and the boomerang are, in our opinion, clear examples of the rotational dynamics justified by the Theory of Dynamic Interactions, TDI, that we propose.

VIII. Lorentz Force

At the beginning of this paper we stated that magnetic confinement is based on the Lorentz force. This force manifests itself when an electric charge in movement moves in a magnetic field. We have stated in other works that, in our opinion, the origin of this force is analogous to that of the inertial forces that manifest themselves in TDI circumstances, and which oblige particles to follow a closed path. Therefore, and in accordance with our hypothesis, the same dynamic effect and the same effective confinement could be obtained doubly, by the magnetic effect and by the dynamic effect.

The Lorentz force is an electromagnetic interaction manifestation, discovered by Hendrik Lorentz, with multiple technological applications. It is the total combination of electric and magnetic force on a point charge due to electromagnetic fields.

[33] Dorado González, M. Dinámica de sistemas con spin: un nuevo enfoque. Fundamentos y aplicaciones. ADI Ser, Ed.: Madrid, 2013.
[34] Dorado González, M. Equation of motion of systems with internal angular momentum – II. arXiv: physics/ 0603207.
[35] Barceló, G. A rotating world (Un mundo en rotación). Editorial Marcombo: Barcelona, 2008. http://www. dinamicafundacion.com/

© 2016 Global Journals Inc. (US)

For a particle of charge subjected to an electric field combined with a magnetic field, the total electromagnetic force or Lorentz force on the particle is given by:

$$\mathbf{f} = q(\mathbf{E} + \mathbf{V} \times \mathbf{B}),$$

Where:
q= particle charge
v= velocity in the presence of an electric field
E= electric field intensity
B= external magnetic field

If we look carefully at the effects a B field exerts on an electrical charge q, it holds that:

- When the particle is at rest, the field B exerts no force on it.
- The force is maximum when the load speed v and the field B are orthogonal and is zero when they are parallel.
- The force is perpendicular to the plane formed by v and B.
- The force is proportional to the charge q, and to de speed v.
- If the load changes sign, the force changes direction

Therefore, the force exerted by a field B on electric charge q moving with velocity v is perpendicular to the field lines B, causes a normal acceleration and change the path charge. Accordingly, a circular movement starts, in which the magnetic force acts as normal force or centripetal, and wherein the modulus of the speed remains constant, there being no tangential force.

This is the same behavior justified by the TDI, to the bodies with intrinsic rotation, when subjected to non-coaxial torque. In these cases, a new curved path also circular is generated, when maintained constants the parameters.

If we simply refer to the magnetic moment generated by a magnet, this determines the force that the magnet can exert on electric currents and magnetic field torque exerted on them. Its formulation is:

$$\boldsymbol{\tau} = \boldsymbol{\mu} \times \mathbf{B}$$

Where:
$\boldsymbol{\tau}$ = is the torque acting on the dipole
$\boldsymbol{\mu}$ = is the magnetic moment.

$\boldsymbol{\tau}$ It is always normal to both, both $\boldsymbol{\mu}$ as B. When B is orthogonal to $\boldsymbol{\mu}$, then it is maximum. Note that this dynamic relationship between two quantities, is analogous to the dynamic equation of the TDI, also the dynamic phenomena occurring in both cases are also similar.

It is assumed, in the case of magnetic materials, that the cause of the magnetic torque are the spin and orbital angular momentum states of the electrons, and whether atoms in one region are aligned with atoms in another.

Magnetic torque and angular momentum have a close connection. It is expressed on a macroscopic scale in the Einstein-de Haas effect, or "rotation by magnetization," and otherwise with its inverse, the Barnett effect, or "magnetization by rotation." Both can be defined as gyromagnetic effects. This applies, for example, when a magnetic moment is subject to a torque in a magnetic field that tends to align it with the applied magnetic field, the moment precesses (rotates about the axis of the applied field).

In these phenomena we can imagine a magnetic dipole as a rotating charged particle. In this case both, the magnetic moment and the angular momentum, increase with the rate of rotation.

Electrons and many atomic nuclei, have intrinsic magnetic moments, which relates to the angular momentum of the particles. This way you can justify the behavior of the plasma. The intrinsic angular momentum (spin) of each type of particle of the plasma is a constant.

The spin magnetic moment is an intrinsic or essential property of particles, such as mass or electrical charge. These intrinsic magnetic moments are what give rise to macroscopic effects of magnetism, and other phenomena such as nuclear magnetic resonance.

In addition, certain orbital distributions, implies an additional magnetic moment, by the movement of electrons as charged particles.

Both ideas being proposed should, we feel, be further explored and, where appropriate, confirmed experimentally; on the one hand, that the Lorentz force is analogous to an inertial dynamic force, similar to TDI ones, and on the other hand, that this theory be applied to the reactor's plasma.[36]

IX. Conclusions

Nuclear fusion is a necessary energy into which countries are researching and in which we trust. Although, to some authors: *At present there is no comprehensive, quantitative explanation for spontaneous rotation,*[37] but in this text we have proposed a dynamic justification of such expontanea plasma rotation.

[36] Barceló, G.: Dynamic Interaction Confinement. World Journal of Nuclear Science and Technology Vol.4 No.4, October 29, 2014DOI: 10.4236/wjnst.2014.44031
http://www.scirp.org/journal/PaperInformation.aspx?paperID=51026&
http://dx.doi.org/10.4236/wjnst.2014.44031

[37] J E Rice: Spontaneous rotation and momentum transport in tokamak plasmas. Plasma Science and Fusion Center Massachusetts Institute of Technology Cambridge MA 02139 USA, 2007, U.S. Department of Energy, Grant DE DE-FC02- 99ER54512, and IOP Publishing Ltd. Journal of Physics: Conference Series, Volume 123, Number 1, 2008.

© 2016 Global Journals Inc. (US)

The focus however is on new theory and simulation of the ExB shear and coriolis angular momentum pinch effects needed to understand the intrinsic toroidal rotation in tokamaks without external torque in the core.[38]

We also understand that should be explored a new theory and simulation, but that's not exactly a Coriolis effect, or the ExB shear where the solution must be found, if not in the TID we propose.

As far as magnetic confinement reactors are concerned, plasma dynamics is still a work in progress. We are as yet ignorant of the exact dynamic behaviour of plasma and, moreover, when designing these generators it is still necessary to reduce their turbulent tendency and improve their momentum transport capacity.

We propose a profound revision of the principles of classical rotational mechanics, especially as regards particles subjected to accelerations by simultaneous and noncoaxial rotations. After conducting our research, we put forward certain specific, alternative dynamic hypotheses.

In addition to the plasma's material container, we can envisage another simultaneous, non-material container, along with the use of magnetic confinement techniques. We suggest the exploring of a new type of dynamic confinement based on the TDI and one that is compatible with magnetic confinement.

Gyrokinetics is the workhorse for modern research on low-frequency micro turbulence in magnetized plasmas. It is a very powerful tool that has been used successfully in many different applications in both fusion and astrophysics.[39]

We recognize that the issue is still unresolved. We understand that it is not correct to speak of the Coriolis effect. The problem lies in the design of the equations of motion applied. We propose the implementation of a new rotational dynamics based on the theory suggested in this text.

Moreover, in our opinion, the TDI could also be employed as a complementary theoretical instrument that would lead to greater reactor efficiency. Consequently, its dynamic hypotheses ought to be studied in the field of Gyrokinetics theory.

The equation for movement that we are proposing in the Theory of Dynamic Interactions for these non-inertial situations is very simple […] it becomes possible to determine the path of plasma particles in translation, when these particles are endowed with intrinsic rotation and are subjected to rotation actions on another, different axis. […]

These new dynamic hypotheses we are putting forward and which we hold applicable to particle systems accelerated by rotation can, we suggest, be used in the interpretation and design of fusion reactors. Accordingly, we persist in our belief that magnetic confinement in a reactor can be likened to confinement by dynamic interactions based on the Theory of Dynamic Interactions. Applying this criterion we are proposing would enable a twin physical-theoretical principle to isolate plasma and try to minimize its turbulence: each plasma particle should have spin and be subjected to a non-coaxial magnetic momentum from without and another gravimetric one. Accordingly, plasma confinement will be based on two different, albeit analogous, physical principles, generating double intransigence in its path. [40]

We understand that the variational analysis used in the applied formulation is currently insufficient, if not accompanied by the dynamic hypothesis that we propose. It is necessary to incorporate on the mathematical formulation, the coupling of the fields of the dynamics magnitudes that are generated in the plasma, because this dynamic coupling defines its true behavior: the plasma circulating along the reactor, in a closed path, generating an additional dynamic confinement, besides already magnetic known. The dynamic model based on a new rotational concept, may represent more correctly the real behavior of the plasma in the rector, even could improve the performance of these energy power plants, minimize turbulence and improve momentum transport.

[38] R. E. Waltz, G.M. Staebler and J. Candy: Gyrokinetic theory and simulation of momentum transport and energy exchange. General Atomics PPPL Theory/NSTX. Seminar October 22, 2007

[39] Barceló, G.: Dynamic Interaction Confinement. World Journal of Nuclear Science and Technology Vol.4 No.4, October 29, 2014DOI: 10.4236/wjnst.2014.44031
http://www.scirp.org/journal/PaperInformation.aspx?paperID=51026&
http://dx.doi.org/10.4236/wjnst.2014.44031

[40] Barceló, G.: *Dynamic Interaction Confinement.* World Journal of Nuclear Science and Technology Vol.4 No.4, October 29, 2014, DOI: 10.4236/wjnst.2014.44031
http://www.scirp.org/journal/PaperInformation.aspx?paperID=51026&
http://dx.doi.org/10.4236/wjnst.2014.44031

© 2016 Global Journals Inc. (US)

Global Journal of Science Frontier Research (A) Volume XVI Issue III Version I Year 2016
10

This page is intentionally left blank

© 2016 Global Journals Inc. (US)

Global Journal of Science Frontier Research: a
Physics and Space Science
Volume 16 Issue 3 Version 1.0 Year 2016
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Bi-Directional EPR Correlation in Cosmology and Planckeon Origin of Dark Energy

By Noboru Hokkyo
Senjikan Institute, Japan

Abstract- A solution of nonlocal EPR correlation between counter-propagating pair of polarization-entangled photons emitted from a common source at S and detected at points P and Q is sought outside the EPR's reality criterion of local causality but within the framework of time-symmetric quantum electrodynamics allowing the bi-directional signal transmission P↔S↔Q on the double-light cone where the future and the past cones share common light paths connecting the photon source S and the detection points P and Q. A cosmlogical implication of the bi-directional signal transmission P↔Q without common source S in the inflationary cosmology and possible Planckeon orgin of dark energy in the upper hemisphere of semiclosed Friedman uiverse, joined on to an asymptotically flat outer space, are also discussed.

Keywords: quantum theory, relativity, causality, locality, correlation cosmology, dark energy.

GJSFR-A Classification : FOR Code: 020109



Strictly as per the compliance and regulations of :

© 2016. Noboru Hokkyo. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Bi-Directional EPR Correlation in Cosmology and Planckeon Origin of Dark Energy

Noboru Hokkyo

Abstract- A solution of nonlocal EPR correlation between counter-propagating pair of polarization-entangled photons emitted from a common source at S and detected at points P and Q is sought outside the EPR's reality criterion of local causality but within the framework of time-symmetric quantum electrodynamics allowing the bi-directional signal transmission P↔S↔Q on the double-light cone where the future and the past cones share common light paths connecting the photon source S and the detection points P and Q. A cosmlogical implication of the bi-directional signal transmission P↔Q without common source S in the inflationary cosmology and possible Planckeon orgin of dark energy in the upper hemisphere of semiclosed Friedman uiverse, joined on to an asymptotically flat outer space, are also discussed.

Keywords: quantum theory, relativity, causality, locality, correlation cosmology, dark energy.

I. Introduction

Since the advent of quantum mechanics in the mid-1920s there have benn continued interpretational controversies surrounding its counter-intuitive nature such as the wave-particle duality and the instantaneous collapse of the particle wave function at the detection point. But the paradox of nonlocal EPR[1] correlation between distant events without nonlocal interactions has been more problematic in recent times by Bell's experimental nonlocality test[2,3] proposed in 1964, though the paradox was first noticed by Schrödinger[4] and discussed in the dialogue between Einstein and Bohr[5] at 1935 Solvay Council. In emphasis of the signal transmission in EPR correlation Cavaicanti and Wiseman[6] asked: "What Bohr could have told Einstein at Solvay had he known about Bell experiments ?" In his recollection in 1990 Bell[7] wrote: "Suppose quantum mechanics were found to *resist* precise formulation. Suppose that when formulation beyond FAPP (For All Practical Purposes)[8] is attempted, we find an unmovable finger obstinately pointing outside the subject....to the Mind of the Observer..., or only Gravitation ?" We here show that the solution of quantum paradoxes can be found outside the EPR's reality criterion of local cauality[9] but within the framework of time-symmetric quantum electrodynamics for finite spacetme.[10] A cosmlogical implication of the bi-directional signal transmission P↔Q without common source S in the inflationary cosmology and a possible orgin of dark energy are also discussed.

Author: Senjikan Institute, Niigata, Japan.
e-mail: noboruhokk@yahoo.co.jp

II. EPR Correlation in EPR Loophole

At the Solvay council EPR asked: "Are there spooky actions at a distance in quantum dmechanics ?" Recently, Yin et al.[11] led by Q. Zhang measured a superluminal speed of spooky acions between counter-propagating pair of photons emitted from an opticaly pumped atom in spin 0 state. During the measurement the locality and the freedom-of-choice loopholes of previous expriments were maximally closed by observing a 12-hour continuous violation of Bell's numerical expression (inequality) to EPR's reality criterion of local causality and separability of distant events. Let the spacetime positions of the photon source and the detection points be $S(x_S, t_S)$, $P(x_P, t_P)$ and $Q(x_Q, t_Q)$. Then the lower bound of the speed c_S of the spooky actions

$$c_S = |x_Q - x_P|/|t_Q - t_P| \qquad (1)$$

can be superluminal as $|t_Q - t_P| \to 0$. Here we can see a local and causal link P←S→Q ($t_S < t_P \approx t_Q$) and the nonlocal and acausal (spooky) link P→Q ($t_P < t_Q$). Let ε_P and ε_Q be the unit polarization vectors of photons measured at P and Q. The experiments verified the quantum expectations of the correlation function $C_{QM(}(\boldsymbol{\varepsilon}_P,\boldsymbol{\varepsilon}_Q) = \boldsymbol{\varepsilon}_P\boldsymbol{\varepsilon}_Q = \cos\theta$, where θ is the Hilbert space angle between $\boldsymbol{\varepsilon}_P$ and $\boldsymbol{\varepsilon}_Q$, and showed a clear rejection of classical theories obeying Bell's inequlities. The experiments also confirmed the insenstivty of C_{QM} to observer's delayed decision as to which direction to measure each photon's polarization at P and Q after the photon left the source at S—too late for a message to reach the opposite photon,[8] making the causal link P←S→Q improbable and the bi-directional link P↔S↔Q probable in the loophole of EPR's reality criterion of local causlity between P and Q,

III. EPR Correlation on Double-Light Cone

Dirac[12] defined the two-point corrlation function or propagator Δ(x, t) between S(0,0) and P(x, t), and visualized the signal transmission S↔P on the light cone with the origin S as vertex:

$$(\partial^2/c^2\partial^2t^2 - \partial^2/\partial^2x^2)\Delta(x,t) = 0, \qquad (2)$$

$$\Delta(x,t) = \alpha(t)\delta(c^2t^2 - x^2)$$

© 2016 Global Journals Inc. (US)

$$= [\delta(ct - x) - \delta(ct - x)]$$
$$= \Delta_{future} - \Delta_{past} = \Delta_{ret} - \Delta_{advt}\ , \qquad (3)$$

where $\alpha(t) = t/|t| = 1$ $(t > 0)$; $= -1$ $(t < 0)$. There an electron at S(0, 0) moves under the retarded (causal) action Δ_{ret} of a charged partcle at P on the past light cone $\delta(ct + x)$ of S as well as the advanced (retrocausal) action Δ_{adv} of a charged particle at Q on the future light cone $\delta(ct - x)$ of S, giving a divergence-free radiation damping of the electron at S. The bi-directional EPR link, $P(x_P, t_P) \leftrightarrow S(x_S, t_S) \leftrightarrow Q(x_Q, t_Q)$, can be visualized on the future light cone of the optically pumped metastble atom at S(0, 0) by replacing the step function $\alpha(t)$ by the square (step-up and down) function $\beta(t) = 0$ $(t < t_S)$; $= 1$ $(t_S < t < t_{P/Q})$; $= 1$ $(t > t_{P/Q})$

$$(t_S < t < t_{P/Q}); = 1\ (t > t_{P/Q})$$
$$\Delta(|x_{P/Q} - x_S|, |t_{P/Q} - t_S|)$$
$$= [\delta(c|t_{P/Q} - t_S| - |x_{P/Q} - x_S|)$$
$$-\delta[(c|t_{P/Q} - t_S| - |x_{P/Q} - x_S|)]/|x_{P/Q} - x_S|\ . \qquad (4)$$

The double-light cone[13] $[\delta_{ret} - \delta_{adv}]$ in Eq.(5) tells that the detection point P/Q on the left/right arms of the future light cone of S is reached by retarded wave $\exp i(\omega t - kt)$ from S while advanced wave $\exp i(\omega t + kx)$ from P/Q reaches S on the right/left arms of the past light cone of P/Q, forming a bi-directional sinusoidal wave, $\exp i\omega t \sin kx$, standing in phase between S and P/Q with nodes fixed in space at $x = n\pi/k$ (n = integer).

IV. Bi-Directional Microscope

"Is the star (moon) there when nobody looks" asked Tetrode (Mermin).[14] At the 1947 Solvay Council Heisenberg proposed a thought experiment measuring the electron position on microscope's object plane. There the photon wave collapsing at S in the retinaof the oberver entails the retrocollapse (appearance) of an elecon at P scattering the photon to be observed at S. That is, the electron is not at P when nobody looks. at S. This point was emphasized by Weizäcker[15] in his delayed-choice thought experiment measuring the transverse photon momentum on the focal plane of Heisenberg's microscope. If the microscope is very long, the observer at S can make choice as to which property of the electron, position or momentum, to measure after the scattering process has taken place at P. To see the bi-directional signal transmission S↔P where $M = \rho_\Lambda V = 4\pi\rho_\Lambda R^3/3$ is the Newtonian mass. in microscope we write Eq.(5) in momentum space[14]

$$\Delta_{\omega,k}(|x_P - x_S|, |t_P - t_S|)$$
$$= [\exp ikc|t_P - t_S|\ |\sin k|x_P - x_S|]/|k|, \qquad (5)$$

getting an uncertainty relation between photon momentum $p = \hbar k$ and the microscope length

$$p|x_P - x_S| = nh/2, \qquad (6)$$

where n is the number of nodes of the sinusoidal wave standing between S and P.

V. Cosmological EPR Correlation

In his aether hypothesis Dirac[17,18] considered a nonlocal EPR connection without common source S between distant events P and Q on the 4-dimensional hyperboloid $(ct)^2 - r^2 = l_{pl}^2$, or the Lorenz sphere of radius l_{pl}, crossing the light cone at $ct = l_{pl}$ at $r = 0$ with spacelike velocity:

$$dr/dt = ct/r = c(1 + l_{pl}^2/r^2)^{1/2}. \qquad (7)$$

Hawking[19] considered a cyclic cosmology where an expanding and contracting Lorenz-de Sitter universe starts and ends on the Lorenz sphere embedded into 4-dimensional Euclidean space $\tau^2 + r^2 \sim l_{pl}^2$ with imaginary time $\tau = it$. There the observable Hubble consant $H = v/d$ relating the relative velocity $v \sim c$ of extragalactic objects at a distance d receding from the Earth, is related to the cycle $C = 2\pi H^{-1}$, temperature $T = H/2\pi$ and entropy $S = \pi H^{-2}$. In high dimensional string theory,[20] the parallel orbifold branes collide periodically in cycle, expanding and contracting with dark energy.

We here consider the embedding of the Planck length l_{pl} in the radial line element ds of the Lorenz-de Sitter-Reissner-Nordstroem universe:

$$ds^2 = c^2 g_{tt} dt^2 - g_{rr} dr^2,$$
$$g_{tt} = g_{rr}^{-1} = c(1 - \Lambda r^2/c^2 + l_{pl}^2/r^2)^{1/2}. \qquad (8)$$

Here Λ is the cosmolgical constant interpreted as the timelike vacuum energy density. Note that the light velocity $dr/dt = c(g_{tt}/g_{rr})^{1/2} = c(1 - \Lambda r^2/c^2 + l_{pl}^2/r^2)$ is spacelike at $r \sim l_{pl}$ but decreases with the increase of r towards $dr/dt = c$ at $r = (cl_{pl})\Lambda^{-1/2} \sim 10^{-23}\Lambda^{-1/2}$cm. In the inflationary cosmology starting from quantum fluctuations of preexisting spacetime metric, the radius of the causally related small region extends from $l_p \sim 10^{-33}$cm to $r \sim 10^{-25}$cm during electroweak and grand unification period followed by a brief interlude of reheating, returning to the pre-inflationary temperature of the universe. After the inflationary period, further evolution of the universe is described by the standard Friedman model universe starting the radiation dominated phase of Hubble's expansion history.

VI. Time-Symmetric and Inflationary Friedman .....Universe

Consider a Friedman universe filled with the uniform disrtribution of constant dark energy density ρ_Λ described by the Lorenz-Reissner-Nordstroem metric:

© 2016 Global Journals Inc. (US)

$$g_{tt} = g_{rr}^{-1} = (1 - r^2/r_g^2)^{1/2} + r_{pl}^2/r^2$$
$$= (1 - r/r_g)^{1/2}(1 + r/r_g)^{1/2} + r_{pl}^2/r^2, \quad (9)$$

where $r_g = 3c^2/8\pi G\rho_\Lambda$ is the gravitational radius of the universe determinig the cosmological event horizon. We find superlminal signal trasmission $dr/dt = c(g_{tt}/g_{rr})^{1/2} >> c$ and causally related small regions at radii $r = r_c$ and $r = r_g - r_c$ where $r_c = (r_{pl}/r_g) \sim 10^{-2}$cm for $r_g = R \sim 10^{28}$cm, that is, the inflationary epochs of expanding and contracting almost closed Friedmanan universe, joined onto asymetrically flat Euclidean spaces through double-valued Schwarzschild bottle-neck.

VII. Mass Defect of Semiclosed Friedman Universe

For $r >> r_{pl}$ the evolutinary history of Friedman universe is determined by the Hubble constant $H = 8\pi G/\rho_\Lambda$ where ρ_Λ is the dark energy density. The dimensionless density parameter Ω_Λ is defined by $\Omega_\Lambda = \rho_\Lambda/\rho_{c\Lambda}$ where $\rho_{c\Lambda}$ is the critical density to make the univese flat. For $r >> r_{pl}$ we have

$$ds^2 = c^2 g_{tt} dt^2 - g_{rr} dr^2,$$
$$g_{tt} = g_{rr}^{-1} = (1 - r^2/r_g^2)^{1/2}. \quad (10)$$

Using the integral $\int dx(1 - x^2)^{1/2} = \sin^{-1}x$ the proper mass M_p and volume V_p of the universe of radius R are calculated ss

$$M_p = \rho_\Lambda V_p = 2\pi\rho_\Lambda \int^R r^2 g_{rr} dr = (3/2)(R/r_g)^3[\sin^{-1}(R/r_g)$$
$$... - (R/r_g)(1 - (R^2/r_g^2)^{1/2})]M, \quad (11)$$

where $M = 4\pi R^3\rho_\Lambda/3$ is the Newtonian mass and $M - M_p$ the mass defect. Equation (11) tells that the proper radius $R_p = \int^R g_{rr} dr$ increases with the increase of the world radius from $r \sim l_{pl}$, where $\sin^{-1}(l_{pl}/r_g) \sim 1$, until V_p fills the lower hemisphere of the closed Friedman universe, where $\sin^{-1}(l_{pl}/r_g) = \pi/4$. With further increase of r, R_p decreases towards $R_p \sim l_{pl}$, where $\sin^{-1}(l_{pl}/r_g) \sim \pi/2$, forming a gravitational semiclosure with $V_p \sim M_p \sim 0$ in the evolutionarily earlier upper hemisphere, creating Planck scale black holes[19] outside the event horizon of almost closed Friedman universe joined onto asymptoticlly flat space through Schwarzschil bottle-neck pulsating with Planck scale period.[21]

VIII. Source of Dark Energy

We note that the negative equation of state $\rho_\Lambda + p_\Lambda c^2 < 0$, where $p_\Lambda c^2 << \rho_\Lambda$ is the pressure, required by the dark energy, is satisfied in the upper hemisphere of the closed Friedman universe where the gravitationally bound pairs of quantized metric fluctuations or Planckeons of mass m_{pl}, or graviational Bohr atoms, creating negative attractive potential $Gm_{pl}^2/l_{pl} = \hbar c/l_{pl} = m_{pl}c^2$ dominate, forming Bose-Einstein condensate, while the single Planckeon excitations with positive rest mass energy $m_{pl}c^2$ prevail in the lower hemisphere where the positive equation of state $\rho_\Lambda + p\ c^2 > 0$ is satisfied by the dark matter. The evolutionarily earlier upper hemishere is chracterized by the density parameter $0.5 < \Omega_\Lambda = (R/r_g)^2 = 1$ and the less earlier lower hemisphere by $0 < \Omega_m < 0.5$, The recently updated density parameters[21] fall into these ranges: $\Omega_\Lambda \sim 0.685$, $\Omega_m \sim 0.266$, $\Omega_{atom} \sim 0.049$ where Ω_{atom} is the density parameter of the evolutionarily recent atomic matter. The fact that $\Omega_\Lambda + \Omega_m + \Omega_{atom} = 0.965 \sim 1$ can be taken an indication of the asymptotical flatness of the outer space detecting asymptotic solutions of Einstein equations.

References Références Referencias

1. A. Einstein, P. Podolsky, and N. Rosen, Phys, Rev.**47**, 777 (1935).
2. J. S. Bell, Physics **1**, 195 (1964).
3. A. Aspect and P. Grangier, Proc. Int. Symp. Foundation of Quantum Mechanics, Experiments on Einsten-Podolsky-Rosen Type Correlation with Visible Photons, Tokyo, 1983, pp. 214-224.
4. E. Schroedinger, Sur la théorie relativiste de l'électron de la méchnique quntique. Annals of the Institute of Henri Poincré **2**. 269-310 (1932).
5. N. Bohr, Phys. Rev. 48, 696 (1935).
6. E. G, Cavaicanti and H. M. Wiseman, Found Phys. 6 July (2012). (on line).
7. J. S. Bell, Physics World, **3**, 33 (1990).
8. R. Penrose, The Shadows of the Mind, Oxford University Press, Oxford (1989).
9. N. Hokkyo, Studies in History and Philosophy of Modern Physics **39** (2008) 762-766.
10. E. C. G. Stueckelberg, Phys. Rev. **81**, 130 (1951).
11. Juan Yin et al. Phys. Rev. Lett. **110,** 260407 (2013)
12. P. M. A. Dirac, The Principles of Quantum Mechanics, Oxford University Press (1947)
13. W. Heitler, The Quantum Theory of Radiation, Oxford University Press (1953).
14. H. Tetrode, Zeits. für Phys. **10**, 317 (1927).
15. K. F. Weizäcker, Zeits. für Phys. **A70**, 114 (1931).
16. O. Costa de Beauregard, Am. J. Phys. **51,** 5 (1983).
17. P. A. M. Dirac, Proc. Roy. Soc. **180A**, 149 (1947)
18. P. A. M. Dirac, Nature, **169**, 702 (1952)
19. S. Hawking and R. Penrose, The Nature of Space and Time, Princeton University Press, Princeton (1996).
20. E. Witten, Nucl. Phys. Phys. B44, 85 (1995).
21. N. Hokkyo, Physics and Space Science **14**, 1 (2014)
22. D. Overbye et al."New Image Reform Views in Infant Universe," Max Planck Meeting held on December 1, 2014 at Ferrela, Italy.
23. N. Hokkyo, Physics and Space Sciences **16**(2) (2016).

© 2016 Global Journals Inc. (US)

This page is intentionally left blank

© 2016 Global Journals Inc. (US)

Global Journal of Science Frontier Research: a
Physics and Space Science
Volume 16 Issue 3 Version 1.0 Year 2016
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

The Dance of the Spinning Top

By Álvarez Martínez, Alejandro & Martín Gutiérrez, Almudena

Abstract- An interpretation of the spinning top movement, the so-called "dance of the spinning top" is proposed. It is based on a mechanical model without a central force, owing to the absence of the same, in spite of the fact that there is curvilinear movement. The path can be interpreted by means of the *Theory of Dynamic Interactions.* This interpretation is valid regardless of the way in which the spinning top has been activated and the manner in which it is bound by the environment.

Following on from the foregoing, we suggest that planetary movement around the sun or that of artificial satellites around the Earth and, in general, all objects endowed with angular momentum on one of their rotating axes in translation and subject to non-coaxial external momenta, could also be interpreted in this way.

Keywords: *Theory of Dynamic Interactions, interaction, inertial field, velocity field, torque, orbiting, angular momentum, angular speed, spinning top, planets, satellites.*

GJSFR-A Classification : *FOR Code: 020399*

THEDANCEOFTHESPINNINGTOP

Strictly as per the compliance and regulations of :

© 2016. Álvarez Martínez, Alejandro & Martín Gutiérrez, Almudena. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

The Dance of the Spinning Top

Álvarez Martínez, Alejandro [α] & Martín Gutiérrez, Almudena [σ]

Summary- An interpretation of the spinning top movement, the so-called "dance of the spinning top" is proposed. It is based on a mechanical model without a central force, owing to the absence of the same, in spite of the fact that there is curvilinear movement. The path can be interpreted by means of the *Theory of Dynamic Interactions.* This interpretation is valid regardless of the way in which the spinning top has been activated and the manner in which it is bound by the environment.

Following on from the foregoing, we suggest that planetary movement around the sun or that of artificial satellites around the Earth and, in general, all objects endowed with angular momentum on one of their rotating axes in translation and subject to non-coaxial external momenta, could also be interpreted in this way.

Keywords: Theory of Dynamic Interactions, interaction, inertial field, velocity field, torque, orbiting, angular momentum, angular speed, spinning top, planets, satellites.

I. Introduction and Grounds

The rotary movement of bodies is a phenomenon that has hitherto been studied insufficiently. Interest in this area dropped in 20th century only to focus on other aspects of physics that were considered to be of greater import at the time.

Notwithstanding, for some years now, the lecturer Gabriel Barceló Rico-Avello, Doctor in Industrial Engineering and Physics Graduate has been proposing a rotational mechanics that we believe better responds to the behaviour of bodies with intrinsic rotation. We refer to the *Theory of Dynamic Interactions* (TDI), which explains that the rotary movement of a body endowed with intrinsic rotation interacts with the translation of the same. *Dynamic interactions are understood as those inertial reactions that appear when a body is subject to accelerations* [1].

There is a host of rotating bodies that could serve as a subject of study. Nonetheless, this paper restricts its scope to that of the spinning top and, more exactly, to the *dance of the spinning top*, which consists of the curvilinear translation movement made by this body when rotating on its symmetrical axis, which in turn, forms a certain angle with the vertical axis perpendicular to the supporting plane. Moreover, we propose to show that the aforementioned *dance of the spinning top*, is in keeping with the basis of the *Theory of Dynamic Interactions* (TDI).

Intuition would suggest a similarity between this *dance of the spinning top* and the movement of any planet around the sun or that of any artificial satellite orbiting the Earth. Consequently, the conclusions to be gleaned from this behaviour will also be extrapolated to these cases of scientific interest.

Furthermore, other particular instances of the spinning top will be analysed to justify the generalisation of this theory and to show that the interaction is independent of the type of effect that provokes the movement, as long as there is rotation and translation.

II. Background

As stated by Dr. Barceló Rico-Avello [2], the peculiarity of a rotating body also warranted the attention of some 20th century scientists. Accordingly, G. Bruhat [3] defined it as the *apparently paradoxicalgyroscopic effect*. Likewise, Paul Appell [4] on referring to rotating bodies also frequently insisted on its paradoxical behaviour. Nonetheless, the principles of classical mechanics failed to delve into this aspect.

We believe the original error in the formulation of the theory of the scientific community as regards rotating bodies could lie in the studies of the scientist Louis Poinsot (1777-1859) from two perspectives.

Firstly, he studied the rotation movement with a fixed point (the case of the spinning top were translation not caused on its point of support on the surface) [5].




Figure 1 : Spinning top rotation axis and the angle it forms with the vertical

In this analysis he proposed that the *Instantaneous Axis of Rotation* defined by D'Alembert (1717-1783) [6] varied at each moment owing to the movement of the body, forming a cone with a fixed point vertex. However, he further claimed that the axis of rotation also varied inside the body itself. We hold that this starting point is erroneous. From experiments on the rotation movement of spinning top it can be observed that, by taking certain fixed axes on it, its axis of rotation does not shift or rotate in any way with respect to them.

Author α σ: Advanced Dynamics, C.B.
e-mail: gestor@advanceddynamics.net

© 2016 Global Journals Inc. (US)

That is to say, as can be seen in figure 1, angle α remains constant.

Secondly, it must be pointed out that Poinsot shows that forces and momenta act on a solid body. However, he further states that "*each one of these dynamic elements produce their forces separately; force determines the translation movement of the body while torque generates rotation around its centre of gravity, without the force exercising any influence*" [7]. Such a reading had already been put forward by Lagrange [8].

This idea, which comes from Classical Mechanics, is the one that has established the method to calculate paths that is still used today. Notwithstanding, we suggest that dissociating translation and rotation in a body where both phenomena occur jointly is mistaken and that the *Theory of Dynamic Interactions* [9] better reflects what really happens, as indicated by Julio Cano in his paper on the pendulum of dynamic interactions. [10]

Other antecedents are known to have questioned the idea of separation. Such is the case of Dr. Shipov, who claims in the context of dynamic energy transfer:

> *Consequently, the law of conservation establishes a connection between translational and rotational forms of movement and makes it possible to transform the translation momentum in a rotational one and vice versa. This means that the law of conservation of momentum in Newtonian mechanics is limited and cannot be generalised to the case of the torsion interactions of colliding bodies.* [11]

Classical Mechanics uses differential equations to calculate paths, the well known "Euler Equations" [12]. Nevertheless, these expressions cannot be integrated even for the case of the movement of the spinning top and, therefore, from this limitation, the need to propose other new ones to provide results for this case arises. Moreover, it must be borne in mind that Newtonian Classical Mechanics is limited to inertial cases. Accordingly, to be able to apply its formula, Coriolis found it necessary to introduce the *Coriolis force* and the *centrifugal force* as fabricated forces for non-inertial cases [13]. Nonetheless, in the *Theory of Dynamic Interactions* we are proposing, there is no need to incorporate fabricated or apparent forces, as the dynamic behaviour of bodies is defined by the coupling of the dynamic fields generated.

Lagrange managed to express the components of angular velocity along the main axes and in function of Euler angles, but he only solved equations for a particular case and was unable to obtain a generalisation or to resolve the case of the spinning top, which is the one that concerns us here. [14].

Subsequently, Sofía Kovalévskaya in her paper entitled *On the rotation of a solid body about a fixed point* [15] attempted to resolve the Euler equations. However, the equations generated four algebraic integrals that it proved impossible to resolve by integration, so she proceeded to resolve them numerically by means of an approximation procedure, thus leaving the general resolution of the equations unresolved.

Some recent studies have been found on the *dance of the spinning top* which in our opinion fall short. Christopher G. Provativis of the Technical University of Athens, published a paper that studies this movement based on the Euler equations and Lagrange's Analytical Mechanics from which he obtained equation systems that cannot be integrated analytically and which he resolved by means of numerical integration, thus arriving at mere approximations. [16]

We came across another thesis in which, on the basis of Lagrange equations and by means of variation calculation, the path of a spinning top in its dance is obtained and visualised. This work compares the different results obtained by different discretizations, but always from the starting point of Lagrangian mechanics [17].

Given the impossibility of resolving the movement equations with the mathematical tools at our disposal and which prevent the confirming of their truthfulness and the relationship between rotation and orbiting, Dr. Barceló proposed the Theory of Dynamic Interactions and a mathematical formula that explains the behaviour of rotating bodies [18].

Indeed, on studying the reactions of a solid rigid body subject to non-coaxial torques, M. Hirn had already concluded in the 19th century that these torques were generating non-homogeneous velocity fields [19]. The *Theory of Dynamic Interactions* also covers the distribution of non-homogeneous fields.

Lastly, it is noteworthy that the Spanish physicist, Miguel A. Catalán, also interested himself in the movement of rotating bodies in the middle of the 20th century, comparing the rotating of an electron around the atom nucleus to spinning top movement [20]. Indeed, he is the lecturer that inspired Dr. Barceló Rico-Avello to develop the theory set forth in this paper.

III. "The Dance of the Spinning Top" and the Theory of Dynamic Interactions

When a spinning top starts to move, it does so owing to a torque acting on it that provokes its rotation.

If the axis of rotation of the spinning top remains perpendicular to the support surface this body stays spinning on a fixed point without shifting. However, what happens when this perpendicularity no longer holds?

By observation, it can be deduced that the spinning top will continue its rotating movement with respect to its rotation axis, which coincides with its symmetrical axis, and will move on the surface tracing a curved path. In this case, it is possible to speak in terms of displacement physics, given that the spinning top

© 2016 Global Journals Inc. (US)

point is not fixed on the surface, i.e., it does not have zero speed but rather shifts position.

This curved path, under surface support and smooth displacement conditions (uniform friction coefficient) will be closed and, the more ideal the conditions in which the experiment is being conducted (absence of air) the closer it will come to tracing a circular path.

We have come across historical descriptions of this so-called *spinning top dance* behaviour accompanied by illustrations reflecting this movement (figure 2), as well as descriptions that explain that the spinning top would rest on the support surface if it was not for its rotation (figure 3). [21].

According to Classical Mechanics, a body in movement traces a curved path if there is a force that modifies its straight movement towards its inside at each point of movement (centripetal force). However, the forces acting on the spinning top are weight and the normal reaction with the surface, both of which are perpendicular to the support plane. Neither of them are, therefore, central forces.




Figure 2 : Dance of the spinning top.
Source: Perry, J. (2010). Spinning Tops. (pg. 21)




Figure 3 : Spinning top with and without rotation
Source: Perry, J. (2010). Spinning Tops. (pg. 21)

What, therefore, causes this path? We suggest that the reason for it is to be found in the momentum generated by the weight of the spinning top with respect to the point of support, which will provoke the overturning of the spinning top but this, instead of falling, if its support point allows it to slide, will start this dance. (See video: www.advanceddynamics.net/spinning-top-video/). [22] This idea can be seen in figure 4. It is the same effect as produced on the flight of the boomerang, the rotational and orbital movements of which have also been studied. Dr. Barceló suggests in this case:

The weight, a force applied at the boomerang's centre of mass, will not coincide with the resulting lift force, determining a couple that acts at the same time with the rotation, but without being coaxial with it. […] The couple will generate a rotation momentum that tends to tilt the boomerang around its flight direction axis. […] The weight couple and that resulting from the lift forces, which is non-coaxial with the intrinsic rotation of the boomerang, will be the dynamic interaction couple that generates the new path. [23]

© 2016 Global Journals Inc. (US)

This situation (see video) is possible owing to the intrinsic rotation of the body, given that were there no rotation, the spinning top would fall onto the surface.

Accordingly, the interaction between the rotation movement and the translation of a body has been explained.




Figure 4 : Forces acting on the spinning top

IV. Similarity with the Planets Orbiting the Sun

Euler [24] was the first to announce the similarity between the Earths' movement around the sun and that of the spinning top. What we intend to do in this paper is not solely to reaffirm this similarity, but rather to explain that the reason behind the Earth's orbit resembles that of the "dance of the spinning top".




Figure 5 : Forces acting on the Earth according to Newton

Source: http://spaceplace.nasa.gov/review/dr-marc-solar-system/planet-orbits.sp.html

Isaac Newton [25] explained that the Earth's orbit around the sun was caused, on the one hand, by a tangential acceleration and, on the other, by a central force, the sun's force of attraction that acts as a centripetal force and which causes the body to deviate from its straight path, thus converting it into a curved one. In this way, Newton also explained why the Earth, or any of the planets, is at a certain distance from the sun and not stuck to it by virtue of its gravitational force. This idea can be seen in figure 5.

However, Newton's explanation of this orbit is brought into question by the *Theory of Dynamic Interactions.* This theory holds that, as is the case with the spinning top, the orbit of any planet around the sun is not necessarily caused by a central force, but rather by the momenta of forces. This is so based on the first corollary of the tenth law that underpins this theory: "*It is possible to infer the existence of inertial dynamic interactions in rotation and orbital phenomena from the momenta of forces.*"[1]and[26]

Obviously the dance of the spinning top is not caused by any central force, owing to the inexistence of the same, as has been explained in the previous section. However, this idea is not as obvious in the case of a solar system planet because the existence of the force of attraction of the sun is a reality. What follows is an attempt to explain and justify it.

The Earth or any planet is, as are the majority of bodies in space, rotating while at the same time moving in translation. As indicated in the principle of inertia formulated in Newton's laws [27], if there is no effect to impede it, such bodies would continue to move in a straight line. The planets orbit around the sun, thus obviously there must be something that provokes the curved path. Accordingly, the *Theory of Dynamic Interactions* is being proposed as a conceptual framework within which to explain the mechanics of celestial bodies understood as rigid solids subjected to successive non-coaxial rotations. [28]

The *Theory of Dynamic Interactions* holds that the orbit is not caused by the gravitational force of the sun, given that, were this the case, the planets would be

© 2016 Global Journals Inc. (US)

stuck to it, but rather by the torque exerted by the sun on the centre of the mass of planets, as a result of its position not coinciding with the centre of the same.

On account of the non-homogeneous constitution of planets, the centre of mass of these is displaced as regards their geometrical centre. Accordingly, the force of the sun's attraction, which will have the direction of a vector that passes through the centre of mass of the sun and that of any planet, will provoke a momentum on the geometrical centre in such a way that the planet, with intrinsic rotation, is subjected to a new non-coaxial momentum. The TDI accepts that the velocity field generated by this new, non-coaxial momentum couples with the translation velocity field thus causing the new orbital path around the sun.

Hitherto, the resemblances have been shown between the spinning top movements and that of the planets. An important difference must now be highlighted: the rotation axis of the dance of the spinning top is inclined with respect to the path. This means that, owing to the effect of the weight, the rotation axis of the spinning top varies with respect to the inertial reference system by tracing a cone. In the case of the Earth, its axis of rotation maintains practically the same inclination with respect to the inertial system along any point of its orbital path. Of course, there will be variations, but much slower and imperceptible, taking some 25,800 years to trace a complete cone. This idea can be seen in figures 6 and 7.

We claim that the oscillation of the Earth's axis, a phenomenon known as *Chandler wobble*, is caused by gravitational forces, as in the case of the spinning top. It must basically be the gravitational force of the moon that makes the axis of rotation of the Earth so stable and showing such extremely slow variations.




Figure 6 : Situation of the Earth in orbit at several positions.
Source: www.geoenciclopedia.com/movimientos-de-la-tierra/




Figure 7 : Situation of the spinning top during its "dance" at several positions

V. Similarity with Artificial Satellites Round the Earth

Even though the case of an artificial satellite orbiting around the Earth resembles that of the Earth around the sun and, therefore, that of the spinning top dance, there will be differences as regards the mechanism that keeps the satellite orbiting around the Earth.

© 2016 Global Journals Inc. (US)

Dr. Barceló Rico-Avello already expounded his intention to extend the Theory of Dynamic Interactions in *Un mundo en rotación* (A Rotating World) [18] to communication satellites which, having an initial speed and endowed with intrinsic rotation if subject to a non-coaxial torque, will assume an orbiting movement.

Obviously, for a satellite to orbit around the Earth at a certain height over its surface, a launch vehicle is required to generate the thrust that will position it at that height. Moreover, it is also clear that, if at that height the thrust no longer exists, the satellite will return to the surface of the Earth, impacting against it, owing to the Earth's force of gravity. Consequently, some type of propellant is needed to generate a force parallel to the Earth's surface that will be sufficient not to represent a simple launch that eventually "spends itself" and returns to the Earth's surface on account of the gravitational action. This force is called thrust.

The thrust will suffice if it provides the satellite with the required orbital speed at that height to orbit the Earth and will be determined by the energy equation. This speed will depend on the Earth's mass [29].

The thrust can be exerted in two different ways, depending on whether it is produced by a motor located on the axis of symmetry of the satellite or from several situated around the same and structurally attached to it. In both cases, a non-coaxial momentum will be caused with respect to the axis of symmetry of the satellite, given that its centre of mass is not going to be located exactly on this axis owing to the heterogeneity peculiar to all satellites, which is inevitable. This will make the satellite rotate owing to the momentum caused and, in turn, endow it with a translation movement on account of the thrust, thus generating a path that will be curved; once again accounting for the *Theory of Dynamic Interactions* and clearly showing that, as is the case with the dance of the spinning top, the movements are not caused by a central force, even though in this case the Earth is indeed the centre of the movement.

Lastly, it must be pointed out that satellites are stabilised by different mechanisms to undertake the mission for which they have been launched and thus are not orbiting with the described characteristics.

VI. Interaction of Fields on the Spinning Top

In all movements there is another physical vectorial magnitude that acts about which no mention has yet been made, namely, translation velocity. In this section, the velocity fields are analysed that are present in the "dance" movement of the spinning top along with inertial fields and, with them, it will be explained how the moving spinning top maintains its position as it does. (See video).

The foregoing is based on the following idea: *"orbital movement is an inertial reaction caused by the distribution of speeds and accelerations that are generated by the action of the second external torque that is acting."* [18]

This idea is based on the four axioms [1] that were announced after long years of research and on which the later field analysis is based:

1. The rotation of space determines the generation of fields.
2. Result of the action of non-coaxial momenta.
3. Inertial fields cause dynamic interactions.
4. The resulting action of the different successive, non-coaxial torques that act on a rigid body cannot be determined by an algebraic sum or be calculated by means of an alleged resulting torque.

The *Theory of Dynamic Interactions* holds that when a rigid solid is subjected to the successive couplings of forces, the first momentum will cause its intrinsic rotation (in this case the rotation of the spinning top on its axis of symmetry) while the subsequent momenta will generate the non-homogenous velocity fields [23].

If the spinning top is placed vertically on its axis and it is endowed with a rotary movement, it will be left rotating on its axis of symmetry and will take longer to fall the faster it rotates. When the spinning top starts to rotate it will also start its dance, even though it has not been endowed with a thrust that provides it with a quantity of movement. It can be understood that at the initial moment, the axis of symmetry undertakes a falling movement endowing the centre of mass with an initial linear velocity, and finally, the spinning top will rotate and orbit at the same time. It can also be supposed that the spinning top's (or those of any rotating body) angular momenta do not add up algebraically or vectorially, resulting in a final momentum and therefore, no general law of the composition of momenta is obtained [18].

Firstly, (see video), a study is made of the fields to which the spinning top is subject at a particular instant of its movement, that is to say, on certain fixed axes to the same, without considering translation.

Figure 8 shows in blue the instantaneous inertial field generated initially on a spinning top that is tilted owing to the effect of the torque action of the forces present on the spinning top: the weight and reaction of the ground on the spinning top. This field will be perpendicular to the radius of the tilt and tangential to the tilt path.

© 2016 Global Journals Inc. (US)




Figure 8 : Inertial fields on the spinning top (without considering its translation movement).

Nevertheless, were only this inertial field to exist, the spinning top would fall and impact against the ground. Therefore, the velocity and inertial fields are coupled thus causing its new path. This has also been observed and mentioned in other studies on bodies in which dynamic interactions are to be found [18] [30].

The body is endowed with intrinsic rotation on its axis of symmetry, which is going to cause a linear velocity on each point of the spinning top. As we are dealing with a rigid solid, this linear velocity value will be constant at each one of its points on the same disc at the same time.

Moreover, the tilt velocity generates a velocity that makes the spinning top fall onto its support surface. The velocity field in yellow in figure 9 that is tilted 90° with respect to the instantaneous tilt velocity is precisely the one that is going to prevent the overturning of the spinning top.

The maximum tilting velocity in the downward direction will correspond to the closest point to the surface (red point in figure 10). The maximum upward tilting velocity will be at 180° from this. Nevertheless, in the first revolution of the spinning top, the amounts of movement will be summed up in successive instants at each point on it. Accordingly, the initial maximum tilting velocity at the following instant will have been shifted by the rotation of the spinning top and the tilting velocity corresponding to its new position will be added to this tilting velocity leading to a change in velocity value. Therefore, on completing the first revolution, the tilt velocity field will have been displaced 90° thus giving rise the situation in figure 9. This situation will hold as long as the spinning top is in a state of dynamic stability. This succession of states until attaining stability (figure 9) can be seen in figure 10.




Figure 9 : Resulting velocities fields (yellow) and tilt velocities fields (blue) on the spinning top (without considering its translation movement)




Figure 10 : Spinning top tilt velocities from the initial instant to maximum stability

© 2016 Global Journals Inc. (US)

It must be pointed out that, owing to friction with the surface and the Magnus effect, the linear velocity that caused rotation will begin to fall. The translation of the velocities field will, then, be contrary to that which was seen when it was increasing, reaching a moment in time in which it coincides with the situation at the initial instant and which causes, therefore, the spinning top to overturn.

In the light of the foregoing, it is understood that the spinning top movement passes from a transitory state in which the linear velocity of rotation of the spinning top increases to a dynamic stability in which this linear velocity is constant, until this speed falls entering once again in a transitory state until it comes to nothing and the spinning top is left overturned on the surface. The states of equilibrium correspond to states of spinning top stability, on the understanding that the concept of equilibrium is not restricted to "static equilibrium". According, Gabriel Barceló claims as follows: "*In accordance with the Theory of Dynamic Interactions proposed, we conceive a universe in dynamic equilibrium in which a constant orbit movement acts inside a closed path.*" [1]

However, it must not be overlooked that the linear translation velocity acts at the same time as the tilting velocity. The yellow velocities field couples with the system's translation velocities field thus generating the movement that is observed. The variation in the time of the two velocities fields gives rise to acceleration fields that serve to generate a force field. The force of these two fields is the effective force produced on the system.

Take the following axes (see video): the green triad symbolises the translation reference system of the solid, where the vector T indicates the direction of translation from the centre of mass of the solid, while R indicates the radial direction that points towards the centre of the curve traced by the dancing spinning top. The yellow triad symbolises the reference system without translation with respect to the foregoing, but tilted in accordance with the inclination of the body to the vertical plane of the Earth. These systems of axes are represented in figure 11.




Figure 11 : Systems of axes on the spinning top

Based on the analysis conducted and the observation of the rotary and translation movement of the spinning top, (see video) the proposed axioms are duly corroborated and in the light of the coupling of the fields of velocities it can be concluded that: "*the body will keep to a closed path (e.g., a an orbital movement), without any need for a central force, and will simultaneously intrinsically rotate around its initial axis.*"[31].

VII. Magnetic Spinning Top




Figure 12 : Magnetic spinning tops

Source: http://cienceandoenelsulayr.blogspot.com.es/2014/09/inicio-del-curso-2014-2015.html

The magnetic spinning top, save for those that can levitate on account of the magnetic repulsion effect, behaves similarly to the conventional spinning top in terms of movement and, therefore, can only be distinguished from it because of the way in which it is activated and in the variations of angular velocity to which its rotation may be subject, owing to the variations in the magnetic forces applied. The spinning top must be magnetised to apply the external forces. These ideas about the magnetic spinning top have already been expounded by Gabriel Barceló in his book *Un mundo en rotación*(A Rotating World) [18]:

> *The disc would be activated remotely by a magnet. Magnetic spinning tops subjected to different external magnetic forces confirm the action of external magnetic torques in the variation of the rotating and orbiting velocity owing to the magnetic induction effect. This is also the case of a magnetic spinning top that keeps a constant rotating movement owing to the effect of an electromagnet acting by impulses, or which levitates in a magnetic field. In these cases, the orbital path generated has been confirmed for different rotating velocities and different levels of friction with the support surface.*

The foregoing, therefore, leads to the following conclusion: it is possible to propose a generalisation of the *Theory of Dynamic Interactions* for any body endowed with intrinsic rotation, regardless of the way in which this rotation is activated or the type of the external

© 2016 Global Journals Inc. (US)

forces and momenta acting on it. We have now seen several cases: the conventional spinning top, planets, satellites and magnetic spinning top in which this theory is confirmed and in different circumstances the reason why the intrinsic rotation was different.

VIII. TDI Equations on the Spinning Top

The paths of several, different sized spinning tops with different masses have been traced to check compliance with the mathematical formula for the *Theory of Dynamic Interactions* in the case of the spinning top; paths that have been calculated applying the TDI. The results obtained are shown below.

The mathematical formula for the *Theory of Dynamic Interactions* is as follows[30]:

$$\vec{v} = \Psi \vec{V_0}$$

Where Ψ is a rotation matrix defined as follows:

$$\Psi = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos\alpha & -\sin\alpha \\ 0 & \sin\alpha & \cos\alpha \end{bmatrix}$$

Where $\alpha = t\frac{M'}{I\omega}$, M' is the external momentum, I the body's inertial momentum, in this case of the spinning top and ω the angular velocity of the rotating body [18].

Figure 13 shows two images of the first experiment. The image on the left shows the real path followed by the spinning top, while the one on the right, the path calculated according to TDI, considering the deceleration caused by dissipative forces (surface friction). This first experiment was done as follows: a spinning top was activated on a wooden table surface by means of a torsion spring. A video was recorded of the spinning top in movement and later images were captured at intervals of a second. Accordingly, the different positions of the spinning top were located during movement, which are the points plotted on the graph to the left of figure 13.

It can be seen that both paths are similar though the second one will be more perfect on having considered the dissipative forces causing an homogenous deceleration, when in reality this will depend on the friction coefficient at each surface point, which may not be homogenous.




Figure 13 : Real path of the spinning top and the path calculated by TDI

Another experiment was conducted to assess the TDI results. In this experiment it was possible to obtain a high number of path points at halfway along the revolution of the dancing spinning top. These points were used to produce a graph that was superimposed on the one obtained by using the TDI. The result of this can be seen in figure 14.A second experiment was designed to obtain a greater number of points. This time, the spinning was activated as before, but it was placed on a sheet covered with powder to trace the path taken.

© 2016 Global Journals Inc. (US)




Figure 14 : Superimposition of real spinning top path graphs and the one calculated in accordance with the TDI (I)

Figure 14 shows the real movement of the spinning top at the start of the same. At this moment in time, the linear velocity begins increasing, which is reflected in the increase of the radius until such time as the dissipative effects cause it to diminish as can be seen in the graph. The difference between the real path and the calculated one arises from the difference in the coefficient of friction on the support surface, because as stated above, this was a sheet covered in powder. Moreover, this gap between the real path and the calculated one came about because the surface had a slight inclination that caused the increase of the radius.

A third experiment was done to reduce the difference among friction coefficients throughout the surface. This obtained the path of a complete revolution owing to the spinning top being able to paint on the surface travelled. However, this spinning top was of greater mass than the previous one so its path radius was smaller. This explains why less points were obtained and greater accuracy attained as regards the tracing of the path. Consequently, in figure 15, a path can be perceived that resembles the one calculated by the TDI, but which has gaps owing to the paucity of the data sample.




Figure 15 : Superimposition of real spinning top path graphs and the one calculated in accordance with the TDI (II)

In the light of the foregoing, it has been seen that the *Theory of Dynamic Interactions* manages to closely approximate the path of the spinning top and surpasses that of the Euler equations that did not allow for integration with respect to the case of the spinning top.

IX. Angular Momentum on the Dance of the Spinning Top

We also analyse the conservation of angular momentum on the movement of the spinning top. The angular momentum is transferred to the spinning top by

© 2016 Global Journals Inc. (US)

means of a thrust that is generated on applying a torque in several ways [32].

As regards this issue, the following reflection by Dr. Barceló Rico-Avello must be mentioned:

> *This behaviour of nature can also be interpreted in the field of physics by the fact that, under such circumstances, **angular momenta are not added up**, so if a body is subject to an angular momentum **L**, its variation ΔL, due to the effects of external forces, will not necessarily be added to the already existing **L**, but rather can generate a new movement, different and simultaneous to the existing one, which we have called **precession movement** or, where appropriate, **orbital movement.** This would be the case of the spinning top, and in the case of a body in space without any restraint, like a boomerang, or numerous other like objects, the body will begin to orbit without the necessary existence of a central force.* [18]

In the case of the spinning top it is possible to differentiate three rotations (figure 16).

The inertial momentum of the spinning top does not vary given that we are dealing with a rigid body and, therefore, it will always have the same mass and be distributed in the same way. Consequently, the variation in the angular momentum is going to be determined by the angular velocity and will be constant in each axis system if the angular velocity in these is also constant.

The conservation of the angular momentum in the rotating of a spinning top (with respect to 0 axes) has already been expounded by A. Fernández-Rañada:

> *When the aforementioned angular velocity is high, the inertia of the spinning top due to its spinning on its own axis is very high and the external torque does not suffice to appreciably change I ω (…). Therefore, the **L** value of the angular momentum remains constant. As must be the conclusion that the angle formed by the axis of rotation and the Z axis must also be constant. In order for these two conditions to be met in the presence of the M torque, the axis of rotation of the spinning top has to trace a cone solid of revolution thus causing a precession movement around the Z axis.* [33]




Figure 16 : Illustration of rotations in the "dance of the spinning top"

X. Future Lines of Research

The most important issues have been dealt with throughout this paper that are considered essential to showing how the *Theory of Dynamic Interactions* explains the dance of the spinning top and other bodies that behave similarly in reality and that may be of interest to the reader. We now deal with other issues related to the subject matter discussed, albeit outside the scope of this paper, but which point to future lines of research.

We know that the Earth makes the following movements: rotation on its own axis, orbiting around the sun, equinoctial precession, along with another we have had cause to mention: the Chandler wobble [34]. It would be interesting were future research to study the latter movement in relation to the others. In the "*El vuelo del bumerán*" (The Flight of the Boomerang) [2], Gabriel Barceló has already hinted at this subject:

> *In 1758, Euler predicted that the axis of rotation of the Earth also had a further movement with respect to a fixed frame of reference. In 1891, Chandler determined the time interval of this free polar movement. The oscillation radius of the Chandler wobble on the Earth's rotation axis is around 6 m.*

It is also proposed that research be done to confirm or refute Poisont's [5] claim that there is an instantaneous axis of rotation that is different at every instant of the movement, both on inertial axes, as well as

© 2016 Global Journals Inc. (US)

on those axes bound up with the solid. This seems to coincide in the first case, but not in the latter, given that this would mean having a different axis of rotation to the axis of symmetry at every moment of rotation, whereas observation seems to reflect otherwise.

We also suggest research be done into the technological applications that could be developed on the basis of the *Theory of Dynamic Interactions* proposed by Dr. Barceló in previous papers. [35].

Lastly, we suggest that the reader have another look at the video to better understand the behaviour of the spinning top: www.advanceddynamics.net/spinning-top-video/

You can consult the following links for further information on the *Theory of Dynamic Interactions:*

http://advanceddynamics.net/
http://dinamicafundacion.com/

References Références Referencias

1. Barceló, G. *Una historia de la concepción humana del cosmos. II. Del Renacimiento a nuestros días.* Ed. Arpegio: Barcelona, 2013.
2. Barceló, G. *El vuelo del bumerán.* Ed. Marcombo: Barcelona, 2005.
3. Bruhat, G. *Mécanique* (París, 1955).
4. Appell, P. *Traité de Mécanique Rationnelle* (Paris, 1909).
5. Poinsot, L. *Théorie générale de l'équilibre et du mouvement des systèmes* (1806).
6. D'Alembert. *Recherches sur la précession des équinoxes et sur la nutation de l'axe de la terre.* Paris, 1749.
7. Poinsot, L.: *Connaissance des temps. Théorie nouvelle de la rotation des corps*. Paris 1852.
8. Lagrange. *Mécanique analytique,* t. I, p. 230.
9. Barceló, G.:*Analysis of Dynamic Fields in Non-inertial Systems.* World Journal of Mechanics, Vol. 2, No. 3, 2012, pp. 175-180. http://dx.doi.org/10.4236/wjm.2012.23021
10. Cano, J. (2015). The Pendulum of Dynamics Interactions. *Journal of Applied Mathematics and Physics*, *1* (3), 1186-1198. DOI: 10.4236/jamp.2015.39146http://www.scirp.org/journal/jamp
11. Shipov, G. (1998) *Una Teoría sobre el vacío físico: Un nuevo paradigma.* Academia de Ciencias Rusa, Instituto Internacional de Física Teórica y Aplicada, Editorial URSS, UR, Moscú.
12. Euler, L. *Du mouvement de rotation d'un corps solide quelconque lorsqu'il tourne autour d'un axe mobile.* Berlin, 1760.
13. Coriolis, G. de. *El nacimiento de la nueva física.* 1989.
14. Lagrange. *Mécanique Analytique.* Tomo I, Edición M. Bertrand. París, 1788.
15. Kovalévskaya, S. K. *Sobre la rotación de un cuerpo sólido sobre un punto fijo.* 1888.
16. Provativis, C. G. (2012) Revisiting the Spinning Top. *International Journal of Material and Mechanical Engineering*, 1 (1), 71-88. DOI:www.ijm-me.org/Download.aspx?ID=2316,www.ijm-me.org
17. Bodack, R. (2008). *Dynamics of the Spinning Top: Discretization and Interactive Visualization.*Diploma Thesis. Technische Universität Berlin. Institut für Mathematik.
18. Barceló, G. *Un mundo en rotación.* Editorial Marcombo: Barcelona, 2008
19. Hirn. M.: *Théorie analytique élémentaire du giroscopio*; en los Annales de l'Observatoire de Paris, t. IX, citado en Gilbert, P.: *Problème de la rotación d'un corps solide autour d'un punto*. Anales de la société scientifique de Bruxelles. 1876.
20. Catalán, M.A y León, A. (1945). *Física Y Química*, séptimo curso.
21. Perry, J. (2010). *Spinning Tops.*
22. Pérez, L. A. and Alvarez, A.: *The dance of the spinning top.* Video script, Valladolid, 2015.
23. Barceló, G. (2014). *Theory of dynamic interactions: The flight of the boomerang.* Journal of Applied Mathematics and Physics.http://dx.doi.org/10.4236/jamp.2014.27063.http://www.scirp.org/journal/jamp
24. Euler, L. *Du mouvement de rotation des corps autour d'un axe variable.* Berlin, 1758.
25. Newton, I: {http://spaceplace.nasa.gov/review/dr-marc-solar-system/planet-orbits.sp.html}
26. Barceló, G. Theory of Dynamic Interactions: Laws of Motion. *World Journal of Mechanics*, 3, 328-338. (2013) http://dx.doi.org/10.4236/wjm.2013.39036
27. Newton, I: *De analysi per aequationes numero terminorum infinitas,* 1711.
28. Barceló, G. (2013). Proposal of New Criteria for Celestial Mechanics. *International Journal of Astronomy and Astrophysics.3* 385-391. http://dx.doi/org/10.4236/ijaa.2013.34044http://www.scirp.org/journal/ijaa
29. Martín, A. *Simulador de vuelo. Viaje a Saturno.* Proyecto de Fin de Carrera. Universidad Politécnica de Madrid, 2015.
30. Álvarez, A. et al. *Mathematical model on rotational dynamics.*First International congress on Maths, I. and A. UP Madrid. ETSICCP: 2007.
31. Barceló, G. (2012). Analysis of Dynamics Fields in No inertial Systems. *World Journal of Mechanics, 2.175-180.*http://dx.doi.org/10.4236/wjm.2012.23021http://www.SciRP.org/journal/wjm
32. Pérez, L. A. (2013). *New evidences of Rotational Dynamics*. World Journal of Mechanics, *1* (3), 174-177. http://dx.doi.org/10.4236/wjm.2013.33016http://www.scirp.org/journal/wjm
33. Fernández- Rañada, A. (1993). *Física Básica.* Madrid.
34. Chandler, S.: *On the variation of latitude*. Astronomical Journal 11(248):59-86.1891, and: *On*

© 2016 Global Journals Inc. (US)

the supposed secular variation of latitudes. Astronomical Journal, 11(254):107-9. 1892.
35. Barceló, G.: *Technological Applications of the New Theory of Dynamic Interactions*. Global Journal of Researches in Engineering: Mechanical and Mechanics Engineering-G, Volume 13, Issue 5, 2013. https://globaljournals.org/GJRE_Volume13/E-Journal_GJRE_(G)_Vol_13_Issue_5.pdf.

© 2016 Global Journals Inc. (US)

This page is intentionally left blank

© 2016 Global Journals Inc. (US)

Global Journal of Science Frontier Research: a
Physics and Space Science
Volume 16 Issue 3 Version 1.0 Year 2016
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Analytical Determination of Subsurface Temperature using Two Layers Model in Part of Niger Delta Sedimentary Basin, Nigeria

By Emujakporue, Godwin Omokenu
University of Port Harcourt, Nigeria

Abstract- The solution of Fourier's one-dimensional steady state conduction heat flow equation for two layers-model has been solved using the analytical techniques and applied to part of Niger Delta sedimentary basin. The temperature-depth profile obtained shows a linear increase in temperature with depth. The internal properties affecting the thermal structures are the thermal conductivity and heat production. Comparison of measured and computed temperatures for some wells showed good agreement with minor variation ranging between -4.0 to 5.0 oC. This model can be used to determine thermal properties of other parts of the basin with little adjustment of the rock properties values.

Keywords: *analytical modelling, two-layer model, geotherm, bottom hole temperature, heat flow.*

GJSFR-A Classification : *FOR Code: 029999*



Strictly as per the compliance and regulations of :

© 2016. Emujakporue, Godwin Omokenu. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Analytical Determination of Subsurface Temperature using Two Layers Model in Part of Niger Delta Sedimentary Basin, Nigeria

Emujakporue, Godwin Omokenu

Abstract- **The solution of Fourier's one-dimensional steady state conduction heat flow equation for two layers-model has been solved using the analytical techniques and applied to part of Niger Delta sedimentary basin. The temperature-depth profile obtained shows a linear increase in temperature with depth. The internal properties affecting the thermal structures are the thermal conductivity and heat production. Comparison of measured and computed temperatures for some wells showed good agreement with minor variation ranging between -4.0 to 5.0 oC. This model can be used to determine thermal properties of other parts of the basin with little adjustment of the rock properties values.**

Keywords: analytical modelling, two-layer model, geotherm, bottom hole temperature, heat flow.

I. Introduction

Geothermal heat flow is a natural process by which heat is transferred from the earth interior to the surface. Heat can be transferred by three different ways, namely conduction, convection, and radiation (Cacace et al., 2010; Beardemore and Cull, 2001).Conduction is the major process by which heat is transfer through the solid part of the earth (crust) while in the fluid part (mantle) heat is usually transfer by convectional method. Geothermal heat flow variation is dependent on subsurface temperature variation and distribution, rock thermal conductivity and depths to heat sources.

Some researchers have worked on the geothermal gradient and heat flow of different parts of the Niger Delta sedimentary basin using bottom-hole or continuous temperatures. Two of the researchers (Odumodu and Ayonma, 2014) calculated the geothermal gradient and heat flow in parts of the eastern Niger delta from bottom hole temperature in 71 wells and obtained geothermal gradient varying between 12 to 24°C/Km with an average of 17.6°C/Km in the coastal swamp, 14°C /Km to 26°C/Km with an average of 20.4 °C/Km in the shallow offshore. They obtained average heat flow value of 42.5mWm^{-2}. Another group (Chukwueke et al., 1992) discovered that the geothermal gradients and heat flow values in the distal part of the Niger delta varies between 19-32°C/Km and 45-85mWm^{-2} respectively.

Author: Department of Physics, University of Port Harcourt, Port Harcourt, Nigeria. e-mail: owin2009@yahoo.com

The major source of information about subsurface temperature is from continuous or bottom hole temperatures measured in oil wells(Rider, 2002). Unfortunately, as a result of high cost and limited depth of drilling, current drilling campaigns in the Niger delta are limited to depths of only a few kilometres and at certain places, rendering direct and complete characterization of the entire subsurface thermal regime impossible to be carried out. Therefore, it is imperative to predict subsurface thermal conditions using alternative means. One method, is by solving Fourier's 1-D steady- state conduction heat flow equation for various layer models and applying it to a given basin.

A steady-state heat flow occurs when the amount of heat arriving and leaving a column of substance is equal over long period of time. Heat conduction across unit area, is a function of thermal property of the material and thermal conductivity. The ability of a substance to transfer its internal heat from one point to another depends on its thermal conductivity(Turcotte and Schubber, 2002; Safanda, 1985; Rudnick et al., 1998). The basin model is based on an extensional sedimentary basin where heat transfer by conduction dominates and the effect of advective transfer of heat by circulating groundwater is negligible due to flat surface topography. The spatial variation of temperature of a sedimentary basin is dependent on thermal conductivity, boundary conditions, water flow, rate of sedimentation and erosion and radiogenicheat production.

The objective of this research is to perform analytical modelling of 1-D conductive steady state temperature distribution of part of the Niger Delta by using two layer techniques and to characterize the geothermal features of the subsurface. The analytical solution of the 1-D Fourier heat flow for two layers model will be used to compute the subsurface temperature of the study area which will be compared with measured temperature from some oil wells.

II. Summary of the Geology of the Niger Delta

Figure 1 shows the location of the study area within the Niger Delta sedimentary basin. The Niger Delta is the youngest sedimentary basin in the Benue

© 2016 Global Journals Inc. (US)

Trough system and it started developing after the Eocene tectonic phase (Ekweozor and Daukoru, 1994; Doust and Omatsola, 1990). The Niger and Benue Rivers is the main supplier of sediments. The thickness of the sediment is approximately 12.0 Km.




Fig. 1 : Map of Niger Delta Showing the Study Area

The Tertiary portion of Niger Delta is divided into three lithostratigraphy (Figure 2.) representing prograding depositional facies that are distinguished based on sand – shale ratios. The lithostratigraphyare Benin, Agbada and Akata Formations(Ekweozor and Omatsola, 1994; Kulke, 1995). The Benin Formation is the youngest of the Delta sequence and it consists mainly of sand and gravels with thickness ranging from 0 - 2100m. The sands and sandstones in this Formation are coarse – fine and commonly granular in texture and partly unconsolidated. Very little oil has been found in the Benin formation, and it is the major source of portable water in the area.

© 2016 Global Journals Inc. (US)




Fig. 2 : Stratigraphic setting of Niger Delta showing the three formations of Niger Delta (modified after [1])

The Agbada Formation consists of alternation of sandstones and shale. The deposition of Agbada Formation began in Eocene and continues into Pleistocene. Shale and sandstone beds were deposited in equal proportions in the lower portion. The upper portion consists of minor shale interbed. The thickness of the Agbada Formation ranges from 300m – 4500m. Most reservoir in the Niger Delta are found in the Agbada Formation.

The Akata Formation forms the basal part of the Delta complex. It is of marine origin and it is made up of thick shale sequence (potential source rock), turbidite sand (potential reservoir in deep water) and minor amount of clay and silt(Avbovbo, 1978). Beginning in the Paleocene and through the recent, the Akata Formation was formed when terrestrial organic matter and clays were transported to deep water areas characterized by low energy conditions and oxygen deficiency. The thickness of the Akata Formation ranges from 0 – 6000m. The formation out lies the entire Delta, and is typically over pressured. It is the major source rock in the basin.

III. Materials and Methods

In order to characterize the process of heat transfer in the subsurface, it is necessary to understand the underlying physical principle. The general conduction equation governing heat transfer in the Earth is the Fourier heat equation (Mussett and Khand, 2000; Lowrie, 2007). The one-dimensional steady state heat conduction equation in vertical direction in a sedimentary column for a two layers model(Fowler, 2005) is given as;

$$\frac{d2T}{dz2} = -\frac{A1}{K} \quad (1)$$

And

$$\frac{d2T}{dz2} = -\frac{A2}{K} \quad (2)$$

Where

T = temperature (°C)

$A\ (1)$ = radioactive heat production for layer 1

$A\ (2)$ = radioactive heat production for layer 2

K = thermal conductivity (Wm^{-1} °C)

z = depth (m).

The radioactive heat production is as expressed as source (mWm^{-3}). The assumed two layers are the Benin and Agbada Formations of Niger Delta sedimentary basin. Assuming the heat production in the Benin and Agbada Formations are A1 and A2

© 2016 Global Journals Inc. (US)

respectively and the boundary between the two lithostratigraphy is Z1, then the ordinary differential equations 1 and 2 can be solved by applying the following assumptions and boundary conditions:

Assumptions

$$A = A1 \text{ for } 0 \leq z < z1 \quad \text{(Benin Formation)} \tag{3}$$

$$A = A2 \text{ for } z1 \leq z < z2 \text{ (Agbada Formation)} \tag{4}$$

Boundary Conditions

$$T = 27^{o}C \text{ on } z = 0 \tag{5}$$

$$Q = -Q2 \text{ on } z = z2. \tag{6}$$

Where

Q2 = heat flow from the base of the Agbada Formation.

The average surface temperature in the Niger Delta is 27°C. The basal heat flow Q=-Q2 is negative because heat flow upwards out of the earth, which is in the negative z direction. It is straightforward to solve the second order temperature differential equations (1 and 2) with two integrations. Integrating the differential equations two times results in two integration constants. The constants can be obtained by applying the two boundary conditions(Stein, 1995; Gibson, 2008). Integrating equations (1 and 2) once, we obtained

$$\frac{dT}{dz} = -\frac{A1}{K}z + C1 \tag{7}$$

And

$$\frac{dT}{dz} = -\frac{A2}{K}z + c2 \tag{8}$$

Integrating equations 7 and 8, then

$$T1 = -\frac{A1Z2}{K} + C1Z + C3 \tag{9}$$

And

$$T2 = -\frac{A2Z2}{K} + C2Z + C4 \tag{10}$$

Where T1 and T2 are the temperature regimes of Benin and Agbada Formations respectively. Applying the first boundary condition (equation 5) to equation 9, at the earth surface, then

$$C3 = 27 \tag{11}$$

Equation 7 is equal to equation 8 at the boundary between the two layers (Z1), therefore

$$-\frac{A1}{K}z1 + C1 = -\frac{A2}{K}z1 + c2 \tag{12}$$

Solving for C1 in terms of C2, then

$$C1 = -\frac{A2}{K}z1 + \frac{A1}{K}z1 + c2 \tag{13}$$

Putting equation 13 into equation 9, we have

$$T1 = -\frac{A1Z2}{K} + \left(-\frac{A2}{K}z1 + \frac{A1}{K}z1 + c2\right)Z + C3 \tag{14}$$

Substituting equation 11 into equation 14, we obtained

$$T1 = -\frac{A1Z2}{K} + \left(-\frac{A2}{K}z1 + \frac{A1}{K}z1 + c2\right)Z + 27 \tag{15}$$

At the base of the Agbada Formation (Z= Z2), applying second boundary condition (equation 5);

$$\frac{dT}{dZ} = \frac{Q2}{K} \tag{16}$$

Comparing equations 8 and 16, then

$$\frac{Q2}{K} = -\frac{A2}{K}z2 + c2 \tag{17}$$

Solving for C2 in equation 17, we obtained

$$c2 = \frac{A2}{K}z2 + \frac{Q2}{K} \tag{18}$$

Substituting equation 18 for C2 in equation 15, we obtained

$$T1 = -\frac{A1Z2}{K} + \left(-\frac{A2}{K}z1 + \frac{A1}{K}z1 + \frac{A2}{K}z2 + \frac{Q2}{K}\right)Z + 27 \tag{19}$$

Similarly, putting equation 18 into 10, we obtained

$$T2 = -\frac{A2Z2}{K} + \left(\frac{A2}{K}z2 + \frac{Q2}{K}\right)Z + C4 \tag{20}$$

At z1, equation 19 is equal to equation 20, therefore

$$-\frac{A2Z2}{K} + \left(\frac{A2}{K}z2 + \frac{Q2}{K}\right)Z1 + C4 = \frac{A1Z2}{K} + \left(-\frac{A2}{K}z1 + \frac{A1}{K}z1 + \frac{A2}{K}z2 + \frac{Q2}{K}\right)Z1 + 27 \tag{21}$$

Solving for C4, we have

$$C4 = \frac{A1Z2}{K} + \frac{A2Z2}{K} + \left(-\frac{A2}{K}z1 + \frac{A1}{K}z1\right)Z1 + 27 \tag{22}$$

Putting equation 22 into equation 20, we have

$$T2 = -\frac{A2Z2}{K} + \left(\frac{A2}{K}z2 + \frac{Q2}{K}\right)Z + \frac{A1Z2}{K} - \frac{A2Z2}{K} + \left(-\frac{A2}{K}z1 + \frac{A1}{K}z1\right)Z1 + 27 \tag{23}$$

© 2016 Global Journals Inc. (US)

The solutions 19 and 23 give the temperature at any position in the depth interval 0 to z1 and z1 to z2.

IV. Result and Discussion

The equilibrium geotherm for the Benin and Agbada Formations was calculated by considering each layer separately. Figure 3depictsa 1-D model for the two layers with its associated boundary conditions (surface temperature and basal heat flow). Thermal conductivity and heat production are the internal properties that influence the temperature structure in the column. The thermal conductivity and heat production were obtained from well logs (Emujakporue, 2009; Emujakporue, 2016).

The model is made up of sediment column with two layers of thickness 1.8 and 4.5 Km respectively. The thermal boundary conditions are surface temperature To = 27°C and basal heat flow of 50.0mWm^{-2} and thermal conductivity. Temperature and temperature gradient were matched across the boundary between the two layers.

Using the above model properties the equilibrium geotherm was calculated for the two lithostratigraphy from equations 21 and 23. The equilibrium geotherms was calculated by assuming heat production and thermal conductivity for layer 1 to be 1.8*10^{-6} and 2.0 Wm^{-1} ∘C^{-1}respectively. Similarly, for the second layer, basal heat flow, heat production and thermal conductivity were 50 mWm^{-2}, 2.1*10^{-6}, and 2.2 Wm^{-1}°C^{-1}.




Fig. 3 : A two-layer model for the Benin and Agbada Formations

The computed temperature depth distribution is shown in Figure 4. The geotherm increases linearly with depth which is normal. The linear relationship between computed temperature and depth from the graph is given as;

$$T = 22.407x + 27.0 \quad (24)$$

The regression coefficient for the equation is 0.9992. The high regression coefficient is an indication of the significant and confidence imposed on the relationship. The equilibrium geotherm for the model rock column changes when the conductivity, radioactive heat generation and basal heat flow are varied.

Equation 24 was applied to part of the Niger Delta and compare with measured bottom hole temperature from some oil wells in the Niger Delta and the result is shown in Table 1.The difference between measured and computed temperature ranges between -4.0 to 5. 0°C.The difference between the measured and computed values are minimum and within tolerance level. The computed average geotherm provides an interesting useful information on the thermal state of the sedimentary basin.

As a result of lateral variation of heat flow due to changes in lithology, thermal conductivity, heat production and groundwater flow, it is necessary to carry out accurate determination of the above parameters in order to use the model equations. The results of the geothermal modelling can be used as an aid to interpreting heat flow patterns in the study area, thermal and hydrocarbon modelling. The surface heat flow can be calculated by multiplying the temperature gradient (slope of the graph) with the thermal conductivity values at a particular depth. Therefore, accurate geothermal heat flow is dependent on accurate determination of subsurface lithology distribution and their corresponding thermal conductivity values.

© 2016 Global Journals Inc. (US)




Fig. 4 : Computed equilibrium geotherm of the studied area

Table 1 : Measured and computed temperatures at some depths for three wells

Well	Depth (Km)	Measured Temp(oC)	Comp. Temp (Oc)	Difference
	3.218	98.89	102.323	-3.43373
	3.21716	98.89	102.304	-3.41406
Well1	3.21564	98.89	102.268	-3.37849
	3.80634	117.22	116.095	1.125
	3.80665	117.22	116.102	1.11774
	3.80695	116.67	116.109	0.56072
	3.80875	117.22	116.151	1.06858
well2	3.039	97.78	98.1338	-0.35387
	3.051	97.78	98.4147	-0.63476
	3.206	107.22	102.042	5.17715
	3.3395	107.22	105.167	2.05232
	3.205	107.22	102.019	5.20056
	3.38	111.11	106.115	4.99434
	3.394	111.11	106.443	4.66664
	3.39	111.11	106.349	4.76027
well3	2.9852	92.78	96.8745	-4.09458
	2.981	92.78	96.7762	-3.99627
	2.985	92.78	96.8699	-4.0899
	2.98	92.78	96.7528	-3.97286
	3.041	101.67	98.1806	3.48931
	3.044	101.67	98.2509	3.41909
	3.176	101.67	101.340	0.32936
	3.1766	101.67	101.354	0.31532

V. Conclusion

In this study the analytical solution of the 1-D steady state heat conduction equation based on two layers model was obtained and applied to the study area. Two equations obtained for the two layers were used to generate the temperature structure. The accuracy of the model depends on the accuracy of thermal conductivity, heat production in the sediment and the basal heat flow. Comparison of the computed and measured temperature shows that the difference is minimal and the model is dependable and can be applied for thermal and hydrocarbon modelling in the Niger Delta. Therefore, it is possible to predict subsurface thermal properties without drilling expensive hole and also carrying out temperature logging.

© 2016 Global Journals Inc. (US)

References Références Referencias

1. Avbovbo, A. A. 1978. Tertiary lithostratigraphy of Niger Delta. AAPG Bull. 62; 695- 306.
2. Beardsmore, G. R. and J. P. Cull (2001). Crustal Heat Flow: A Guide to Measurement and Modelling. New York, NY: Cambridge University Press.
3. Cacace, M., B.O. Kaiser, B. Lewerenz, M. Scheck-Wenderoth, (2010). "Geothermal energy in sedimentary basins: what we can learn from numerical models," Chemie der Erde vol. 70 (3), 33-46.
4. Chukwueke, C, G. Thomas, and J. Delfraud (1992). "Sedimentary processes, eustatism, subsidence and heat flow in the distal Part of the Niger Delta". Bull. Centres Rech. Explor.-Prod. Elf-Acquitaine. 16(1):137 – 186.
5. Doust, H and E. Omatsola (1990). Niger Delta in, Divergent/ Passive Margin Basin, AAPG Memoi 48; Tulsa, American Association Petroleum Geologist; 239-248.
6. Ekweozor, C. M. and E. M. Daukoru (1994). Northern delta depobelt portion of the Akata – Abgada (1) petroleum system, Niger Delta, Nigeria, in Magoon L B, Dow WG.eds. The petroleum system from source to Trap, AAPG. Memoir 60; 599 – 614.
7. Emujakporue, G. O. (2009). Subsidence and geothermal history in the eastern Niger delta with implications for hydrocarbons. Unpublished PhD. Thesis U.P.H.
8. Emujakporue, G. O. (2016). Radiogenic heat production from well logs in part of Niger Delta sedimentary basin, Nigeria. Global Journal of Science Frontier Research: Physics and Space Science Vol.16 (1), 57-64
9. Fowler, C. M. R. (2005) The Solid Earth: an introduction to global geophysics, 2nd edn. Cambridge University Press, Cambridge,
10. Gibson, H., K. Stuwe, R. Seikel, D. FitzGerald, P. Calcagno, A. Guillen, P. McInerney, (2008). "Forward prediction of temperature distribution direct from 3D geology models", Australian Geothermal Energy Conference.
11. Kulke, H. (1995). "Nigeria", in Kulke H.,ed, Regional Petroleum Geology of the World, part 11, Gebruder, Bombraeger, Berlin, Africa, America, Australia and Antarctica; 143-172
12. Lowrie, W. (2007). Fundamentals of Geophysics. Cambridge: Cambridge University Press, New York. 1st, edition.
13. Mussett. A. E. and M. A. Khan (2000).*Looking into the Earth: An Introduction to Geological Geophysics*, Cambridge: Cambridge University Press.
14. Odumodu, C. F. R. and W. M. Ayonma (2014). Present Day Geothermal Regime in Parts of the Eastern Niger Delta, Nigeria. Petroleum Technology Development Journal (ISSN 1595-9104): An International Journal; Vol. 1 (1); 7-26.
15. Rider, M. (2002). The geological interpretation of well logs. Rider-French consulting, Sotland. 2nd edition.
16. Rudnick, R. L, W. F. McDonough and R. J. O'Connell (1998). Thermal structure, thickness and composition of continental lithosphere. *Chem. Geol.* 145; 395–411.
17. Safanda, J. (1985). Calculation of temperature distribution in two-dimensional geothermal profile," Studiageoph. et geod. 29; 197-207.
18. Stein. C. A. (1995) Heat flow of the Earth. In Global Earth Physics: A Handbook of Physical Constants. AGU Reference Shelf 1. Washington: American Geophysical Union, 144–58.
19. Turcotte, D. L. and G. Schubert, (2002). *Geodynamics*, 2nd edn. Cambridge: Cambridge University Press.

© 2016 Global Journals Inc. (US)

Global Journals Inc. (US) Guidelines Handbook 2016

www.GlobalJournals.org

FELLOWS

FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (FARSS)

Global Journals Incorporate (USA) is accredited by Open Association of Research Society (OARS), U.S.A and in turn, awards "FARSS" title to individuals. The 'FARSS' title is accorded to a selected professional after the approval of the Editor-in-Chief/Editorial Board Members/Dean.



- The "FARSS" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSS or William Walldroff, M.S., FARSS.

FARSS accrediting is an honor. It authenticates your research activities. After recognition as FARSB, you can add 'FARSS' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, and Visiting Card etc.

The following benefits can be availed by you only for next three years from the date of certification:

FARSS designated members are entitled to avail a 40% discount while publishing their research papers (of a single author) with Global Journals Incorporation (USA), if the same is accepted by Editorial Board/Peer Reviewers. If you are a main author or co-author in case of multiple authors, you will be entitled to avail discount of 10%.

Once FARSB title is accorded, the Fellow is authorized to organize a symposium/seminar/conference on behalf of Global Journal Incorporation (USA). The Fellow can also participate in conference/seminar/symposium organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent.





You may join as member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer. In addition, it is also desirable that you should organize seminar/symposium/conference at least once.

We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



© Copyright by Global Journals Inc.(US) | Guidelines Handbook

The FARSS can go through standards of OARS. You can also play vital role if you have any suggestions so that proper amendment can take place to improve the same for the benefit of entire research community.



As FARSS, you will be given a renowned, secure and free professional email address with 100 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.



The FARSS will be eligible for a free application of standardization of their researches. Standardization of research will be subject to acceptability within stipulated norms as the next step after publishing in a journal. We shall depute a team of specialized research professionals who will render their services for elevating your researches to next higher level, which is worldwide open standardization.



The FARSS member can apply for grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.
Once you are designated as FARSS, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria. After certification of all your credentials by OARS, they will be published on your Fellow Profile link on website https://associationofresearch.org which will be helpful to upgrade the dignity.

The FARSS members can avail the benefits of free research podcasting in Global Research Radio with their research documents. After publishing the work, (including published elsewhere worldwide with proper authorization) you can upload your research paper with your recorded voice or you can utilize chargeable services of our professional RJs to record your paper in their voice on request.



The FARSS member also entitled to get the benefits of free research podcasting of their research documents through video clips. We can also streamline your conference videos and display your slides/ online slides and online research video clips at reasonable charges, on request.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook



The FARSS is eligible to earn from sales proceeds of his/her researches/reference/review Books or literature, while publishing with Global Journals. The FARSS can decide whether he/she would like to publish his/her research in a closed manner. In this case, whenever readers purchase that individual research paper for reading, maximum 60% of its profit earned as royalty by Global Journals, will be credited to his/her bank account. The entire entitled amount will be credited to his/her bank account exceeding limit of minimum fixed balance. There is no minimum time limit for collection. The FARSS member can decide its price and we can help in making the right decision.

The FARSS member is eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get remuneration of 15% of author fees, taken from the author of a respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account.



MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (MARSS)

The ' MARSS ' title is accorded to a selected professional after the approval of the Editor-in-Chief / Editorial Board Members/Dean.



The "MARSS" is a dignified ornament which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., MARSS or William Walldroff, M.S., MARSS.

MARSS accrediting is an honor. It authenticates your research activities. After becoming MARSS, you can add 'MARSS' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, Visiting Card and Name Plate etc.

The following benefitscan be availed by you only for next three years from the date of certification.



MARSS designated members are entitled to avail a 25% discount while publishing their research papers (of a single author) in Global Journals Inc., if the same is accepted by our Editorial Board and Peer Reviewers. If you are a main author or co-author of a group of authors, you will get discount of 10%.

As MARSS, you will be given a renowned, secure and free professional email address with 30 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.



© Copyright by Global Journals Inc.(US) | Guidelines Handbook



We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

The MARSS member can apply for approval, grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.





Once you are designated as MARSS, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria.

It is mandatory to read all terms and conditions carefully.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

AUXILIARY MEMBERSHIPS

Institutional Fellow of Global Journals Incorporation (USA)-OARS (USA)

Global Journals Incorporation (USA) is accredited by Open Association of Research Society, U.S.A (OARS) and in turn, affiliates research institutions as "Institutional Fellow of Open Association of Research Society" (IFOARS).



The "FARSC" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSC or William Walldroff, M.S., FARSC.

The IFOARS institution is entitled to form a Board comprised of one Chairperson and three to five board members preferably from different streams. The Board will be recognized as "Institutional Board of Open Association of Research Society"-(IBOARS).

The Institute will be entitled to following benefits:

The IBOARS can initially review research papers of their institute and recommend them to publish with respective journal of Global Journals. It can also review the papers of other institutions after obtaining our consent. The second review will be done by peer reviewer of Global Journals Incorporation (USA) The Board is at liberty to appoint a peer reviewer with the approval of chairperson after consulting us.

The author fees of such paper may be waived off up to 40%.

The Global Journals Incorporation (USA) at its discretion can also refer double blind peer reviewed paper at their end to the board for the verification and to get recommendation for final stage of acceptance of publication.





The IBOARS can organize symposium/seminar/conference in their country on behalf of Global Journals Incorporation (USA)-OARS (USA). The terms and conditions can be discussed separately.

The Board can also play vital role by exploring and giving valuable suggestions regarding the Standards of "Open Association of Research Society, U.S.A (OARS)" so that proper amendment can take place for the benefit of entire research community. We shall provide details of particular standard only on receipt of request from the Board.



The board members can also join us as Individual Fellow with 40% discount on total fees applicable to Individual Fellow. They will be entitled to avail all the benefits as declared. Please visit Individual Fellow-sub menu of GlobalJournals.org to have more relevant details.

© Copyright by Global Journals Inc.(US) | Guidelines Handbook

We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



After nomination of your institution as "Institutional Fellow" and constantly functioning successfully for one year, we can consider giving recognition to your institute to function as Regional/Zonal office on our behalf.
The board can also take up the additional allied activities for betterment after our consultation.

The following entitlements are applicable to individual Fellows:



Open Association of Research Society, U.S.A (OARS) By-laws states that an individual Fellow may use the designations as applicable, or the corresponding initials. The Credentials of individual Fellow and Associate designations signify that the individual has gained knowledge of the fundamental concepts. One is magnanimous and proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice.

Open Association of Research Society (US)/ Global Journals Incorporation (USA), as described in Corporate Statements, are educational, research publishing and professional membership organizations. Achieving our individual Fellow or Associate status is based mainly on meeting stated educational research requirements.



Disbursement of 40% Royalty earned through Global Journals : Researcher = 50%, Peer Reviewer = 37.50%, Institution = 12.50% E.g. Out of 40%, the 20% benefit should be passed on to researcher, 15 % benefit towards remuneration should be given to a reviewer and remaining 5% is to be retained by the institution.

We shall provide print version of 12 issues of any three journals [as per your requirement] out of our 38 journals worth $ 2376 USD.

Other:

The individual Fellow and Associate designations accredited by Open Association of Research Society (US) credentials signify guarantees following achievements:

- The professional accredited with Fellow honor, is entitled to various benefits viz. name, fame, honor, regular flow of income, secured bright future, social status etc.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

- In addition to above, if one is single author, then entitled to 40% discount on publishing research paper and can get 10%discount if one is co-author or main author among group of authors.
- The Fellow can organize symposium/seminar/conference on behalf of Global Journals Incorporation (USA) and he/she can also attend the same organized by other institutes on behalf of Global Journals.
- The Fellow can become member of Editorial Board Member after completing 3yrs.
- The Fellow can earn 60% of sales proceeds from the sale of reference/review books/literature/publishing of research paper.
- Fellow can also join as paid peer reviewer and earn 15% remuneration of author charges and can also get an opportunity to join as member of the Editorial Board of Global Journals Incorporation (USA)
- • This individual has learned the basic methods of applying those concepts and techniques to common challenging situations. This individual has further demonstrated an in–depth understanding of the application of suitable techniques to a particular area of research practice.

Note :

"

- In future, if the board feels the necessity to change any board member, the same can be done with the consent of the chairperson along with anyone board member without our approval.
- In case, the chairperson needs to be replaced then consent of 2/3rd board members are required and they are also required to jointly pass the resolution copy of which should be sent to us. In such case, it will be compulsory to obtain our approval before replacement.
- In case of "Difference of Opinion [if any]" among the Board members, our decision will be final and binding to everyone.

"

Process of submission of Research Paper

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (*.DOC,*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission.Online Submission: There are three ways to submit your paper:

(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.

(II) Choose corresponding Journal.

(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.

(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.

(C) If these two are not conveninet , and then email the paper directly to dean@globaljournals.org.

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Preferred Author Guidelines

MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11'"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

You can use your own standard format also.
Author Guidelines:

1. General,

2. Ethical Guidelines,

3. Submission of Manuscripts,

4. Manuscript's Category,

5. Structure and Format of Manuscript,

6. After Acceptance.

1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global

Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.

2) Drafting the paper and revising it critically regarding important academic content.

3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.

Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.



To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

5.STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also.Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

Papers: These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

(a)Title should be relevant and commensurate with the theme of the paper.

(b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.

(c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.

(d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.

(e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.

(f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;

(g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.

(h) Brief Acknowledgements.

(i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.

The Editorial Board reserves the right to make literary corrections and to make suggestions to improve briefness.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

Format

Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than 1.4 × 10-3 m3, or 4 mm somewhat than 4 × 10-3 m. Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

Abstract, used in Original Papers and Reviews:

Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art.A few tips for deciding as strategically as possible about keyword search:

- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

Acknowledgements: Please make these as concise as possible.

References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

Tables, Figures and Figure Legends

Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.

Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.

Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.

Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.

6. AFTER ACCEPTANCE

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

6.1 Proof Corrections

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

www.adobe.com/products/acrobat/readstep2.html. This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at dean@globaljournals.org within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

6.3 Author Services

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

6.4 Author Material Archive Policy

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

6.5 Offprint and Extra Copies

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: editor@globaljournals.org .



Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

1. Choosing the topic: In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

2. Evaluators are human: First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

3. Think Like Evaluators: If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

4. Make blueprints of paper: The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

5. Ask your Guides: If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

6. Use of computer is recommended: As you are doing research in the field of Computer Science, then this point is quite obvious.

7. Use right software: Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

8. Use the Internet for help: An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

9. Use and get big pictures: Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

10. Bookmarks are useful: When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

11. Revise what you wrote: When you write anything, always read it, summarize it and then finalize it.

12. Make all efforts: Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

13. Have backups: When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

14. Produce good diagrams of your own: Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

15. Use of direct quotes: When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

16. Use proper verb tense: Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

17. Never use online paper: If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

18. Pick a good study spot: To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

19. Know what you know: Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

20. Use good quality grammar: Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

21. Arrangement of information: Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

22. Never start in last minute: Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

23. Multitasking in research is not good: Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

24. Never copy others' work: Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

25. Take proper rest and food: No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

26. Go for seminars: Attend seminars if the topic is relevant to your research area. Utilize all your resources.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

27. Refresh your mind after intervals: Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

28. Make colleagues: Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

29. Think technically: Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

30. Think and then print: When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

31. Adding unnecessary information: Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

32. Never oversimplify everything: To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

33. Report concluded results: Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

34. After conclusion: Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium though which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

Informal Guidelines of Research Paper Writing

Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.

Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

General style:

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

· Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

· Use standard writing style including articles ("a", "the," etc.)

· Keep on paying attention on the research topic of the paper

· Use paragraphs to split each significant point (excluding for the abstract)

· Align the primary line of each section

· Present your points in sound order

· Use present tense to report well accepted

· Use past tense to describe specific results

· Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives

· Shun use of extra pictures - include only those figures essential to presenting results

Title Page:

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript--must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for briefness. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

Approach:

- Single section, and succinct
- As a outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an conceptual must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.

- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

Procedures (Methods and Materials):

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

Methods:

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

Approach:

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

What to keep away from

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

Results:

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently.You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

Segment Draft and Final Research Paper: You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptive of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.

- Do not give permission to anyone else to "PROOFREAD" your manuscript.

- Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Criterion for Grading a Research Paper (Compilation) by Global Journals Inc. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

Topics	Grades		
	A-B	**C-D**	**E-F**
Abstract	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form Above 200 words	No specific data with ambiguous information Above 250 words
Introduction	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
Methods and Procedures	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
Result	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
Discussion	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
References	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring

Index

G

Gyrokinetics · 1, 2, 3, 10

H

Heuristic · 3
Hitherto · 17

I

Instantaneous · 17

L

Levitate · 26
Lorentz · 1, 8, 9

P

Paucity · 28
Pleistocene · 34
Propellant · 23

T

Toroidal · 1, 2, 3, 4, 5, 6, 7, 8, 9
Torque · 3, 4, 5, 8, 9, 17, 18, 19, 21, 22, 23, 28, 29



Global Journal of Science Frontier Research

Visit us on the Web at www.GlobalJournals.org | www.JournalofScience.org
or email us at helpdesk@globaljournals.org

ISSN 9755896



© Global Journals