



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH
MATHEMATICS AND DECISION SCIENCES

Volume 13 Issue 6 Version 1.0 Year 2013

Type : Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Characterization of Partial Lattices on Countable Boolean Lattice

By D.V.S.R. Anil Kumar, Y.V.Seshagiri Rao, Y Narasimhulu &
Venkata Sundaranand Putcha

Nagarjuna University, India

Abstract - In this paper new concepts countable join property, countable meet property, P_σ – lattice and P_δ – lattice are introduced. We established that P_σ – lattice and P_δ – lattice are measurable partial lattices and characterized partial lattices of a lattice through countable join and meet properties. We also established some interesting result on the injective property of the lattice measurable functions defined over countable Boolean lattices.

Keywords : lattice, partial lattice, σ - algebra, measure.


GJSFR-F Classification : ASM 03G10, 28A05



Strictly as per the compliance and regulations of :



© 2013. D.V.S.R. Anil Kumar, Y.V.Seshagiri Rao, Y Narasimhulu & Venkata Sundaranand Putcha. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License (<http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.



Characterization of Partial Lattices on Countable Boolean Lattice

D.V.S.R. Anil Kumar ^α, Y.V.Seshagiri Rao ^σ, Y Narasimhulu ^ρ & Venkata Sundaranand Putcha ^ω

Abstract - In this paper new concepts countable join property, countable meet property, P_{σ} -lattice and P_{δ} -lattice are introduced. We established that P_{σ} -lattice and P_{δ} -lattice are measurable partial lattices and characterized partial lattices of a lattice through countable join and meet properties. We also established some interesting result on the injective property of the lattice measurable functions defined over countable Boolean lattices.

Keywords : lattice, partial lattice, σ -algebra, measure.

I. INTRODUCTION

The origin of a lattice concept can be traced back to Boole's analysis of thought and Dedekind's study of divisibility, Schroder and Pierce contributed substantially to this area. Though some of the work in this direction was done around 1930, much momentum was gained in 1967 with the contributions of Birkhoff's [2]. In 1963, Gabor szasz [9] introduced the generalization of the lattice measure concepts. To study σ -additive set functions on a lattice of sets, Gena A. DE Both [3] introduced σ -lattice in 1973. The concept of partial lattices was introduced by George Gratzer [5] in 1978. In 2000, Pao - Sheng Hus [8] characterized outer measures associated with lattice measure. The Hann decomposition theorem of a signed lattice measure by Jun Tanaka [10] defined a signed lattice measure on a lattice σ -algebras and the concept of sigma algebras are extensively studied by [4]. D.V.S.R. Anil Kumar et al [1] introduce the concept of measurable Borel lattices, σ -lattice and δ -lattice to characterize a class of Measurable Borel Lattices. This paper is organized as follows. Section 2 presents the preliminaries definitions and results. In Section 3 we proved that P_{σ} -lattice and P_{δ} -lattice are measurable partial lattices and all partial lattices of a lattice satisfy both countable join and meet properties. Some interesting result on the injective property of the lattice measurable functions defined over countable Boolean lattices are established in Section 4.

Author α : Nizam Institute of Engineering and Technology, Deshmukhi, Nalgonda district, Hyderabad, A.P., India.

E-mail : anilkumardaita@yahoo.in

Author σ : Teegala Krishna Reddy Engineering College, Meerpet, Hyderabad, A.P. India. E-mail : yangalav@gmail.com

Author ρ : Professor of Mathematics, University of Hyderabad. E-mail : yakkalan@uohyd.ernet.in

Author ω : Center for Mathematical Sciences-DST, CR Rao Advanced Institute of Mathematics, Statistics and Computer Science, University of Hyderabad Campus, Hyderabad, 500 046, India. E-mail : anand_putcha@yahoo.com

II. PRELIMINARIES

Consider a lattice $(L, \wedge, \vee)$ with the operations meet $\wedge$ and join $\vee$ and usual ordering $\leq$, where L is a collection of subset of a non empty set X . Now this lattice $(L, \wedge, \vee)$ is denoted by L and satisfy the commutative law, the associative law and the absorption law. A lattice L is called distributive if the distributive law is satisfied. The zero and one elements of the lattice L are denoted by 0 and 1 respectively. A distributive lattice L is called a Boolean lattice if for any element x in L , there exists a unique complement x^c such that $x \vee x^c = 1$ and $x \wedge x^c = 0$. An operator $C: L \rightarrow L$, where L is a lattice is called a lattice complement in L if the law of complementation, the law of contra positive and the law of double negation are satisfied. The following are very important examples of Boolean lattice.

Example2.1. Let $(\{0,1\}, \leq)$ be the set consisting of the two elements $0,1$ equipped with the usual order relation $0 \leq 1$. This poset is a Boolean lattice with respect to the operations presented in the tables below (at the left the lattice operations and at the right the complementation):

a	b	$a \wedge b$	$a \vee b$
0	0	0	0
0	1	0	1
1	0	0	1
1	1	1	1

x	x^c
0	1
1	0

This is usually known as the two valued or two elements Boolean lattice, denoted by $B = (\{0,1\}, \vee, \wedge, ^c, 0, 1)$.

Example2.2. The power set $P(X)$ of a universe X a Boolean lattice if we choose the set theoretic complement $A^c = X \setminus A := \{x \in X: x \in X \text{ and } x \notin A\}$ as the complement of a given set A in the universe X . Such a Boolean lattice is $P = (P(X), \vee, \wedge, ^c, \phi, X)$.

Example2.3. $E = (2^X, \vee, \wedge, ^c, 0, 1)$ is the collection 2^X of all two valued functional on the universe X is a Boolean lattice if we choose the functional $\chi^c = 1 - \chi$ as the complement of a given functional χ .

Example2.4. Let $(D, \vee, \wedge, ^c, 1, 70)$ is a Boolean lattice where $D = \{1, 2, 5, 7, 10, 14, 35, 70\}$ is the set of all divisors of 70 , $x \wedge y = \text{Greatest Common Divisor of } x \text{ and } y$, $x \vee y = \text{Least Common Multiple of } x \text{ and } y$ and $x^c = \frac{70}{x}$.

Definition2.1. A Boolean lattice L is called a countable Boolean lattice if L is closed under countable join and is denoted by $\sigma(L)$.

Example2.5. $\{\text{empty set } \phi, X\}$, Power set of X , Let $X = \mathfrak{R}$, $L = \{\text{measurable subsets of } \mathfrak{R}\}$ with usual ordering $(\leq)$ are all countable Boolean lattice.

Definition2.2. The entire set X together with countable Boolean lattice is called lattice measurable space and is denoted by the ordered pair $(X, \sigma(L))$.

Example2.6. $X = \mathfrak{R}$, where $\mathfrak{R}$ is extended real number system and $L = \{\text{All Lebesgue measurable sub sets of } \mathfrak{R}\}$, $(\mathfrak{R}, \sigma(L))$ is a lattice measurable space.

Definition2.3. If $\mu: \sigma(L) \rightarrow \mathbb{R} \cup \{\infty\}$ satisfies the following properties (i) $\mu(\phi) = \mu(0) = 0$ (ii) for all $h, g \in \sigma(L)$, such that $\mu(h), \mu(g) \geq 0$; $h \leq g \Rightarrow \mu(h) \leq \mu(g)$ (iii) for all $h, g \in \sigma(L)$: $\mu(h \vee g) + \mu(h \wedge g) = \mu(h) + \mu(g)$ (iv) If $h_n \in \sigma(L)$, $n \in \mathbb{N}$ such that $h_1 \leq h_2 \leq \dots \leq h_n \leq \dots$, then $\mu(\bigvee_{n=1}^{\infty} h_n) = \lim \mu(h_n)$ then μ is called a lattice measure on the countable Boolean lattice $\sigma(L)$.

The following is definition given in [5]

Definition2.4. Let $\sigma(L)$ be a countable Boolean lattice, $H \subseteq \sigma(L)$, and restrict $\wedge$ and $\vee$ to H as follows. For $a, b, c \in H$, if $a \wedge b = c$ (dually, $a \vee b = c$), then we say that in H , $a \wedge b$ (dually $a \vee b$) is defined and it equals c , if, for $a, b \in H$, $a \wedge b \notin H$ (dually $a \vee b \notin H$), then we say that $a \wedge b$ (dually $a \vee b$) is not defined in H . Thus $(H, \wedge, \vee)$ is a set with two binary partial operations. $(H, \wedge, \vee)$ is called a partial lattice, a relative sublattice of $\sigma(L)$.

Observation2.1. Every subset of a countable Boolean lattice determines a partial lattice. Every sublattice of $\sigma(L)$ is a partial lattice and the converse need not be true.

Definition2.5.[7] A set A is said to be measurable partial lattice, if A is in $\sigma(L)$.

Example2.7. $(\mathfrak{R}, \sigma(L))$ be lattice measurable space. Then the interval (a, ∞) is a measurable partial lattice under usual ordering.

Example2.8. $[0, 1) \subset \mathfrak{R}$ is a measurable partial lattice under usual ordering.

Definition2.9. A P_σ -lattice is a poset for which sup exist for any countable collection of its partial lattices.

Example2.9. $R = \bigvee_{n=1}^{\infty} (-n, n)$ is a p_σ -lattice.

Definition2.10. A P_δ -lattice is a poset for which inf exist for any countable collection of its partial lattices.

Example2.11. (i) $\bigwedge_{n=1}^{\infty} (-n, n) = (-1, 1)$ and (ii) $\bigwedge_{n=1}^{\infty} \left(-\frac{1}{n}, \frac{1}{n}\right) = \{0\}$ are P_δ -lattices.

Definition2.11.Countable join property (CJP): If $\{E_k\}$ is monotonic increasing sequence of partial lattices of a lattice L and $E = \bigvee_{k=1}^{\infty} E_k$. Then $\mu(E) = \lim_{n \rightarrow \infty} \mu(E_n)$.

Definition2.12.Countable meet property (CMP): If $\{E_k\}$ is a monotonic decreasing sequence of partial lattices of a lattice L and $E = \bigwedge_{k=1}^{\infty} E_k$. Then $\mu(E) = \lim_{n \rightarrow \infty} \mu(E_n)$.

Result2.1.[1]. If E is measurable lattice so is E^c .

III. P_σ -LATTICE AND P_δ -LATTICE

Theorem3.1. Every P_σ -lattice is lattice measurable.

Proof. Let $E_1, E_2, \dots$ are pair wise disjoint measurable partial lattices and $E = \bigvee_{k=1}^{\infty} E_k$,

Evidently,

$$\mu \left(\bigvee_{k=1}^{\infty} E_k \right) \leq \sum_{k=1}^{\infty} \mu(E_k) \quad (1)$$

and

$$\mu \left(\bigvee_{k=1}^{\infty} E_k \right) \geq \mu \left(\bigvee_{k=1}^n E_k \right) \quad (2)$$

From definition 2.3. We have $\mu(E_1 \vee E_2) = \mu(E_1) + \mu(E_2)$. By the principle of mathematical induction on number of pair wise disjoint measurable partial lattices, n , we have $\mu \left(\bigvee_{k=1}^n E_k \right) = \sum_{k=1}^n \mu(E_k)$. As $n \rightarrow \infty$, from (2) it follows that

$$\mu \left(\bigvee_{k=1}^{\infty} E_k \right) \geq \sum_{k=1}^{\infty} \mu(E_k) \quad (3)$$

From (1) and (3), we have $\mu \left(\bigvee_{k=1}^{\infty} E_k \right) = \sum_{k=1}^{\infty} \mu(E_k)$. Now $E = \bigvee_{k=1}^{\infty} E_k = E_1 \vee (E_2 \wedge E_1^c) \vee \dots \vee (E_k \wedge (\bigvee_{k=1}^{n-1} E_k^c) \vee \dots$. Since $E_1, E_2 \wedge E_1^c, \dots$ are disjoint measurable partial lattices, we have, $\bigvee_{k=1}^{\infty} E_k$ is a measurable partial lattice. Hence every P_{σ} -lattice is a lattice measurable.

Theorem 3.2. Every P_{σ} -lattice satisfies CJP.

Proof. Suppose that $\{E_k\}$ is monotonic increasing sequence of partial lattices of a $\sigma(L)$ and $E = \bigvee_{k=1}^{\infty} E_k$. Write $E = E_1 \vee (E_2 \wedge E_1^c) \vee \dots \vee (E_k \wedge (\bigvee_{k=1}^{n-1} E_k^c) \vee \dots$

So we have $E = E_1 \vee (\bigvee_{k=1}^{\infty} (E_{k+1} \wedge E_k^c))$ (a disjoint joint). By Theorem 3.1.

$$\begin{aligned} \text{Now, } \mu(E) &= \mu(E_1) + \sum_{k=1}^{\infty} \mu(E_{k+1} - E_k) = \mu(E_1) + \lim_{n \rightarrow \infty} \sum_{k=1}^n [\mu(E_{k+1}) - \mu(E_k)] = \mu(E_1) \\ &+ \lim_{n \rightarrow \infty} [\mu(E_2) - \mu(E_1) + \dots + \mu(E_n) - \mu(E_{n-1})] = \mu(E_1) + \lim_{n \rightarrow \infty} [-\mu(E_1) + \mu(E_n)] = \mu(E_1) - \mu(E_1) + \lim_{n \rightarrow \infty} \mu(E_n) = \lim_{n \rightarrow \infty} \mu(E_n). \end{aligned}$$

Theorem 3.3. Every P_{δ} -lattice is lattice measurable.

Proof. Let $E_1, E_2, \dots$ are measurable partial lattices.

By theorem 3.1. $E = \bigvee_{k=1}^{\infty} E_k$ is a measurable partial lattice. Let $G = \bigwedge_{k=1}^{\infty} E_k$.

Then $G^c = (\bigwedge_{k=1}^{\infty} E_k)^c = \bigvee_{k=1}^{\infty} E_k^c$. Given that each E_k is a measurable partial lattice.

Hence by Result 2.1., each E_k^c is a measurable partial lattice. Which implies $\bigvee_{k=1}^{\infty} E_k^c$ is a measurable partial lattice (Every P_{σ} -lattice is a measurable partial lattice). This leads to G^c is measurable partial lattice. Hence G is measurable partial lattice (By Result 2.1.).

Theorem 3.4. Every P_δ -lattice satisfies CMP.

Proof. Suppose that $\{E_k\}$ is a monotonic decreasing sequence of partial lattices of $\sigma(L)$ and $E = \bigwedge_{k=1}^{\infty} E_k$. Let $E = \bigwedge_{k=1}^{\infty} E_k$. Evidently $E_1 = E \vee (E_1 \wedge E_2^c) \vee (E_2 \wedge E_3^c) \vee \dots$

$$\begin{aligned} \text{Then } \mu(E_1) &= \mu(E) + \sum_{k=1}^{\infty} \mu(E_k) - \mu(E_{k+1}) = \mu(E) + \lim_{n \rightarrow \infty} \sum_{k=1}^n \mu(E_k) - \mu(E_{k+1}) \\ &= \mu(E) + \lim_{n \rightarrow \infty} [\mu(E_1) - \mu(E_2) + \dots + \mu(E_n) - \mu(E_{n+1})] = \mu(E) + \lim_{n \rightarrow \infty} [\mu(E_1) - \mu(E_{n+1})] \\ &= \mu(E) + \mu(E_1) - \lim_{n \rightarrow \infty} \mu(E_{n+1}). \text{ Which implies } \mu(E) = \lim_{n \rightarrow \infty} \mu(E_n). \end{aligned}$$

IV. THE INJECTIVE AND PROJECTIVE PROPERTIES OF LATTICE MEASURABLE FUNCTIONS

Definition 4.1. An extended real value function f defined on a lattice measurable E is said to be lattice measurable function if the set $\{x \in E / f(x) > \alpha\}$ is lattice measurable for all real numbers α .

Example 4.1. Constant functions, Continuous functions and Characteristic functions are lattice measurable functions.

Result 4.1. If f and g are lattice measurable functions then $f \vee g$ and $f \wedge g$ are also lattice measurable functions.

Proof. For any real number α we have $\{x \in L / (f \vee g)(x) > \alpha\} = \{x \in L / f(x) > \alpha\} \vee \{x \in L / g(x) > \alpha\}$ and $\{x \in L / (f \wedge g)(x) > \alpha\} = \{x \in L / f(x) > \alpha\} \wedge \{x \in L / g(x) > \alpha\}$. Since $\{x \in L / f(x) > \alpha\}$ and $\{x \in L / g(x) > \alpha\}$ are lattice measurable sets implies the sets of RHS are lattice measurable implies $f \vee g$ and $f \wedge g$ are lattice measurable functions.

The following interesting property can easily be verified from the works of [6] by considering lattice measurable functions f and g defined over countable Boolean lattice.

Property 4.1. A Countable Boolean lattice A is a Retrace of Countable Boolean lattice B if there exist homomorphism $g: A \rightarrow B$ and $f: B \rightarrow A$ such that fg is the identity on A . Here g and f are necessarily a monomorphism (injection) and epimorphism (projective) respectively. That is A Countable Boolean lattice is Retrace injective if it is a Retrace of every Countable Boolean lattice that contains it

V. CONCLUSION

New concepts like countable join property, countable meet property, P_σ -lattice and P_δ -lattice are introduced. Characterized partial lattices of a lattice through countable join and meet properties and proved that P_σ -lattice and P_δ -lattice are measurable partial lattices. Interesting result on the injective property of the lattice measurable functions defined over Countable Boolean lattices are established.

VI. ACKNOWLEDGEMENTS

Venkata Sundaranand Putcha is supported by DST-CMS project Lr.No.SR/S4/MS: 516/07, Dt.21-04-2008 and the support is gratefully acknowledged.

Ref

6. Halmos, R Paul, Injective and Projective Boolean Algebras, Research supported in part by a grant from the National science foundation. Lattice Theory, Proceedings of Symposia in pure mathematics Volume II, American Mathematical Society.

REFERENCES RÉFÉRENCES REFERENCIAS

1. D.V.S.R. Anil kumar, J. Venkateswara rao, E.S.R. Ravi kumar, Charecterization of Class of Measurable Borel Lattices, International Journal of Contemporary Mathematical Sciences, ISSN: 1312-7586, Volume 6(2011), no. 9, 439-446.
2. Birkhoff. G, Lattice Theory 3rd ed., AMS Colloquim Publications, Providence, RI, 1967.
3. Gene A. DE Both, Additive Set Functions on Lattices of Sets, Transactions of The American Mathematical Society, Volume 178, Apr 1973 pp 341-355.
4. Sen Gupta, Chapter1sigma Algebras www.math.lsu.edu/~sengupta/7312S02/sigmaalg.pdf.
5. Gratzner. G, General Lattice Theory, Academic Press Inc., 1978.
6. Halmos. R Paul, Injective and Projective Boolean Algebras, Research supported in part by a grant from the National science foundation. Lattice Theory, Proceedings of Symposia in pure mathematics Volume II, American Mathematical Society.
7. Halmos. P.R., Measure Theory (Springer, New York, 1974).
8. Pao-Sheng Hus Characterization of outer measures associated with lattice measures, International Journal of Mathematics and Mathematical Sciences. Vol.24, No.4 (2000) 237-249.
9. Szasz Gabor, Introduction to lattice theory, academic press, New York and London 1963.
10. Tanaka. J, Hahn Decomposition Theorem of Signed Lattice Measure, arXiv:0906.0147 Vol1 [Math.CA] 31 May 2009.

Notes