



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH
AGRICULTURE AND VETERINARY
Volume 13 Issue 8 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Determination of Genetic Diversity in Bread Wheat (*T.Aestivum* L.) by Agronomic and Quality Traits and SDS-PAGE Method

By Murat Olgun, Imren Kutlu, N. Gözde Ayter, Özcan Yorgancilar
& Zekiye Başçiftçi

Osmangazi University, Turkey

Abstract - The aim of this study was examine genetic diversities in bread wheat based on Agronomic and Quality Traits and SDS-PAGE Method. Bread wheat genotypes are Prostor, Eser, Daphan, Kinacı 97, Bağcı 2002, Tosunbey, Karahan 99, Katea 1, Nenehatun, Lancer, Karasu 90, Alparslan, Palandöken 97, Konya 2002 and Doğu 88 were used. Results revealed that Prostor wheat genotype differed from the other genotypes in three traits (gliadin subunits, agronomic and quality traits); Eser, Tosunbey and Kinacı 97 genotypes showed similarities in agronomic traits and gliadin subunits. Some genotypes presented similarities in each analysis and these similarities in terms of genotypes was different for each traits. Determination of genetic characteristic of genotypes in breeding programs provides genotype classification, genetic purity, similarities and dissimilarities early material selection. Gliadin pattern of wheat genotypes are free from environment, they are easy and forceful method in evaluation of genetic materials, breeding programs, pure seed productions in hexaploid wheat.

Keywords : *agronomic and quality traits, bread wheat, cluster analysis, correlation, SDS-PAGE method.*

GJSFR-D Classification : *FOR Code: 820507*



Strictly as per the compliance and regulations of :



Determination of Genetic Diversity in Bread Wheat (*T.Aestivum* L.) by Agronomic and Quality Traits and SDS-PAGE Method

Murat Olgun ^α, Imren Kutlu ^σ, N. Gözde Ayter ^ρ, Özcan Yorgancilar ^ω & Zekiye Başçiftçi [✱]

Abstract - The aim of this study was examine genetic diversities in bread wheat based on Agronomic and Quality Traits and SDS-PAGE Method. Bread wheat genotypes are Prostor, Eser, Daphan, Kinacı 97, Bağcı 2002, Tosunbey, Karahan 99, Katea 1, Nenehatun, Lancer, Karasu 90, Alparslan, Palandöken 97, Konya 2002 and Doğu 88 were used. Results revealed that Prostor wheat genotype differed from the other genotypes in three traits (gliadin subunits, agronomic and quality traits); Eser, Tosunbey and Kinacı 97 genotypes showed similarities in agronomic traits and gliadin subunits. Some genotypes presented similarities in each analysis and these similarities in terms of genotypes was different for each traits. Determination of genetic characteristic of genotypes in breeding programs provides genotype classification, genetic purity, similarities and dissimilarities early material selection. Gliadin pattern of wheat genotypes are free from environment, they are easy and forceful method in evaluation of genetic materials, breeding programs, pure seed productions in hexaploid wheat.

Keywords : agronomic and quality traits, bread wheat, cluster analysis, correlation, SDS-PAGE method.

1. INTRODUCTION

Wheat is the cultivated crop in the world. With the purpose of meet the increasing demand of food in tremendously increasing human population, aim in breeding programs is to develop high yielding cultivars possessing high quality, resistant to cold, drought and heat stress (Blum, 1988; Wilson, 1984). Success to overcome these troubles is not sufficient, sustainability in these properties of is also important, a number of studies have been made to increase for this purpose (Metakovsky and Branlard, 1998). Success in breeding programs is closely related to opulence in number of breeding materials affecting released cultivars. Reductions in the number of germplasm materials by using limited number of parents in breeding programs and diminish variation in new characters. Besides, keeping genetic purity in cultivars, in other words sustaining production pure seed production will create opportunity to address different market's demands, consequently, classification certain

physiological, physiological and quality traits vital (Akcura, 2006). Certain traits have been used to classify wheat genotypes such as physiological and quality traits and gliadin subunits (Weegels et al., 1996; Metakovsky et al., 1997; Metakovsky and Branlard, 1998; Ojaghi and Akhundova, 2010). Combinations for each trait create opportunity to distinguish genotypes each other (Ojaghi and Akhundova, 2010). Orth et al. (1996) stressed that many of physical and chemical tests could make the assessment of wheat genotypes for quality and these measurements strongly depend upon physical and chemical composition of wheat seed. Once physiological, physiological and quality traits are under genotype x environment interactions and a trait could be showed variations in different year or locations (Nevo et al., 1988). But gliadin sequence for each genotype is not affected by environment to identify genotypes (Zillman and Bushuk, 1979). Gliadins are the important seed proteins, evaluating and identifying genetic differences of genotypes could be safely made by them (Bushuk and Zillman 1978; Nevo and Payne 1987; Pfluger et al.2001). Reductions in the number of germplasm materials by using limited number of parents in breeding programs and diminish variation in new characters (Porceddu et al., 1988). Even tough morphometric characters (yield, yield components ect.) have been intensively used studies related to genetically diversity headed for protein bands such as gliadins (Metakovsky and Branlard, 1988). Gliadin (dissolved in etile alcohol, 70%) important part of wheat protein having high level of proline its named as prolamine (Zillman and Bushuk 1979). Gliadins as monomeric proteins could be classified to ω , γ , β , α groups (Jones et al.,1959). Most gliadin genes placed at first (Gli-1) and sixth (Gli-2) homology groups (Payne, 1987) and Gli-4 and Gli-5 groups in short arm of D chromosome (Rodrigve and Carrillo, 1996). Quality and agronomic traits have been commonly used in breeding programs, but determining quality and agronomic traits in genotypes needs long-term measurements to reach reliable results, since they are widely affected from environmental conditions (Metakovsky and Branlard, 1998; Ojaghi and Akhundova, 2010). Gliadin pattern of wheat genotypes are free from environment, they are easy and forceful method in evaluation of genetic materials, breeding programs, pure seed productions in hexaploid wheat.

Authors ^{α σ ρ ✱} : Osmangazi University, Faculty of Agriculture, Department of Field Crops, ESKİŞEHİR/ TURKEY.

E-mail : ikutlu@ogu.edu.tr

Author ^ω : Transitional Zone Agricultural Research Institute, ESKİŞEHİR/ TURKEY. E-mail : burakolg@hotmail.com

The aim of this study was to reveal genetic diversity by examining agronomic and quality traits, and gliadin patterns in bread wheat.

II. MATERIALS AND METHODS

Study was conducted in experimental station Osmangazi University Agricultural Faculty during 2009-2010 seasons. Long-term annual precipitation in Eskişehir province is 411.1 mm. Precipitation was 286.9 mm in 2009-2010 and 384.0 mm in 2010-2011 growing seasons. The soil had loamy-clay texture; 2.85 dS/m electrical conductivity, 7.21 pH (H₂O), 0.38% CaCO₃, 42,24 kg P₂O₅/ha available phosphorus, 3342 kg K₂O ha⁻¹ and 1.70% organic matter. In the study, ammonium sulphate (21%N) and triple super phosphate (46%) were used as fertilizers. Experimental design was randomised completed block design with three replication. Plot size was 6 m x 1.2 m (7.2m²). Seed were sown in 15 October and seed rate was 500 seed/m². 60 kg N/ ha (½ at sowing stage and ½ at tillering stage) and 60 kg P₂O₅/ha (at sowing) were applied. Wheat was sown in 1-15 September at the rate of 475 seed/m². Plot size was 6 m x 1.2 m (7.2m²) at sowing. No K fertilization was made. Plots received (2000 cc/ha) applications of 2,4-D ester [(2,4-dichlorophenoxy)acetic acid] in early spring to control winter annual broadleaf weeds. Plot size at harvest for determining grain yield was 0.80 m x 5 m (5.0

m²). Plots were harvested in 15th of July in 2009-2010 and 14th of July in 2010-2011. Bread wheat genotypes are Prostor, Eser, Daphan, Kinacı 97, Bağcı 2002, Tosunbey, Karahan 99, Katea 1, Nenehatun, Lancer, Karasu 90, Alparslan, Palandöken 97, Konya 2002 and Doğu 88 were used. Pedigree of wheat genotypes are given in Table 1. Agronomic characters; seed yield and plant height, seed number/spike, seed weight/spike, spikelet number, harvest index and 1000 grain weight (Slafer and Miralles, 1992) were evaluated. Quality characters; protein and gluten contents (Poehlman, 1987), zeleny sedimentation (Zeleny, 1947), farinograph, water absorbtion (Lehmensiek et al., 2006), alveograph, W_{energy} (Bettge et al., 1989), test weight (Sade et al., 1999), 1000 seed weight (Uluöz 1965) and PSI (Hruskova and Svec, 2009) were evaluated. Cluster analyses were made by MINITAB 16 Pocket Program. For gliadin subunits, sodium dodecylsulfate polyacrylamide gel electrophoresis (SDS-PAGE) analysis was used. Gliadin proteins were performed on vertical slab (Rashed et al., 2007). Relative mobility measures were used to evaluate gliadin band models. Relative motilities of bands were measured by formula (Keskin et al., 1996).

$Rm = \frac{\text{distance of bands measured from origin}}{\text{distance of reference band from origin} \times 45,5}$.

Table 1 : Pedigree of wheat genotypes

GENOTYPE	PEDIGREE	Releas. Year
Prostor	Russalka imp./Nadadores63	1999
Eser	AGRI/NAC//LIRA	2003
Daphan	JUP/4/CLLF/3/II4.53/ODIN//CI14431/WA00477	2002
Kinacı 97	YAMHILL/TOBARI-66//MCDERMID/3/LIRA	1997
Bağcı 2002	HN7/OROFEN//BJN8/3/SERI82/4/74CB462/TAPPER/VONA	2002
Tosunbey	EÇVD-12/KRÇ-66//CROW"S"	2004
Karahan 99	C-126-15/COLLAFEN/3/NORIN-10-BREVOR/P-14//PULLMAN-101/4/KIRAÇ-66	1999
Kate a 1	Hebros/Bez-1	1998
Nenehatun	ND//P101/BLUEBOY SWM584	2001
Lancer	TK/CNO//NE-451406	1977
Karasu 90	LOM11/B12973//MİR264	1990
Alparslan	TX69A509-2//BBY2/FOX	2001
Palandöken 97	AU//YT54*2/N10B/3/II8260/5/PNC/CM//NB6977/3/CC/INIA//BB/4/MXP//KR/FUNO	1997
Konya 2002	KANRED/TENMARG//P211-6/3/2183/CO652643/LANCER	2002
Doğu 88	Bez/Dane//CO725052	1990

III. RESULTS AND DISCUSSION

Genotypic/phenotypic diversity or classification in bread wheat genotypes are successfully made by using physiological, quality traits and gliadin subunits (Clarke et al., 1991; Yang et al., 1991; Metaovsky et al., 1990; Vaquero et al., 1990). Although physiological and quality traits are substantially connected to the environmental conditions, gliadin subunits are not

affected by environmental conditions (Zillman and Bushluk, 1979; Metakovsky and Branlard, 1998). Maximum, minimum values and means in agronomic and quality traits and relative mobility of gliadin subunits, are given Table 2. Introduction and selection are milestones in wheat breeding. To obtain information about genotypes are commonly revealing characteristics of genotypes by agronomic and quality and electrophoresis methods including protein analysis

(Soizow and Poperelya, 1980; Metakovsky and Branlard, 1998). Once gliadin structure remains unchanged through generations, agronomic and quality traits are under genotype x environment interactions, and electrophoresis is therefore safe method to describe protein characteristics of genotypes (Bushuk and Zillman, 1978; Metakovsky and Branlard, 1998). Table 1 sows that relative mobility of gliadin subunits were varied from 20,33 to 76,47 (20,33-25,29 in ω , 44,33-49,49,08 in γ , 54,33-62,35 in β and 71,00-76,47 in α). Depending upon increasing mobility, gliadin protein fractions are grouped in four main groups ω , γ , β , α gliadin subunits (Bietz and Wall, 1973; Kharebian et al., 2008). Gepts (1990) stressed that genotypes can be successfully grouped or evaluated according to relative mobility of different gliadin groups.

Researchers stated that gliadin structures (Nevo and Payne, 1987; Bushuk and Zillman, 1978), agronomic and quality traits of genotypes are differed (Lee and Ronalds, 1967; Wrigley, 1970). Whole three traits could safely be used in genotype classification (Metakovsky and Branlard, 1998; Ojaghi and Akhundova, 2010), but the safest method is gliadin electrophoresis method (Zillman and Bushuk, 1979). A number of researches demonstrated the relationship of gliadin polymorphism with genetic diversity (Nevo and Payne, 1987). In our study each gliadin band fractions are evaluated in subgroups, ω , γ , β and α and are shown in Figure 1.




Figure 1 : Gliadin band subunits, ω , γ , β and α in bread wheat genotypes

Table 2: Maximum, minimum values and means in agronomic and quality traits and relative mobility of gliadin subunits in bread wheat genotypes

Traits	Minimum	Maximum	Mean
Rm of Gliadin subunits			
ω	20,33	25,29	22,87±1,60
γ	44,33	49,08	46,19±1,96
β	54,33	62,35	63,38±1,40
α	71,00	76,47	73,45±1,59
Agronomic Traits			
Plant Height	85,40	142,47	109,12±14,83
Spike Heigh	7,73	11,35	9,61±1,02
Spikelet Number	14,40	18,67	16,38±1,27
Seed Number per Spike	28,47	57,87	38,92±8,36
Seed Weight per Spike	1,23	2,23	1,73±0,30
Harvest Index	42,34	56,67	25,08±7,21
Grain Yield	218,11	582,67	362,30±100,13

Quality Traits

PSI	54,34	75,33	67,99±6,31
Thousand Seed Weight	38,02	55,04	44,94±5,38
Test Weight	72,49	79,28	76,88±1,65
Protein Content	10,81	13,36	12,17±0,62
Zeleny Sedimentation	27,66	43,80	35,23±4,96
Alveograph W _{Energy}	96,01	263,42	161,72±44,25
Farihograph Water Absorbion	53,40	62,70	57,37±2,63
Gluten Content	23,00	38,00	28,00±3,87

We found notable polymorphisms among bread wheat genotypes on ω , γ , β and α gliadin subunits where γ and β have the heaviest molecular weights in Figure 1. Spectrum and density of molecular changed among ω , γ , β and α gliadin subunits in genotypes (Bietz and Wall, 1973; Nevo and Payne, 1987; Keskin et al., 1999). Kharabian et al. (2008) focused that ω has heaviest molecular weights. In our study, heavier molecular weights on γ , β gliadin subunits in Eser, Daphan, Kinacı 97, Tosunbey and Karahan 99 were found than that of the other genotypes. Moreover, differences in gliadin subunits assign differences of wheat genotypes, and ω , γ , β and α gliadin subunits are mostly located in Gli-A₁ and Gli-B₁ locus (Lafiandra and Kasarda, 1985; Kharabian et al., 2008), new gliadin genes in Gli-D₄ Gli-D₅ (Rodriguez and Carrillo, 1996). Rashed et al. (2007) focused that ω region were so rich in the number of bands. The dendrogram tree showed the similarity index of wheat cultivars detected by Rm of gliadin pattern (Figure 2).

The dendrogram divided wheat genotypes in seven clusters. Prostor, Konya 2002, Palandöken 97 and Karasu 90 placed in separate groups. Once Eser, Tosunbey, Karahan 99, Kinacı 97 and Daphan genotypes placed one group; Lancer, Alparslan, Doğu88 and Bağcı 2002 another group. Nevertheless,

Katea 1 and Nenehatun genotypes showed similar characteristics and placed same group (Figure 2). Gepts (1990) stressed that genotypes can be successfully grouped or evaluated according to relative mobility of different gliadin groups. According to Tanaka et al. (2003) wheat genotypes in Japan mainly different from other countries for gliadin pattern. Being main protein fragment in storage proteins, gliadins are important for bread making quality in hexaploid wheats (Kasarda et al., 1984). Nizar (2002) stated that determining gliadin pattern of wheat genotypes is safe method to classify cultivars and he successfully classified *T. durum* cultivars. Until recently, morphometric characters had been used in genetic diversities of wheat genotypes. Now, electrophoresis based techniques have been used for the determination and description of genotypes in cereals (Persson and Von Bothmer, 2000).

Genotypes segregate their own gliadin fraction; band pattern are characteristic of the genotype and aren't affected from environmental conditions (Bushuk and Zillman, 1978). Gliadin proteins having ω , γ , β , α gliadin subunits with combination of glutenins are play important role agronomic and bread making traits (Ojaghiani and Akhundova, 2010).




Figure 2 : Dendrogram assigning genetic relationships among bread wheat genotypes for gliadin patterns

Agronomic characters are essential corner stones of breeding programs (Darwinkel et al., 1977; Genc, 1977; Poehlman, 1987). A number of yield components are used not only in breeding programs but genotype description. Monitoring such yield and yield components separately or their relationships each other lights the way for breeding programs, agronomic studies and genotype classification (Hsu and Walton, 1971; Nass, 1973; McVetty and Evans, 1980). Plant and spike heights, spikelet and seed numbers per spike, seed weight per spike, harvest index and grain yield are some of the most used characters and they could be used in genotypic classification (Ojaghi and Akhundova, 2010). Similar to gliadin patterns, agronomic traits ranged 85,40-142, 47 cm in plant height, 7,73-11,35 cm in spike height, 14,40-18,67 in spikelet number per spike, 28,47-57,87 in seed number per spike, 1,23-2,23 gr in seed weight per spike, 42,34-56,6 in harvest index and 218,11-582,67 kg/da in grain yield (Table 2). Agronomic characters are commonly are named as yield components. The dendrogram tree of agronomic traits is given in Figure 3.

Evaluated in eight were wheat genotypes (Figure 3). Prostor, Konya 2002 and Katea 1 drew their own single groups. The other wheat genotypes formed dual or multiple groups: dual groups; Nenehatun and Doğu 88, Karasu 90 and Karahan 99, Alparslan and Lancer wheat genotypes; multiple group; Eser, Tosunbey, Daphan, Bağcı 2002 and Kınacı 97 wheat genotypes. Even though agronomic traits are under genotype x environment interactions, with certain approaches they are used to show genotypic differences (Metakovsky and Branlard, 1998; Ojaghi and Akhundova, 2010). Similar to our findings, Nenehatun, Doğu 88, Alparslan and Karasu 90 have similar yield

capacity and they are suggested to Eastern Anatolia climatic conditions. In the same way, Tosunbey, Eser, Kınacı 97 and Bağcı 2002 were developed for Central Anatolia climatic conditions.

The aim of bread wheat breeding programs is to develop high yielding and high-quality wheat cultivars. To achieve this aim, in addition of yield potential, wheat genotypes are evaluated for high quality such as for protein content, zeleny sedimentation, energy value such as alveograph W_{energy} , water absorption and test weight (Dexter et al. 1981; Atli 1999; Akcacak, 2006). Genotypes having higher values in protein content, zeleny sedimentation, alveograph W_{energy} , farinograph water absorption and test weight will be promising genotypes in breeding programs (Arat, 1949., Seçkin, 1970., Ünal, 1991., Atli, 1999). The dendrogram tree of quality traits is given in Figure 4.

Genotypes could be divided into nine groups (Figure 4). Prostor, Daphan, Konya 2002, Bağcı 2002 and Eser wheat genotypes had single groups. Nenehatun, Palandöken 97 The other wheat genotypes formed dual or multiple groups: dual groups; and Doğu 88, Karasu 90 and Karahan 99, Alparslan and Lancer wheat genotypes; multiple group; Eser, Tosunbey, Daphan, Bağcı 2002 and Kınacı 97 wheat genotypes. Even though agronomic traits are under genotype x environment interactions, with certain approaches they are used to show genotypic differences (Metakovsky and Branlard, 1998; Ojaghi and Akhundova, 2010). Similar to our findings, Nenehatun, Doğu 88, Alparslan and Karasu 90 have similar yield capacity and they are suggested to Eastern Anatolia climatic conditions. In the same way Tosunbey, Eser, Kınacı 97 and Bağcı 2002 were developed for Central Anatolia climatic conditions.




Figure 3 : Dendrogram of agronomic traits showing relationships among bread wheat genotypes




Figure 4 : Dendrogram of quality traits showing relationships among bread wheat genotypes

Table 3 shows joint analysis of wheat genotypes for gliadin subunits, agronomic and quality traits. Prostor wheat genotype differed from the other genotypes in three traits (gliadin subunits, agronomic and quality traits); Eser, Tosunbey and Kinacı 97 genotypes showed similarities in agronomic traits and gliadin subunits. As described earlier, some genotypes presented similarities in each analysis and these similarities in terms of genotypes was different for each trait (gliadin subunits, agronomic and quality traits).

Interesting phenomenon is that some genotypes could exhibit similarities by some traits (Eser, Tosunbey and Kinacı 97 genotypes in agronomic traits and gliadin subunits). or a genotype could significantly differ in all traits (Prostor genotype in all three traits). Studies revealed that agronomic and quality traits in wheat genotypes subject to genotype x environment interactions and this interaction differs in each genotype (Rodrigue and Carrillo, 1996; Şehirali and Özgen, 2007). Gliadin subunits stable with genotypes and don't change with changing environmental conditions (Bushuk and Zillman 1978; Rodrigue and Carrillo, 1996; Persson and Von Bothmer, 2000). Genetic diversity in genotypes can be controlled by polymeric markers and electrophoretic patterns of gliadin subunits in genotypes

are therefore very useful not only in genotype identification but in breeding programs (Nevo and Payne 1987; Metakovsky and Branlard, 1998; Pfluger et al., 2001). Studies stressed that agronomic and quality traits can be used in genotype classification (Kün, 1985; Şehirali, 2007), but both are influenced from genotype and environment conditions (Darwinkel et al., 1977; Genc, 1977; Poehlman, 1987; Gupta et al., 1993; Indrani and Venkateswara, 2000). Correlations between gliadin subunits, agronomic and quality traits are shown in Table 4. Positive and significant correlation ($P < 0.05$) was found between protein content and α gliadin pattern. Besides correlations between seed number per spike and ω gliadin pattern, seed number per spike and α gliadin pattern, seed number per spike and β gliadin pattern, seed weight per spike and ω gliadin pattern, spikelet number per spike and test weight, spikelet number per spike and thousand seed weight were determined as negative and significant ($P < 0.05$). Moreover, correlations between seed number per spike and γ gliadin pattern, seed number per spike and β gliadin pattern, seed weight per spike and ω , γ , β , α gliadin patterns, spike height and test weight were negative and significant at 1% (Table 4).

Table 3 : Joint analysis of wheat genotypes for gliadin subunits, agronomic and quality traits

Quality Traits	Prostor	Nenehatun	Palandöken 97	Lancer	Daphan	Kinacı 97	Karahan 99	Katea 1	Alparslan	Tosunbey	Karasu 90	Doğu 88	Konya 2002	Bağcı 2002	Eser
Gliadin Subunits	Prostor	Eser	Tosunbey	Karahan 99	Kinacı 97	Daphan	Konya 2002	Katea 1	Nenehatun	Lancer	Alparslan	Doğu 88	Bağcı 2002	Palandöken 97	Karasu 90
Agron. Traits	Prostor	Eser	Tosunbey	Bağcı 2002	Kinacı 97	Daphan	Nenehatun	Doğu 88	Konya 2002	Palandöken 97	Karahan 99	Karasu 90	Alparslan	Lancer	Katea 1

Table 4 : Correlations between gliadin subunits, agronomic and quality traits

	ω	γ	β	α		PSI	TeWe	ThSW	WAb
PSI	0,119	-0,010	0,185	0,308	Plant Height	-0,192	-0,194	0,154	0,297
Test Weight	0,104	0,210	-0,205	0,031	Spike Height	0,009	-0,689**	-0,246	-0,464
Thou. Seed We.	-0,049	-0,008	0,038	-0,320	Splt per Spk.	-0,071	-0,552*	-0,685*	-0,250
Protein Con	0,201	0,409	0,329	0,635*	Seed per Spk	-0,057	-0,553*	-0,177	-0,426
Zeleny Sedim.	0,128	0,178	-0,115	0,281	Seed W.per Spk	-0,093	-0,465	0,091	-0,347
Alveo. W _{en}	-0,061	0,237	-0,156	0,243	Harvest Index	-0,093	-0,189	0,392	-0,020
Farin. Water Ab	0,036	0,404	0,048	0,221	Yield	0,170	-0,219	0,439	-0,015
Gluten Index	-0,052	-0,035	-0,092	-0,105	PrCo	Sed	Wen	Gl	
Plant Height	-0,025	0,357	0,497	0,265	Plant Height	0,010	0,061	-0,078	0,086
Spike Height	-0,239	-0,367	0,155	-0,069	Spike Height	-0,279	-0,195	-0,410	0,198
Splt per Spk.	-0,105	-0,240	-0,163	0,009	Splt per Spk.	-0,378	0,357	-0,134	0,177
Seed per Spk	-0,623*	-0,717**	-0,633*	-0,584*	Seed per Spk	-0,361	-0,053	-0,256	0,010
Seed W.per Spk	-0,767**	-0,853**	-0,698**	-0,663**	Seed W.per Spk	-0,378	-0,279	-0,330	-0,133
Harvest Index	0,184	-0,176	-0,284	-0,370	Harvest Index	-0,208	-0,231	-0,318	-0,315
Yield	0,219	0,330	0,322	0,156	Yield	0,142	-0,397	-0,312	-0,330

Determinations in correlation between gliadin patterns and agronomic and quality traits are hard due to high molecular weight subunits of glutenin (Fido et al., 1997). Branlard and Dardevet (1985) found a negative correlation between ω gliadin and a number of quality traits. Whereas, positive correlation was found between γ gliadin subunit and gluten quality (Rashed et al., 2007). Significant positive effects of gliadin isoforms were found on agronomic traits and environmental adaptation (Metakosky and Branlard, 1998). Metakovsky and Branlard (1998) stated that genetic diversity in genotypes can be controlled by polymeric markers and electrophoretic patterns of gliadin subunits in genotypes. Determination of gliadin subunits is therefore very useful not only in genotype identification but in breeding programs (Bushuk and Zillman 1978; Persson and Von Bothmer, 2000).

So, quality and agronomic traits, and gliadin electrophoretic profiles revealed that comprehensive diversity occurs among genotypes. Determination of quality and agronomic traits in genotypes needs long-term measurements to reach reliable results and they are widely affected from environmental conditions (Vaquero et al., 1990; Metakovsky and Branlard, 1998; Ojaghi and Akhundova, 2010). Gliadin pattern of wheat genotypes are free from environment, they are easy and forceful method in evaluation of genetic materials, breeding programs, pure seed productions in hexaploid wheat. It is well known that yield is controlled polygenic activities and it is difficult to increase directly, however certain traits are significantly correlated with yield. Assigning and monitoring traits could help to increase yield and quality.

REFERENCES RÉFÉRENCES REFERENCIAS

1. Akçacık, A.G. (2006). Determination of suitable bread wheat and general and specific combination

ability and heritability parameters for quality and yield by using linextester analyses method. Ph. D. Thesis, Selçuk University, Department of Field Crops, Konya, 161 p.

2. Akçura M (2006) Characterization of turkish winter bread wheat landraces genetic resource. Ph. D. Thesis, Selcuk University, Department of Field Crop, Konya, 2006.
3. Arat SO (1949) Wheat technology. Ministry of Agriculture Directorate of Publications, Number: 654.
4. Atli A (1999) Quality of wheat and products. Symposium on Problems and Solutions in Agriculture in Central Anatolia Grain, 8-11 June, Konya.
5. Bettge A, Rubenthaler GL and Pomeranz Y (1989) Alveograph algorithms to predict functional properties of in bread and cookiebaking. Cereal Chemistry, 66: 81-86.
6. Bietz JA and Wall JS (1973) Isolation and characterization of gliadin-like subunits from glutenin. Cereal Chem. 50:537.
7. Blum A (1988) Plant breeding for stress environments. CRC Press Inc., Boca Raton, FL, 233 pp.
8. Branlard G and Dardevet M (1985) Diversity of grain protein and bread quality. I. Correlation between gliadin bands and technological characteristics. J Cereal Sci, 3:329-343.
9. Bushuk W and Zillman RR (1978) Wheat cultivar identification by gliadin electrophoregrams. I. apparatus, method and nomenclature. Can. J. Plant Sci., 58:505-515.
10. Clarke JM, Romagosa I and DePauw RM (1991) Screening durum wheat germplasm for dry growing conditions: Morphological and physiological criteria. Crop Sci., 31:770-775.

11. Darwinkel A, Hag BA and Kuizenga J (1977) Effect of sowing date and seed rate on crop development and grain production of winter wheat. Netherlands J. Agric. Sci. 24:83-94.
12. Dexter JE and Matsuo RR (1981) Effect of starchy kernels, immaturity, and shrunken kernels on durum wheat quality. Cereal Chem. 58:395-400.
13. Fido RJ, Bekest F, Grast PW and Tahtam AS (1997) Effects of α -, β -, γ -, and ω - Gliadins on Dought Mixing Properties of Wheat Flour. Journal of Cereal Science, 26:271-277.
14. Genç I (1977) Physiological and morphological basis in cereals. Çukurova Uni. Agric. Fac. Press. No: 22, Ankara.
15. Gepts P (1990) Biochemical evidence bearing on the domestication of *Phaseolus* (Fabaceae) beans. Econ. Bot. 44 (3 Suppl.):28-38.
16. Hruskova M, and Svec I (2009) Wheat hardness in relation to other quality factors. Czech Journal of Food Sciences 27(4):240-248.
17. Hsu P and Walton PD (1970) The inheritance of morphological and agronomic characters in spring wheat. Euphytica. 19:54-60.
18. Jones RW, Taylor NW and Senti FR (1959) Electrophoresis and fractionation of wheat gluten. Arch. Biochem. Biophys. 84:363-376.
19. Kasarda DD, Okita TW, Bernardin JE, Baecker PA, Nimmo CC, Lew EJL, Dietler MD and Greene FC (1984) Nucleic acid (dDNA) and amino acid sequences of α -type gliadins from wheat (*Triticum aestivum*). Proc. Natl. Acad. Sci. USA, 81: 4712-4716.
20. Keskin S, Asal S and Kavuncu O (1996) Genetic analysis and gliadin banding patterns in some bread wheat varieties grown in turkey and their hybrids. Tr. J. of Agriculture and Forestry 23: 291-298.
21. Kharabian A, Jones A and Kaviani B (2008) Electrophoretic profiles of gliadin subunits to evaluate genetic diversity of *Triticum persicum* Boiss. and *Triticum pyramidale* Pervical. Plant Omics Journal, 1(1):9-16.
22. Kün E, (1985) Hot climate cereals. Ankara University Faculty of Agriculture Publications, Ankara. 1985
23. Lafiandra D and Kasarda DD (1985) One- and two-dimensional (two pH) polyacrylamide gel electrophoresis in a single gel: separation of wheat proteins. Cereal Chem 62:314-319.
24. Lee JW and Ronalds JA (1967) Effect of environment on wheat gliadin. Nature (London) 213: 844-846.
25. Lehmensiek A, Eckermann PJ, Verbyla AP, Appels R, Sutherland MW, Martin D, Daggard GE, (2006) Flour yield QTLs in three Australian doubled haploid populations. Australian Journal of Agricultural Research 57:1115-1122.
26. McVetty PBE and Evans LE (1980) Breeding methodology in wheat. I. Determination of characters measured on F spaced plants for yield selection in spring wheat. Crop Sci. 20:584-586.
27. Metakovsky EV, Wrigley CW, Bekes F and Gupta RB (1990) Gluten polypeptides as useful genetic markers of dough quality in Australian wheats. Aust J Agr Res 41:289-306.
28. Metakovsky EV, Annicchiarico P, Baggini G, Pogana NE (1997) Relationship between gliadin alleles and dough strength in Italian bread wheat cultivars. J. Cereal Sci. 25:229-236.
29. Metakovsky EV, Branlard G (1998) Genetic diversity of French common wheat germplasm based on gliadin alleles. Theor. Appl. Genet. 96:209-218.
30. Nass HG (1973) Determination of characters for yield selection in spring wheat. Can. J. Plant Sci. 53: 755-762
31. Nevo E, Payne PL (1987) Wheat storage proteins:diversity of HMWglutenin subunits in wild emmer from Israel.Geographical patterns and ecological predictability. Theoretical and Appl. Genet. 74: 827-836.
32. Nevo E, Beiles A and Krugman T (1988) Natural selection of allozyme polymorphisms: a microgeographical differentiation by edaphic, topographical, and temporal factors in wild emmer wheat (*Triticum dicoccoides*). Theor Appl Genet, 76: 737-752
33. Nizar MA (2002) Gliadins polymorphism and cluster analyses of Syrian grown durum wheat. Plant Breeding and Seed Science, 46: 45-56,
34. Ojaghi J and Akhundova E (2010) Genetic diversity in doubled haploid wheat according to the acid-PAGE method, morphological traits and baking quality.Turk J Biol. 34: 343-353
35. Orth RA, O'Brien L and Jardine C (1976) A factor analysis of bread wheat quality tests. Aust. J. Agric. Res.,27:575-82
36. Payne PI (1987) Genetics of wheat storage protein and the effect of allelic variation on bread baking quality. Annu. Rev. Plant physiol.38: 141-153.
37. Persson K, Von Bothmer R (2000) Assessing the variation in cultivars and Swedish landraces of rye (*Secale cereale* L.) Hereditas, 13: 7-17.
38. Pflüger LA, D'Ovidio R, Margiotta B and Pena R (2001) Characterisation of high- and low molecular weight glutenin subunits associated to the D genome of *Aegilops tauschii* in a collection of synthetic hexaploid wheats. Theor Appl Genet. 103: 1293-1301.
39. Poehlman JM (1987) Breeding field crops, Van Nostrand Reinhold Company Inc. 115 Fifth Avenue New York.
40. Porceddu E, Ceoloni C, Lafiandra D, Tanzarella OA and Scarascia Mugnozza GT (1988) Genetic

- resources and plant breeding: problems and prospects. In: Miller TE, Koebner RMD (Eds.), Proceedings of the Seventh International Wheat Genetics Symposium, Institute of Plant Science Research, Cambridge, pp. 7–22.
41. Rashed MA, Abou-Deif MH, Sallam MAA, Aida Rizkalla A, Walaa Ramada A (2007) Identification and prediction of the flour quality of bread wheat by gliadin electrophoresis, Journal of Applied Sciences Research, 3 (11): 1393-1399.
 42. Sade B, Topal A and Soylu S (1999) Determination of durum wheat varieties grown under irrigated conditions in Konya. Turkey 3. Field Crops Congress: 91-96. 15-20 November, Adana.
 43. Seçkin R (1970) The factors that affect of wheat quality and composition. Ankara University Agricultural Faculty Publications.430 Conferences Series 8., Ankara.
 44. Slafer GA and Miralles DJ (1992) Green area duration during the grain filling period of wheat as affected by sowing date, temperature and sink strength. J Agron Crop Sci 168: 191–200.
 45. Şehirali S and Özgen M (2007) Plant breeding. Ankara University Agricultural Faculty Publications, 1553/506, 270 p, Ankara.
 46. Tanaka H, Tomita M, Tsujimoto H, Yasumuro Y (2003) Limited but specific variation of seed storage proteins in Japanese common wheat (*Triticum aestivum* L.). Euphytica, 132:167-174.
 47. Uluöz M (1965) Analysis methods of wheat, flour and bread. A.U. Agricultural Faculty Publications 29, 91 s .
 48. Ünal S (1991), Cereal techtonology. A.U. Agricultural Faculty Publications, 29. İzmir
 49. Vaquero F, Rebordinos, L, Vences FJ and Perez de la Vega M (1990) Genetic mapping of isozyme loci in *Secale cereale* L. Theor. Appl. Genet., 80: 88-94.
 50. Weegels PL, Hamer RJ, Schofield JD (1996) Functional properties of wheat glutenins. J. Cereal Sci. 23:1-17.
 51. Wilson JA (1984) Hybride wheat breeding and commercial seed development. Plant Breeding Reviews,2: 303-319.
 52. Wrigley CW (1970) Protein mapping by combined gel electrofocusing and electrophoresis: Application to the study of genotypic variations in wheat gliadins. Biochem. Genet. 4: 509.
 53. Yang RC, Jana S and Clarke JM (1991) Phenotypic diversity and association of some potentially drought responsive characters in durum wheat. Crop Sci., 31:1484-1491
 54. Zeleny L (1947) A simple sedimentation test for estimating the bread-baking and gluten qualities of wheat flour. Cereal Chem. 24: 465-475.
 55. Zillman RR, Bushuk W (1979) Wheat cultivar identification by gliadin electrophoregrams. II. Effects of environmental and experimental factors on the gliadin electrophoregrams. Can. J. Plant Sci. 59: 281-286.

