



GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH
MATHEMATICS AND DECISION SCIENCES

Volume 13 Issue 11 Version 1.0 Year 2013

Type : Double Blind Peer Reviewed International Research Journal

Publisher: Global Journals Inc. (USA)

Online ISSN: 2249-4626 & Print ISSN: 0975-5896

A Note on Basic Hypergeometric Function of N-Variable

By Pankaj Srivastava & Mohan Rudravarapu

Motilal Nehru National Institute of Technology, India

Abstract- In this paper an attempt has been made to establish new transformation formula for the basic hypergeometric function of n variable.

Keywords: q-series, basic hypergeometric function, transformation.

GJSFR-F Classification : MSC 2010: 33C05, 33D70



Strictly as per the compliance and regulations of:



© 2013. Pankaj Srivastava & Mohan Rudravarapu. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License <http://creativecommons.org/licenses/by-nc/3.0/>), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.



R_{ef}

A Note on Basic Hypergeometric Function of N-Variable

Pankaj Srivastava^α & Mohan Rudraravapu^σ

Abstract- In this paper an attempt has been made to establish new transformation formula for the basic hypergeometric function of n variable.

Keywords: q-series, basic hypergeometric function, transformation.

I. INTRODUCTION

Basic hypergeometric functions are among the most important functions with very diverse applications to Engineering, Physics and Mathematical Analysis. Nowadays, the importance of basic hypergeometric function and its unique role as the strategic resource in the Ramanujan's Mathematics became more important than the past time. The importance given to the Ramanujan's Mathematics has increased considerably in recent years and great deal of attention in basic hypergeometric function's literature is being given for the numerous topics that have been addressed by mathematicians working in the field of Basic hypergeometric functions, notably R.P Agarwal [1], G.E. Andrews and B.C. Berndt [2], G.E. Andrews [3, 4], R.Askey [5, 6], W.N. Bailey [7], S. Bhargava and Chandrashekar Adiga [8], R.Y.Denis [9], R.Y. Denis *et al.* [10], G.Gasper [11], V.K.jain [12, 13], M.S. Mahadeva Naika and B.N. Dharmendra [14], T.H.M.Rassias and S.N. Singh [15], L.J.Slater [16], Pankaj Srivastava [17, 18], Pankaj Srivastava and Mohan Rudraravapu [19], A. Verma [20], G.N.Watson [21] and many others published large number of studies. In this paper, we are interested to develop certain new transformation formula for the basic hypergeometric function of n variable with the help of technique developed by Andrews [4], special cases also developed.

II. NOTATIONS AND DEFINITIONS

A basic hypergeometric series of n-variables is defined as

$$\phi \left[\begin{matrix} (a_p) : (b_{M_1}^1); (b_{M_2}^2); \dots; (b_{M_n}^n) \\ (c_t) : (d_{N_1}^1); (d_{N_2}^2); \dots; (d_{N_n}^n) \end{matrix} ; x_1, x_2, \dots; x_n \right]$$

Author α: Department of Mathematics, Motilal Nehru National Institute of Technology, Allahabad - 211 004, Uttar Pradesh, India.
e-mail: mohanrudraravapu@hotmail.com

Author σ: Department of Mathematics, Motilal Nehru National Institute of Technology, Allahabad - 211 004, Uttar Pradesh, India
e-mail: mohanrudraravapu@hotmail.com

[1] R.P.Agarwal, Certain basic hypergeometric identities associated with Mock theta functions, *Quarterly Journal Of Mathematics*, **20(1)** (1969), 121-128.

Year 2013

123

Version I

XI

Issue

Volume XIII

(F)

Research

Frontier

Journal of

Science

Global

Inc.

(US)

$$= \sum_{m_1, m_2, \dots, m_n \geq 0}^{\infty} \frac{[(a_p)]_{m_1+m_2+\dots+m_n} [(b_{M_1}^1)]_{m_1} \dots [(b_{M_n}^n)]_{m_n} x_1^{m_1} \dots x_n^{m_n}}{[q]_{m_1} \dots [q]_{m_n} [(c_t)]_{m_1+m_2+\dots+m_n} [(d_{N_1}^1)]_{m_1} \dots [(d_{N_n}^n)]_{m_n}}, \quad (2.1)$$

Where (a_p) stands for p -parameters $a_1, a_2, \dots, a_p$. For the convergence of this series we require $\max(|q|, |x_1|, \dots, |x_n|) < 1$. The q -shifted factorial is defined as

$$(a; q)_n = \begin{cases} 1, & n = 0 \\ (1-a)(1-aq) \dots (1-aq^{n-1}), & n = 1, 2, 3, \dots \end{cases} \quad (2.2)$$

We also define

$$(a)_{\infty} = (a; q)_{\infty} = \prod_{k=0}^{\infty} (1-aq^k), \text{ for } |q| < 1. \quad (2.3)$$

The infinite product diverges when $a \neq 0$.

Also

$$(a_1, a_2, \dots, a_r; q)_n = (a_1; q)_n (a_2; q)_n \dots (a_r; q)_n. \quad (2.4)$$

And

$${}_1\phi_0 [a; -; q, z] = \frac{(az; q)_{\infty}}{(z; q)_{\infty}}, |z| < 1, |q| < 1. \quad (2.5)$$

The product formula

$${}_1\phi_0 (a; -; q, z) {}_1\phi_0 (b; -; q, az) = {}_1\phi_0 (ab; -; q, z). \quad (2.6)$$

$${}_1\phi_1 [a; c; q, c/a] = \frac{(c/a; q)_{\infty}}{(c; q)_{\infty}}. \quad (2.7)$$

III. MAIN RESULTS

In this section we shall establish the following main result .

$$\begin{aligned} & \phi \left[\begin{matrix} a, (a_p) : (b_{M_1}^1); (b_{M_2}^2); \dots; (b_{M_n}^n) \\ c, (c_t) : (d_{N_1}^1); (d_{N_2}^2); \dots; (d_{N_n}^n) \end{matrix} ; x_1, x_2, \dots, x_n \right] \\ &= \frac{[a]_{\infty}}{[c]_{\infty}} \sum_{r=0}^{\infty} \frac{[c/a]_r a^r}{[q]_r} \phi \left[\begin{matrix} (a_p) : (b_{M_1}^1); \dots; (b_{M_n}^n) \\ (c_t) : (d_{N_1}^1); \dots; (d_{N_n}^n) \end{matrix} ; x_1 q^r, x_2 q^r, \dots, x_n q^r \right]. \quad (3.1) \end{aligned}$$

IV. PROOF OF (3.1)

By using (2.1) the left hand side of (3.1) can be put in the following form

$$\begin{aligned}
 & \sum_{m_1, m_2, \dots, m_n \geq 0}^{\infty} \frac{[a]_{m_1+m_2+\dots+m_n} [(a_p)]_{m_1+m_2+\dots+m_n}}{[q]_{m_1} \dots [q]_{m_n} [c]_{m_1+m_2+\dots+m_n} [(c_t)]_{m_1+m_2+\dots+m_n}} \times \\
 & \quad \frac{[(b_{M_1}^1)]_{m_1} \dots [(b_{M_n}^n)]_{m_n} x_1^{m_1} \dots x_n^{m_n}}{[(d_{N_1}^1)]_{m_1} \dots [(d_{N_n}^n)]_{m_n}} \\
 &= \frac{[a]_{\infty}}{[c]_{\infty}} \sum_{m_1, m_2, \dots, m_n \geq 0}^{\infty} \frac{[(a_p)]_{m_1+m_2+\dots+m_n} [(b_{M_1}^1)]_{m_1} \dots [(b_{M_n}^n)]_{m_n}}{[q]_{m_1} \dots [q]_{m_n} [(c_t)]_{m_1+m_2+\dots+m_n}} \times \\
 & \quad \frac{(cq^{m_1+\dots+m_n})_{\infty} x_1^{m_1} \dots x_n^{m_n}}{[(d_{N_1}^1)]_{m_1} \dots [(d_{N_n}^n)]_{m_n} (aq^{m_1+m_2+\dots+m_n})_{\infty}} \\
 &= \frac{[a]_{\infty}}{[c]_{\infty}} \sum_{m_1, \dots, m_n \geq 0}^{\infty} \frac{[(a_p)]_{m_1+\dots+m_n} [(b_{M_1}^1)]_{m_1} \dots [(b_{M_n}^n)]_{m_n} x_1^{m_1} \dots x_n^{m_n}}{[q]_{m_1} \dots [q]_{m_n} [(c_t)]_{m_1+m_2+\dots+m_n} [(d_{N_1}^1)]_{m_1} \dots [(d_{N_n}^n)]_{m_n}} \\
 & \quad \times \sum_{r=0}^{\infty} \frac{(c/a)_r a^r q^{r(m_1+m_2+\dots+m_n)}}{[q]_r}.
 \end{aligned}$$

By changing the order of summation in the above equation, the right hand side of (3.1) can be obtained.

V. PARTICULAR CASES

Putting $p = t = M_1 = \dots = M_n = N_1 = \dots = N_n = 0$ in (3.1), we get

$$\begin{aligned}
 & {}_1\phi_1 \left[\begin{matrix} a : -; \dots; -; \\ c : -; \dots; -; \end{matrix} ; x_1, x_2, \dots; x_n \right] \\
 &= \frac{[a]_{\infty}}{[c]_{\infty}} \sum_{r=0}^{\infty} \frac{[c/a]_r a^r}{[q]_r} \times {}_0\phi_0 [-; -; x_1 q^r] \dots {}_0\phi_0 [-; -; x_n q^r] \\
 &= \frac{[a]_{\infty}}{[c]_{\infty}} \sum_{r=0}^{\infty} \frac{[c/a]_r a^r}{[q]_r [x_1 q^r]_{\infty} \dots [x_n q^r]_{\infty}} \\
 &= \frac{[a]_{\infty}}{[c]_{\infty}} \sum_{r=0}^{\infty} \frac{[c/a]_r a^r [x_1]_r \dots [x_n]_r}{[q]_r [x_1]_{\infty} \dots [x_n]_{\infty}} \\
 &= \frac{[a]_{\infty}}{[c]_{\infty} [x_1]_{\infty} \dots [x_n]_{\infty}} {}_{n+1}\phi_0 [x_1, \dots, x_n, c/a; -; a],
 \end{aligned}$$

$$\text{which is valid if } |a| < 1. \quad (5.1)$$

Putting $p = t = N_1 = \dots = N_n = 0, M_1 = \dots = M_n = 1, b_1^1 = b_1, b_1^2 = b_2, \dots, b_1^n = b_n$ in (3.1), we get:

$$\begin{aligned}
 & \phi \left[\begin{matrix} a : b_1; \dots; b_n \\ ; x_1, x_2, \dots; x_n \\ c : -; \dots; - \end{matrix} \right] \\
 &= \frac{[a]_{\infty}}{[c]_{\infty}} \sum_{r=0}^{\infty} \frac{[c/a]_r a^r}{[q]_r} \times {}_1\phi_0 [b_1; -; x_1 q^r] \dots {}_1\phi_0 [b_n; -; x_n q^r] \\
 &= \frac{[a]_{\infty}}{[c]_{\infty}} \sum_{r=0}^{\infty} \frac{[c/a]_r a^r}{[q]_r} \frac{[b_1 x_1 q^r]_{\infty}}{[x_1 q^r]_{\infty}} \dots \frac{[b_n x_n q^r]_{\infty}}{[x_n q^r]_{\infty}} \\
 &= \frac{[a]_{\infty} [b_1 x_1]_{\infty} \dots [b_n x_n]_{\infty}}{[c]_{\infty} [x_1]_{\infty} \dots [x_n]_{\infty}} \times \sum_{r=0}^{\infty} \frac{[c/a]_r a^r [x_1]_r \dots [x_n]_r}{[q]_r [b_1 x_1]_r \dots [b_n x_n]_r} \\
 &= \frac{[a]_{\infty} [b_1 x_1]_{\infty} \dots [b_n x_n]_{\infty}}{[c]_{\infty} [x_1]_{\infty} \dots [x_n]_{\infty}} \times {}_{n+1}\phi_n [x_1, \dots, x_n, c/a; b_1 x_1, \dots, b_n x_n; a], \quad (5.2)
 \end{aligned}$$

Putting $x_3 = x_4 = \dots = x_n = 0$ in (5.2), we get

$$\begin{aligned}
 & \phi \left[\begin{matrix} a : b_1; b_2; \dots; \\ ; x_1, x_2 \\ c : -; \dots; - \end{matrix} \right] = \phi^{(1)} [a; b_1, b_2; c; x_1, x_2] \\
 &= \frac{[a]_{\infty} [b_1 x_1]_{\infty} [b_2 x_2]_{\infty}}{[c]_{\infty} [x_1]_{\infty} [x_2]_{\infty}} \times {}_3\phi_2 \left[\begin{matrix} x_1, x_2, c/a; \\ ; a \\ b_1 x_1, b_2 x_2 \end{matrix} \right], \quad (5.3)
 \end{aligned}$$

This is the result due to Denis ([9], 5.5).

Now, putting $b_2 = x_1/x_2, b_1 x_1 = c x_2$ in (5.3) and evaluating ${}_2\phi_1$ series on the right hand side with the help of Slater ([16]; Appendix.IV.2), we get

$$\phi^{(1)} [a; c x_2/x_1, x_1/x_2; c; x_1, x_2] = \frac{[x_2 a]_{\infty}}{[x_2]_{\infty}}. \quad (5.4)$$

Now, substituting $p = t = 0, N_1 = \dots = N_n = 1, M_1 = \dots = M_n = 1, b_1^1 = b_1, b_1^2 = b_2, \dots, b_1^n = b_n, d_1^1 = d_1, d_1^2 = d_2, \dots, d_1^n = d_n$ in (3.1), we get:

$$\begin{aligned}
 & \phi \left[\begin{matrix} a : b_1; \dots; b_n \\ ; x_1, x_2, \dots; x_n \\ c : d_1; \dots; d_n \end{matrix} \right] \\
 &= \frac{[a]_{\infty}}{[c]_{\infty}} \sum_{r=0}^{\infty} \frac{[c/a]_r a^r}{[q]_r} \times {}_1\phi_1 [b_1; d_1; x_1 q^r] \dots {}_1\phi_1 [b_n; d_n; x_n q^r], \quad (5.5)
 \end{aligned}$$

Taking $d_1 = b_1 x_1 q^r, d_2 = b_2 x_2 q^r, \dots, d_n = b_n x_n q^r$ in (5.5), we get the following result after simplification

$$\phi \left[\begin{matrix} a : b_1; \dots; b_n \\ c : b_1 x_1 q^r; \dots; b_n x_n q^r \end{matrix} ; x_1, x_2, \dots; x_n \right] \\ = \frac{[a]_{\infty} [x_1]_{\infty} \dots [x_n]_{\infty}}{[c]_{\infty} [b_1 x_1]_{\infty} \dots [b_n x_n]_{\infty}} \times {}_{n+1}\phi_n [b_1 x_1, \dots, b_n x_n, c/a; x_1, \dots, x_n; a], \quad (5.6)$$

Now, putting $x_3 = x_4 = \dots = x_n = 0$ in (5.6), we get

$$\phi \left[\begin{matrix} a : b_1; b_2 \\ c : b_1 x_1 q^r; b_2 x_2 q^r \end{matrix} ; x_1, x_2 \right] \\ = \frac{[a]_{\infty} [x_1]_{\infty} [x_2]_{\infty}}{[c]_{\infty} [b_1 x_1]_{\infty} [b_2 x_2]_{\infty}} \times {}_3\phi_2 \left[\begin{matrix} b_1 x_1, b_2 x_2, c/a; \\ x_1, x_2 \end{matrix} ; a \right], \quad (5.7)$$

Now, substituting $b_1 x_1 = x_2, b_2 x_2 = x_1/c$ in (5.7) and evaluating ${}_2\phi_1$ series in the right hand side with the help of Slater ([16]:Appendix IV.2), we get

$$\phi \left[\begin{matrix} a : x_2/x_1; x_1/cx_2 \\ c : x_2 q^r; x_1 q^r/c \end{matrix} ; x_1, x_2 \right] = \frac{(ax_1/c; q)_{\infty}}{(x_1/c; q)_{\infty}}. \quad (5.8)$$

A variety of similar interesting results can be scored.

REFERENCES RÉFÉRENCES REFERENCIAS

- [1] R.P.Agarwal, Certain basic hypergeometric identities associated with Mock theta functions, *Quarterly Journal Of Mathematics*, **20(1)** (1969), 121-128.
- [2] G.E.Andrews and B.C.Berndt, Ramanujan's Lost Notebook, Part1, *Springer, New York*, (2005).
- [3] G.E.Andrews, Applications of basic hypergeometric functions, *SIAM Rev*, **16** (1974a), 441-484.
- [4] G.E.Andrews, Problems and properties for basic hypergeometric function, theory and applications of special functions, *Academic press, New York*, (1975).
- [5] R.Askey, Some basic hypergeometric extensions of integrals of Selberg and Andrews, *SIAM J.Math.Anal*, **11** (1980b), 938-951.
- [6] R.Askey, Ramanujan's ${}_1\phi_1$ and formal Laurent series, *Indian J.Math*, **29** (1987), 101-105.
- [7] W.N.Bailey, Well poised basic hypergeometric series, *Quart.J.Math, Oxford*. **18** (1947b), 157-166.

- [8] S.Bhargava and Chandrashekar Adiga, On some continued fraction identities of Srinivasa Ramanujan, *Proc. Amer. Math. Soc.*, **92** (1984), 13-18.
- [9] R. Y. Denis, On certain transformations of basic hypergeometric functions, *Ranchi Univ. Math.*, **7.13** (1983), 44-51.
- [10] R.Y.Denis, S.N.Singh and S.P.Singh, On single series representation of mock theta functions of fifth and seventh order, *Italian Journal Of Pure and Applied Mathematics-N*, **23** (2008), 67-74.
- [11] G.Gasper, Summation formulas for basic hypergeometric series, *SIAM J. Math. Anal.*, **12** (1981a), 196-200.
- [12] V.K.Jain, Some expansions involving basic hypergeometric functions of two variables, *Pac. J. Math.*, **91** (1980a), 349-361.
- [13] V.K.Jain, Some transformations of basic hypergeometric series and their applications, *Proc. Amer. Math. Soc.*, **78** (1980b), 375-384.
- [14] M.S.Mahadeva Naika and B.N.Dharmendra, On Some new general theorems for the explicit evaluations of Ramanujan's remarkable product of Theta functions, *The Ramanujan Journal*, **15(3)** (2008), 349-366.
- [15] TH.M.Rassias and S.N.Singh, Certain transformations involving basic hypergeometric series, *Complex Variables*, **18** (1992), 3-6.
- [16] L. J. Slater, *Generalized hypergeometric functions*, Cambridge At the university Press, (1966).
- [17] Pankaj Srivastava, Certain transformations of generalized Kampe' de' Fe'riet function, *Vijnana Parishad Anusandan Patrika*, **45(4)** (2002), 383-392.
- [18] Pankaj Srivastava, Resonance of continued fractions related to ${}_2 {}_2$ basic bilateral hypergeometric series, *Kyungpook Math. J.*, **51** (2011), 419-427.
- [19] Pankaj Srivastava and Mohan Rudravarapu, Certain Transformation Formulae for the Generalized Hypergeometric Series of Multi Variables, *Australian Journal of Basic and Applied Sciences*, **5(11)** (2011), 496-517.
- [20] A.Verma, Certain expansions of the basic hypergeometric functions, *Math. Comp.*, **20** (1966), 151-157.
- [21] G.N.Watson, A new proof of the Rogers-Ramanujan identities, *J. London Math. Soc.*, **4** (1929a), 4-9.