Online ISSN: 2249-4626
Print ISSN: 0975-5896

Global Journal

of Science Frontier Research: B

Chemistry



Banana Peels Powder

Hydrophobic Carbazole Ring

Highlights

Water Quality of River

Phenyl Sulfonyl Pyrazoline

Discovering Thoughts, Inventing Future

VOLUME 14 ISSUE 1 VERSION 1.0

© 2001-2014 by Global Journal of Science Frontier Research , USA

Volume 14 Issue 2 was originally issued as B1

Global Journal of Science Frontier Research: B
Chemistry

Volume 14 Issue 2
Originally issued as B1

Global Journal of Science Frontier Research: B
Chemistry

Volume 14 Issue 1 (Ver. 1.0)

Open Association of Research Society

© Global Journal of Science Frontier Research. 2014.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Science Frontier Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Science Frontier Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Science Frontier Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website http://globaljournals.us/terms-and-condition/menu-id-1463/

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089
License No.: 42125/022010/1186
Registration No.: 430374
Import-Export Code: 1109007027
Employer Identification Number (EIN):
USA Tax ID: 98-0673427

Global Journals Inc.

*(A Delaware USA Incorporation with "Good Standing"; **Reg. Number: 0423089**)*

Sponsors: Open Association of Research Society
Open Scientific Standards

Publisher's Headquarters office

Global Journals Headquarters
301st Edgewater Place Suite, 100 Edgewater Dr.-Pl,
Wakefield MASSACHUSETTS, Pin: 01880,
United States of America
USA Toll Free: +001-888-839-7392
USA Toll Free Fax: +001-888-839-7392

Offset Typesetting

Global Journals Incorporated
2nd, Lansdowne, Lansdowne Rd., Croydon-Surrey,
Pin: CR9 2ER, United Kingdom

Packaging & Continental Dispatching

Global Journals
E-3130 Sudama Nagar, Near Gopur Square,
Indore, M.P., Pin:452009, India

Find a correspondence nodal officer near you

To find nodal officer of your country, please email us at *local@globaljournals.org*

eContacts

Press Inquiries: *press@globaljournals.org*
Investor Inquiries: *investors@globaljournals.org*
Technical Support: *technology@globaljournals.org*
Media & Releases: *media@globaljournals.org*

Pricing (Including by Air Parcel Charges):

For Authors:
22 USD (B/W) & 50 USD (Color)
Yearly Subscription (Personal & Institutional):
200 USD (B/W) & 250 USD (Color)

Integrated Editorial Board
(Computer Science, Engineering, Medical, Management, Natural Science, Social Science)

John A. Hamilton,"Drew" Jr.,
Ph.D., Professor, Management
Computer Science and Software Engineering
Director, Information Assurance Laboratory
Auburn University

Dr. Henry Hexmoor
IEEE senior member since 2004
Ph.D. Computer Science, University at Buffalo
Department of Computer Science
Southern Illinois University at Carbondale

Dr. Osman Balci, Professor
Department of Computer Science
Virginia Tech, Virginia University
Ph.D.and M.S.Syracuse University, Syracuse, New York
M.S. and B.S. Bogazici University, Istanbul, Turkey

Yogita Bajpai
M.Sc. (Computer Science), FICCT
U.S.A.Email:
yogita@computerresearch.org

Dr. T. David A. Forbes
Associate Professor and Range Nutritionist
Ph.D. Edinburgh University - Animal Nutrition
M.S. Aberdeen University - Animal Nutrition
B.A. University of Dublin- Zoology

Dr. Wenying Feng
Professor, Department of Computing & Information Systems
Department of Mathematics
Trent University, Peterborough,
ON Canada K9J 7B8

Dr. Thomas Wischgoll
Computer Science and Engineering,
Wright State University, Dayton, Ohio
B.S., M.S., Ph.D.
(University of Kaiserslautern)

Dr. Abdurrahman Arslanyilmaz
Computer Science & Information Systems Department
Youngstown State University
Ph.D., Texas A&M University
University of Missouri, Columbia
Gazi University, Turkey

Dr. Xiaohong He
Professor of International Business
University of Quinnipiac
BS, Jilin Institute of Technology; MA, MS, PhD,. (University of Texas-Dallas)

Burcin Becerik-Gerber
University of Southern California
Ph.D. in Civil Engineering
DDes from Harvard University
M.S. from University of California, Berkeley & Istanbul University

Dr. Bart Lambrecht
Director of Research in Accounting and FinanceProfessor of Finance
Lancaster University Management School
BA (Antwerp); MPhil, MA, PhD (Cambridge)

Dr. Carlos García Pont
Associate Professor of Marketing
IESE Business School, University of Navarra
Doctor of Philosophy (Management), Massachusetts Institute of Technology (MIT)
Master in Business Administration, IESE, University of Navarra
Degree in Industrial Engineering, Universitat Politècnica de Catalunya

Dr. Fotini Labropulu
Mathematics - Luther College
University of ReginaPh.D., M.Sc. in Mathematics
B.A. (Honors) in Mathematics
University of Windso

Dr. Lynn Lim
Reader in Business and Marketing
Roehampton University, London
BCom, PGDip, MBA (Distinction), PhD, FHEA

Dr. Mihaly Mezei
ASSOCIATE PROFESSOR
Department of Structural and Chemical Biology, Mount Sinai School of Medical Center
Ph.D., Etvs Lornd University
Postdoctoral Training,
New York University

Dr. Söhnke M. Bartram
Department of Accounting and FinanceLancaster University Management SchoolPh.D. (WHU Koblenz)
MBA/BBA (University of Saarbrücken)

Dr. Miguel Angel Ariño
Professor of Decision Sciences
IESE Business School
Barcelona, Spain (Universidad de Navarra)
CEIBS (China Europe International Business School).
Beijing, Shanghai and Shenzhen
Ph.D. in Mathematics
University of Barcelona
BA in Mathematics (Licenciatura)
University of Barcelona

Philip G. Moscoso
Technology and Operations Management
IESE Business School, University of Navarra
Ph.D in Industrial Engineering and Management, ETH Zurich
M.Sc. in Chemical Engineering, ETH Zurich

Dr. Sanjay Dixit, M.D.
Director, EP Laboratories, Philadelphia VA Medical Center
Cardiovascular Medicine - Cardiac Arrhythmia
Univ of Penn School of Medicine

Dr. Han-Xiang Deng
MD., Ph.D
Associate Professor and Research
Department Division of Neuromuscular Medicine
Davee Department of Neurology and Clinical NeuroscienceNorthwestern University
Feinberg School of Medicine

Dr. Pina C. Sanelli
Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic Radiology
M.D., State University of New York at Buffalo,School of Medicine and Biomedical Sciences

Dr. Roberto Sanchez
Associate Professor
Department of Structural and Chemical Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University

Dr. Wen-Yih Sun
Professor of Earth and Atmospheric SciencesPurdue University Director
National Center for Typhoon and Flooding Research, Taiwan
University Chair Professor
Department of Atmospheric Sciences, National Central University, Chung-Li, TaiwanUniversity Chair Professor
Institute of Environmental Engineering, National Chiao Tung University, Hsin-chu, Taiwan.Ph.D., MS The University of Chicago, Geophysical Sciences
BS National Taiwan University, Atmospheric Sciences
Associate Professor of Radiology

Dr. Michael R. Rudnick
M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and Hypertension Division (PMC)
Penn Medicine, University of Pennsylvania
Presbyterian Medical Center, Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of Internal Medicine

Dr. Bassey Benjamin Esu
B.Sc. Marketing; MBA Marketing; Ph.D Marketing
Lecturer, Department of Marketing, University of Calabar
Tourism Consultant, Cross River State Tourism Development Department
Co-ordinator , Sustainable Tourism Initiative, Calabar, Nigeria

Dr. Aziz M. Barbar, Ph.D.
IEEE Senior Member
Chairperson, Department of Computer Science
AUST - American University of Science & Technology
Alfred Naccash Avenue – Ashrafieh

President Editor (HON.)

Dr. George Perry, (Neuroscientist)
Dean and Professor, College of Sciences
Denham Harman Research Award (American Aging Association)
ISI Highly Cited Researcher, Iberoamerican Molecular Biology Organization
AAAS Fellow, Correspondent Member of Spanish Royal Academy of Sciences
University of Texas at San Antonio
Postdoctoral Fellow (Department of Cell Biology)
Baylor College of Medicine
Houston, Texas, United States

Chief Author (HON.)

Dr. R.K. Dixit
M.Sc., Ph.D., FICCT
Chief Author, India
Email: authorind@computerresearch.org

Dean & Editor-in-Chief (HON.)

Vivek Dubey(HON.)
MS (Industrial Engineering),
MS (Mechanical Engineering)
University of Wisconsin, FICCT
Editor-in-Chief, USA
editorusa@computerresearch.org

Sangita Dixit
M.Sc., FICCT
Dean & Chancellor (Asia Pacific)
deanind@computerresearch.org

Suyash Dixit
(B.E., Computer Science Engineering), FICCTT
President, Web Administration and
Development , CEO at IOSRD
COO at GAOR & OSS

Er. Suyog Dixit
(M. Tech), BE (HONS. in CSE), FICCT
SAP Certified Consultant
CEO at IOSRD, GAOR & OSS
Technical Dean, Global Journals Inc. (US)
Website: www.suyogdixit.com
Email:suyog@suyogdixit.com

Pritesh Rajvaidya
(MS) Computer Science Department
California State University
BE (Computer Science), FICCT
Technical Dean, USA
Email: pritesh@computerresearch.org

Luis Galárraga
J!Research Project Leader
Saarbrücken, Germany

Contents of the Volume

i. Copyright Notice
ii. Editorial Board Members
iii. Chief Author and Dean
iv. Table of Contents
v. From the Chief Editor's Desk
vi. Research and Review Papers

1. Transition of Morphology between Nanosheet and Nanosphere of Ternary Comb Copolymers with Hydrophobic Carbazole Ring Induced by Changes in Surface Pressure. *1-5*
2. Analysis by GC-MS of an Aza-Michael Reaction Catalyzed by CALB on an Orbital Shaker and under Microwave Irradiation. *7-21*
3. Water Quality of River Tungabhadra Due to the Discharge of Industrial Effluent at Harihar, District Davanagere, Karnataka State, India. *23-31*
4. Synthesis, Characterization and Antimicrobial Activity of Azetidin-2-One based Phenyl Sulfonyl Pyrazoline Derivatives. *33-38*
5. Elimination of Methylene Blue Dye with Natural Adsorbent « Banana Peels Powder ». *39-44*

vii. Auxiliary Memberships
viii. Process of Submission of Research Paper
ix. Preferred Author Guidelines
x. Index

Global Journal of Science Frontier Research: B
Chemistry
Volume 14 Issue 1 Version 1.0 Year 2014
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Transition of Morphology between Nanosheet and Nanosphere of Ternary Comb Copolymers with Hydrophobic Carbazole Ring Induced by Changes in Surface Pressure

By Yohei Kaneko, Takahiro Kikkawa & Atsuhiro Fujimori
Saitama University, Japan

Abstract- Transition behavior from monolayer on the water surface to single particle layer of ternary comb copolymer containing carbazole ring have investigated by surface pressure-area isotherm, atomic force microscopy, X-ray diffraction. This polymer monolayer is able to form the "polymer nanosheet" with amorphous layers and strong interaction between main-chains fabricated by a Langmuir-Blodgett (LB) technique. Further, single particle layer is able to form the multi-particle layered organization of "polymer nanosphere" by LB method. This ternary comb copolymer synthesized by radical copolymerization with hydrogenated and fluorinated long-chain vinyl compounds. Poly(N-vinylcarbazole) homopolymer formed single particle layer on the water surface after spreading, immediately. Spontaneous "particulation" behavior is accelerated by incorporation of carbazole units in ternary copolymer films. Monolayer of these copolymers on the water surface formed below 10 mN m–1, and this monolayer transited to single particle layer above 15 mN m–1.

Keywords: *organized molecular films, ternary comb copolymer, fluorocarbon, hydrocarbon, n-vinyl carbazole.*

GJSFR-B Classification : *FOR Code: 039999*



Strictly as per the compliance and regulations of :

© 2014. Yohei Kaneko, Takahiro Kikkawa & Atsuhiro Fujimori. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Transition of Morphology between Nanosheet and Nanosphere of Ternary Comb Copolymers with Hydrophobic Carbazole Ring Induced by Changes in Surface Pressure

Yohei Kaneko [α], Takahiro Kikkawa [σ] & Atsuhiro Fujimori [ρ]

Abstract- **Transition behavior from monolayer on the water surface to single particle layer of ternary comb copolymer containing carbazole ring have investigated by surface pressure-area isotherm, atomic force microscopy, X-ray diffraction. This polymer monolayer is able to form the "polymer nanosheet" with amorphous layers and strong interaction between main-chains fabricated by a Langmuir-Blodgett (LB) technique. Further, single particle layer is able to form the multi-particle layered organization of "polymer nanosphere" by LB method. This ternary comb copolymer synthesized by radical copolymerization with hydrogenated and fluorinated long-chain vinyl compounds. Poly(N-vinylcarbazole) homopolymer formed single particle layer on the water surface after spreading, immediately. Spontaneous "particulation" behavior is accelerated by incorporation of carbazole units in ternary copolymer films. Monolayer of these copolymers on the water surface formed below 10 mN m–1, and this monolayer transited to single particle layer above 15 mN m–1.**

Keywords: organized molecular films, ternary comb copolymer, fluorocarbon, hydrocarbon, n-vinyl carbazole.

I. Introduction

Organized molecular films1 fabricated using functional polymers with precisely controlled structures can be developed as candidates for biomimetic models2 and molecular electronic devices,3 which have attracted considerable interest in fundamental science, as well as for their potential applications. Generally, it is well known that crystalline polymers construct hierarchical structures from lamellae on the nanometer scale to spherulite on the mesoscopic scale. 4 Crystals in these crystalline polymers are generally formed by the folding of the main-chain, although there main-chain, although there are two types of structural characteristics for side-chain crystalline ternary comb polymers. It is known that ternary comb polymers, which have long n-alkyl side-chains with crystallization ability attached to an amorphous main-chain, are packed into a layer structure along the c-axis and sub-cell of the side-chains in the ab-plane.5 This "sub-cell" structure6 of the side-chain is formed as a two-dimensional lattice, and the "layer structure" along the c-axis often reflects the long spacing between the main-chains in the accumulated double layer structures. With regard to the industrial application of these type of ternary comb polymers, poly(octadecyl)-based materials that use co-crystallization with another hydrocarbon to incorporate the material into the same crystalline lattice have found numerous applications such as pour-point depressants for lubricating oils or fuels, rheological modifiers, additives in petroleum products, and smart gels.7

On the other hand, many researchers have attempted to utilize poly-N-vinylcarbazole (Poly-NVCz) as organic light emitting diodes, organic transistors, and host compounds for organic electroluminescence8 in recent years, because it is well known that Poly-NVCz exhibits properties identical to those of organic semiconductors.9 Further, it is expected that Poly-NVCz can be used to realize a hologram memory material because of the high refractive index (greater than 1.68) of its monomer.10

As mentioned above, it is essential to control the molecular orientation and the arrangement at the monomolecular level in order to drive the functionality of these organic molecular devices in an efficient manner. However, in many cases, it is difficult to control the molecular arrangement and the homopolymer packing because Poly-NVCz itself easily forms an amorphous polymer.11

In previous study, we have investigated the control of solid-state structures and the formation of organized films for the newly synthesized comb copolymers containing hydrogenated and fluorinated side-chains.12 These comb polymers form a side-chain crystal in the bulk state by van der Waals interaction between the side-chains and the extremely stable condensed monolayers at the air/water interface. If bulky and flat functional groups are incorporated in the main-chains of side-chain crystalline polymers, it will be possible to control the arrangement of bulky groups at

Authors α σ ρ: Graduate School of Science and Engineering, Faculty of Engineering, Saitama University, 255 Shimo-okubo, Sakura-ku, Saitama, Japan. e-mail: fujimori@fms.saitama-u.ac.jp

© 2014 Global Journals Inc. (US)

the molecular level, regardless of the low crystallinity of the polymers.

Recently, the fine structures in the solid-state of the synthesized ternary comb polymers containing N-vinylcarbazole (NVCz) in the main-chains obtained by co-polymerization with both hydrogenated and fluorinated long-chain compounds were investigated by wide-angle X-ray diffraction (XRD). Further, the molecular arrangements of the organized molecular films in ternary comb copolymers containing NVCz units were estimated by in-plane XRD.13

In this study, transition behavior from monolayer on the water surface to single particle layer of ternary comb copolymer containing carbazole ring have investigated by surface pressure-area (π-A) isotherm, atomic force microscopy (AFM), out-of-plane XRD.

II. Experimental

a) Materials

The ternary comb copolymers used in this study were obtained by the copolymerization of NVCz with octadecyl acrylate (OA) and 2-(perfluorodecyl)ethyl acrylate (FF10EA) at various monomer ratios. Copolymerizations were carried out in an acetone solution at 50 °C for 48 h using azobisisobutyronitrile (AIBN) as an initiator. These monomers and the initiator were purchased from Tokyo Kasei Co. Ltd. and Daikin Fine Chemicals Co. Ltd., and were used without further purification. The precipitated polymers were washed with acetone until they were free of monomers. The ternary comb copolymer compositions were determined by 1H NMR (Nihon Densi Co. Ltd. EX270 NMR) spectroscopy. The tacticity of fluorinated homopolymer obtained by 1H NMR analysis according to the reference was found to be almost syndiotactic (Diad: 58 %). In this study, we synthesized ternary copolymers with two series of molecular weights estimated to be approximately Mw = 4.43 × 104 and 4.93 × 103 (Mw/Mn ≈ 2.14 and 1.17 respectively) on the basis of GPC (JASCO-860-CO) measurements.

According to the theory of the Q-e scheme proposed by T. Alfrey and C.C. Price,14 these ternary comb polymers form alternating copolymers in the view of relation of NVCz and long-chain acrylates. In this case, the e values of NVCz, OA and FF10EA are –1.40, +1.12 and +0.66 respectively. Thus, it appears that the NVCz : long-chain acrylates copolymers almost form ideal alternating copolymers, especially for the 2:1:1 monomer ratio.

b) Formation of copolymer monolayers on water surface and estimation of molecular arrangement in the films

The monolayers of the ternary comb copolymer with NVCz were spread from the chloroform or chloroform/trifluoroacetic acid = 90/10 (v/v) mixed solutions (about 10–4 M) onto the distilled water (about 18 MΩ•cm). The π–A isotherms were measured by a USI-3-22 film balance (USI Co. Ltd.) at 15 °C. These monolayers were transferred onto glass (XRD sample), quartz (UV-vis and fluorescence spectroscopy), and mica (AFM) substrates by the Langmuir-Blodgett (LB) method to obtain alternating Y-type film at 15 °C.

The long spacing between the layer structures along the c-axis estimated by an out-of-plane X-ray diffractometer (Rigaku, Rint-Ultima III, CuKα radiation, 40 kV, 40 mA) equipped with a graphite monochromator.

III. Results and Discussion

Figure 1 shows π-A isotherm of "single particle layer" on the water surface and corresponding AFM image of single particle layer on solid substrate of Poly-NVCz. This homopolymer formed single particle layer on the water surface after spreading, immediately. This




Fig.1 : π-A isotherm of single particle layer on the water surface and AFM images of single particle layer on solid of PNVCz homopolymer, and out-of-plane XRD of multi-particle layers of Poly-NVCz.

© 2014 Global Journals Inc. (US)

figure also shows out-of-plane XRD of multi-particle layers of Poly-NVCz. From the result of this measurement, it is found that inter-particle spacing along the c-axis correspond to 52 Å.

Figure 2 shows π-A isotherm of "monolayer" on the water surface and corresponding AFM image of monolayer on solid substrate of NVCz:FF10EA:OA = 2:1:1 copolymer. This figure also shows out-of-plane XRD of multilayers of this copolymer. This profile clearly indicates higher order reflection of (00l) plane. Therefore, it is found that this LB multilayer formed highly ordered layer structure.




Fig. 2 : π-A isotherm of monolayer on the water surface and AFM images of monolayer on solid of VCz:FF_{10}EA:OA = 2:1:1 copolymer, and out-of-plane XRD of multilayers of this copolymer

Figure 3 showsπ -A isotherm of monolayer on the water surface of NVCz:FF10EA:OA = 3:1:1, 4:1:1, and 6:1:1 copolymers. All isotherms indicate an existence of plateau region based on the two-dimensional phase transition.




Fig. 3 : π-A isotherm of monolayer on the water surface of NVCz: FF_{10}EA:OA = 3:1:1, 4:1:1, and 6:1:1 copolymers

Figure 4 shows AFM images of Z-type monolayer and single particle layer on the water surface NVCz:FF10EA:OA = 3:1:1, 4:1:1, and 6:1:1 copolymers transferred at before and after transition. This figure indicates monolayer of these copolymers on the water surface formed below 10 mN m–1, and this monolayer transited to single particle layer above 15 mN m–1. This figure also means that spontaneous "particulation" behavior is accelerated by incorporation of carbazole units in ternary copolymer films.




Fig. 4 : AFM images of Z-type monolayers and single particle layers on the water surface of NVCz: FF_{10}EA: OA = 3:1:1, 4:1:1, and 6:1:1 copolymers transferred at before (5 mN m^{-1}, lower column) and after (35 mN m^{-1}, upper column) transition.

Figure 5 shows out-of-plane XRD profiles of LB multilayers of NVCz:FF10EA:OA = 4:1:1 copolymer transferred at 35 and 5 mN m–1. It is found that both organizations commonly formed ordered layer structure along the c-axis.

© 2014 Global Journals Inc. (US)

Figure 6 shows schematic illustration of polymer nanosheet and polymer nanosphere layered




Fig. 5 : Out-of-plane XRD profiles of LB multilayers of NVCz:FF_{10}EA:OA = 4:1:1 copolymer transferred at (a) 5 mN m^{-1} and (b) 35 mN m^{-1}.

organization, and their formation mechanism. It seems that polymer particles are formed by piled up molecular chain, like a bilayer structure in high surface pressure region.




Fig. 6 : Schematic illustration of polymer nanosheet and polymer nanosphere layered organization, and their formation mechanism

Figure 7 shows UV-vis and fluorescence spectra of Poly-NVCz and NVCz:FF10EA:OA = 4:1:1 copolymer in solution. Further, fig. 8 shows ones of




Fig. 7 : UV-vis and fluorescence spectra of Poly-NVCz and NVCz:FF_{10}EA:OA = 4:1:1 copolymer in solution




Fig. 8 : UV-vis and fluorescence spectra of multi-particle layered organization of Poly-NVCz and NVCz: FF_{10}EA:OA = 4:1:1 copolymer.

© 2014 Global Journals Inc. (US)

multi-particle layered organization of same materials.

The "polymer nanosphere" of these ternary comb copolymers showed the absorption band around 250 nm and emitted fluorescence at 400 nm. The emitted band in fluorescence spectrum showed red-shift to the one of their solution. It is speculated that this result means formation of aggregated structure by π-π stacking of NVCz rings.[15-17]

IV. Conclusion

Formation behavior of single particle layer of ternary comb copolymer containing NVCz have investigated. This copolymer monolayer is able to form the "polymer nanosheet" fabricated by a LB technique. Further, single particle layer is able to form the multi-particle layered organization of "polymer nanosphere". This ternary comb copolymer synthesized by radical copolymerization. Spontaneous particle formation is attained by incorporation of hydrophobic carbazole. Monolayer of these copolymers on the water surface transited to single particle layer above 15 mN m–1.

References Références Referencias

1. Gaines, Jr. G. L., "Insoluble Monolayers at Liquid Gas Interfaces", Wiley, New York, 1966.
2. Kuhn, H.; Möbius, D.; Bucher, H., Spectroscopy of Monolayer Assemblies, in "Physical Methods of Chemistry" (Weissberger, A., Rossiter, B. W., Eds.), Vol. 1, Part IIIB, pp. 577-702, Wiley, New York, 1972.
3. Ulman, A., "Ultrathin Organic Films", Academic Press, London, 1991.
4. Keller, A., Phil. Mag. 1957, 2, 1171.
5. Neugebauer, D.; Theis, M.; Pakula, T.; Wegner, G.; Matyjaszewski, K., Macromolecules 2006, 39, 584.
6. Vand, A., Acta Cryst. 1951, 4, 465.
7. Baskar, G.; Ramya, S.; Mandal, A., Colloid Polym. Sci. 2002, 280, 886.
8. Pearson, J. M.; Stolka, M., "Poly(N-vinyl carbazole)" in Polymer monograph, vol. 6, New York: Gordon and Breach, 1981.
9. Choudhury, K. R.; Samoc, M.; Patra, A.; Prasad, P. N., J. Phys. Chem. B 2004, 108, 1556.
10. Laguna, M. T.; Gallego, R. J.; Mendicuti, F.; Saiz, E.; Tarazona, M. P., Macromolecules 2006, 35, 7782.
11. Maertens, C.; Dubois, P.; Jerome, R.; Blanche, P.A.; Lemaire, P.C., J. Polym. Sci., Part B, Polym. Phys. 2000, 38(1), 205.
12. Masuya, R.; Ninomiya, N.; Fujimori, A.; Nakahara, H.; Masuko, T., J. Polym. Sci., Part B, Polym. Phys. 2006, 44, 416.
13. Fujimori, A.; Hoshizawa, H.; Kobayashi, S.; Sato, N.; Kanai, K.; Ouchi, Y., J. Phys. Chem. B 2010, 114, 2100.
14. Alfrey, T.; Price, C. C., J. Polym. Sci. 1947, 2, 101.
15. Fujimori, A.; Kaneko, Y.; Kikkawa, T. Chiba, S.; Shibasaki, Y., J. Colloid Interf. Sci. 2014, 418, 338.
16. Kaneko, Y.; Fujimori, A., Chem. Lett. 2012, 41, 1183.
17. Kaneko, Y.; Fujimori, A., Trans. Mater. Res. Soc. Jpn., 2012, 37, 291.

© 2014 Global Journals Inc. (US)

This page is intentionally left blank

© 2014 Global Journals Inc. (US)

Global Journal of Science Frontier Research: B
Chemistry
Volume 14 Issue 1 Version 1.0 Year 2014
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Analysis by GC-MS of an Aza-Michael Reaction Catalyzed by CALB on an Orbital Shaker and under Microwave Irradiation

By Sandra S. Ribeiro, Marciana P. Uliana, Timothy J. Brocksom & André L. M. Porto

Universidade De São Paulo, Brazil

Abstract- In this study, aza-Michael reactions between 1-phenylmethanamine and *α,β*-unsaturated cyclohexenones were investigated, using lipase from *Candida antarctica.* The reactions were performed in various organic solvents (CH_2Cl_2, hexane, MeOH, toluene, THF) under mild conditions, with orbital shaking and microwave irradiation. The reactions showed good results in the presence of CALB, yielding the Michael adducts and imines. The reaction products were analyzed by GC-MS and in some cases it was found that the reverse aza-Michael reaction had occurred.

Keywords: *biocatalysis; amines; aza-Michael addition; enones; imines.*

GJSFR-B Classification : *FOR Code: 100505, 250106*



Strictly as per the compliance and regulations of :

© 2014 Sandra S. Ribeiro, Marciana P. Uliana, Timothy J. Brocksom & André L. M. Porto. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/ licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Analysis by GC-MS of an Aza-Michael Reaction Catalyzed by CALB on an Orbital Shaker and under Microwave Irradiation

Sandra S. Ribeiro [α], Marciana P. Uliana [σ], Timothy J. Brocksom [ρ] & André L. M. Porto [ω]

Abstract- **In this study, aza-Michael reactions between 1-phenylmethanamine and *α,β*-unsaturated cyclohexenones were investigated, using lipase from *Candida antarctica*. The reactions were performed in various organic solvents (CH_2Cl_2, hexane, MeOH, toluene, THF) under mild conditions, with orbital shaking and microwave irradiation. The reactions showed good results in the presence of CALB, yielding the Michael adducts and imines. The reaction products were analyzed by GC-MS and in some cases it was found that the reverse aza-Michael reaction had occurred.**

Keywords : biocatalysis; amines; aza-Michael addition; enones; imines.

I. Introduction

Chiral amines are an important class of organic compounds, on account of their utility in the preparation of pharmaceutical and industrial products of considerable interest [1,2]. These compounds are intermediates in the synthesis of a large number of organic compounds and are widely used to prepare derivatives of natural products, antibiotics, chiral auxiliaries and lactams. Owing to their vast range of applications, the modification and improvement of techniques for synthesizing *β*-amino carbonyl compounds has been a research objective in recent years [2, 3].

The conjugate addition of the nucleophilic nitrogen in compounds such as amines to *α,β*-unsaturated carbonyl or nitrile compounds, namely the aza-Michael reaction, constitutes a key reaction for the construction of C-N bonds in the preparation of *β*-amino esters and nitriles [4]. This reaction is highly versatile as it can occur between various *N*-nucleophiles (aliphatic and aromatic amines, amides, carbamates and azides) and Michael acceptors (enones, acrylates, unsaturated nitriles, amides, sulfones, phosphonates, trifluoromethylalkenes and nitroalkenes) [5].

Recently, an efficient enzymatic protocol was reported for the synthesis of *β*-amino esters via aza-Michael addition of primary and secondary amines to acrylates; CALB (*Candida antarctica* lipase B) was the biocatalyst and the corresponding *β*-amino esters were produced in good yields [6]. Green chemistry, in which harmful organic solvents and extreme conditions are avoided, is growing in importance; furthermore, the catalyst can be recycled. An efficient and simple protocol for aza-Michael addition of aliphatic and aromatic amines to electron-deficient alkenes has been established, using TMG-based ionic liquids as catalyst under solvent-free conditions [7]. Solvent-free aza-Michael reactions between a variety of amines and *α,β*-unsaturated carbonyl compounds under microwave irradiation, catalyzed by perchloric acid impregnated on silica gel ($HClO_4$=SiO_2), produced the corresponding adducts [8]. Another method has been developed for the aza-Michael addition of acrylonitrile to 2-aryl-oxymethylbenzimidazole derivatives, in the presence of anhydrous potassium carbonate under microwave irradiation, to synthesize a novel series of 1-cyanoethyl-2-aryloxymethylbenzimidazole derivatives [9]. The present paper summarizes the results of our investigation into the conjugate addition of nitrogen-containing nucleophiles to electron-deficient ketones, in which environmentally-friendly enzymatic catalysis was carried out in an orbital shaker and under microwave irradiation, yielding aza-Michael imine adducts and reverse aza-Michael compounds.

II. Materials and Methods

a) General procedure

The Michael addition reactions were carried out on a Tecnal TE-421 orbital shaker. The performance of the enzymatic reactions was measured by analyzing the products in a Shimadzu GC 2010 gas chromatograph equipped with an AOC 20i auto injector, a flame ionization detector (FID) and a J&W Scientific DB-5 column (30 m x 0.25 mm x 0.25 μm). The conditions employed in the gas chromatograph were as follows: carrier gas: nitrogen (60 kPa); injector temperature: 250 °C; injector split ratio: 1:20; detector temperature: 300 °C; initial oven temperature: 80 °C for 2 min; final oven temperature: 250 °C for 3 min; heating rate: 10 °C min^{-1}, and total time of analysis: 22.0 min (Supplementary Information – SI). Gas chromatography–mass spectrometry (GC-MS): a Shimadzu GC2010 Plus

Authors ω α: Instituto de Química de São Carlos, Universidade de São Paulo, Av. João Dagnone, 1100, Ed. Química Ambiental, J. Santa Angelina, 13563-120, São Carlos, SP, Brazil.
e-mail: almporto@iqsc.usp.br
Authors σ ρ: Universidade Federal de São Carlos, Rod. Washington Luís, km 235, 13565-905, São Carlos, SP, Brazil.

© 2014 Global Journals Inc. (US)

gas-chromatography system coupled to a mass-selective detector (Shimadzu MS2010plus) was used in the electron ionisation (EI) mode. The GC-MS oven was fitted with a DB-5 fused silica column (J&W Scientific 30 m x 0.25 mm x 0.25 μm). The oven temperature was programmed from 50 °C to 270 °C at a heating rate of 10 °C/min, and a total time of analysis: 32.0 min. The injector and detector temperatures were maintained at 300 °C; injector split ratio was 1:20 and helium was used as the carrier gas at a pressure of 60 kPa. The compounds were identified from the fragmentation ions (70 eV) detected in the GC-MS spectra (Supplementary Information – SI). The structures were confirmed after comparison with Mass Spectral Database (NIST 5.0) and by co-injection of authentic standards. Reactions were purified by column chromatography on silica gel (230–400 mesh) eluted with mixtures of *n*-hexane and EtOAc (8:2). Column effluents were monitored by TLC, using aluminium-backed pre-coated silica gel.

b) Enzyme and chemicals

Novozym 435® (component B of the lipase from *Candida antarctica* immobilized on macroporous poly-acrylate resin, with 10,000 propyl laurate units per gram), was a gift from Novo Nordisk (Curitiba-Paraná, Brazil). Solvents (EtOAc, CH_2Cl_2, *n*-hexane, MeOH, toluene, THF) were commercially available and of analytical grade. Cyclohex-2-en-1-one **1**, 1-phenylmethanamine **2** and 3-methylcyclohex-2-en-1-one **5** were purchased from Sigma-Aldrich and 2,5-dimethyl-*para*-benzoquinone **8** was synthesized from 2,5-dimethyl-phenol [10, 11].

c) Lipase-catalyzed aza-Michael reactions under conventional conditions (orbital shaker)

CALB lipase (80 mg), the unsaturated ketone [2-cyclohexen-1-one **1** (40 μL, 0.41 mmol), 3-methylcyclohex-2-en-1-one **5** (40 μL, 0.35 mmol) or 2,5-dimethyl-*para*-benzoquinone **8** (40 μL, 0.33 mmol)] and the amine [1-phenylmethanamine **2** (40 μL, 0.36 mmol)] were separately added to 10 mL of the organic solvent (*n*-hexane, toluene, dichloromethane, tetrahydrofuran or methyl alcohol) in 50 mL Erlenmeyer flasks. The flasks were sealed with a rubber stopper and the reaction mixture was shaken in an orbital shaker at 33 °C and 133 rpm. The reaction progress was monitored by collecting samples (0.5 mL) at intervals, between 2 and 48 hours from the start, and analyzed by gas chromatography GC-FID and GC-MS. The enzymatic aza-Michael reactions were carried out in triplicate.


d) Lipase-catalyzed aza-Michael with microwave irradiation

The microwave irradiation (MW) experiments were performed with a Discover System from CEM Corporation. Organic solvents (10 mL, EtOAc, CH_2Cl_2, *n*-hexane, MeOH, toluene, THF), primary amine **2** (40 μL, 0.36 mmol), Michael acceptors **1** (40 μL, 0.41 mmol), **5** (40 μL, 0.35 mmol) and **8** (40 μL, 0.33 mmol) and CALB (80 mg) were separately added to a 50 mL flat bottom flask. The whole reaction mixture was placed in the microwave oven at a frequency 2.45 GHz and irradiated at 33, 40, 50 and 60 °C, power output of about 70 W. The reaction progress was monitored by collecting samples (0.1 mL) and analyzed by gas chromatography with a J & W Scientific DB-5 (30 m x 0.25 mm x 0.25 μm). After 3 hours of reaction, the flask was removed and the immobilized lipase was filtered off. The organic solvent was evaporated under reduced pressure. The residue was purified by column chromatography on silica gel with 8:2 hexane and ethyl acetate as eluent. The reactions were carried out in triplicate and the results are presented and discussed next.

III. Results and Discussion

The lipase from *Candida antarctica* (CALB) is often used to promote asymmetric reactions via transesterification of racemic alcohols and amines by chemo-enzymatic resolution [12-15]. The enzymes used to catalyze unusual reactions, commonly named promiscuous enzymes, such as the Michael addition reaction catalyzed by lipases, are being investigated with interest [6, 16-17].

The aim of this study was to carry out Michael additions with the immobilized lipase from *C. antarctica*, on account of its proven catalytic efficiency and thermal stability, with high activity at 70-90 °C [18]. Our study began with cyclohex-2-enone **1**, due to the absence of steric hindrance in the 1,4-addition, and the achiral primary amine, 1-phenylmethanamine **2**. Under these conditions, the reaction can favor two products, the imine *N*-(cyclohex-2-en-1-ylidene)-1-phenylmethanaminne **3** and the chiral aza-Michael adduct 3-(benzylamino) cyclohexanone **4** (Scheme 1).



Scheme 1 : Aza-Michael reactions using CALB on an orbital shaker and microwave irradiation

© 2014 Global Journals Inc. (US)

Initially, this reaction was tested in five organic solvents (*n*-hexane, dichloromethane, methanol, tetrahydrofuran and toluene). It was also performed in the absence of CALB, since primary amine **2** is a strong nucleophile and *α,β*-unsaturated ketone **1** is a good Michael acceptor, which could favor the spontaneous formation of products **3** and **4**.

Under orbital shaking, in the presence of CALB, this reaction gave the aza-Michael adduct **4** at low conversion rates, varying with the solvent used (c 9-37%), and imine **3** at conversions of 5-23%, in 48 h. The non-enzymatic reaction gave similar results for adduct **4** (c 3-29%) and imine **3** (4-34%) in 48 h. The reaction in the presence of CALB and *n*-hexane produced adduct **4** at 25% conversion and imine **3** at 23% in 48 h, while under the same conditions without CALB lower yields of both adduct **4** (c 10%) and imine **3** (c 9%) in 48 h were observed. In this case, there is a visible influence of lipase in the formation of the aza-Michael adduct **4** (SI-1). In methanol, there was good conversion to the adduct **4** (c 38%) and imine **3** (c 14%) in 48 h. In the absence of CALB, a decrease occurred in the conversion of adduct **4** (c 29%) and an increase in conversion to imine **3** (c 35%) in 48 h (SI-2).

It is reported in the literature that CALB exhibits good catalytic activity in hydrophobic solvents [12, 14, 19]. In our studies the reactions in *n*-hexane and methanol were not efficient. Factors such as the polarity of the substrates (ketone **1** and amine **2**) may have influenced these results, and Michael addition reactions may be occurring spontaneously and reversibly, yielding unstable products, without being promoted by CALB [17].

In dichloromethane, with CALB, the reaction showed low conversion to adduct **4** (c 10%) and imine **3** (c 6%) after 48 h. In the absence of lipase we observed similar values to those obtained enzymatically for adduct **4** (c 7%) and imine **3** (c 4%), at 48 h (SI-3).

In toluene, adduct **4** showed low conversion (c <14%), as did imine **3** (c 11%), after 48 h. In the non-enzymatic reaction, adduct **4** was obtained at a conversion of less than 3% and the imine at 2% in 48 h (SI-4). The choice of solvent depends strongly on the solubility of the catalyst and of the donor and acceptor in the Michael addition, as well as the ability of the solvent to prevent the occurrence of side reactions. For example, if the reactant or products are susceptible to alcoholysis (ester hydrolysis), transesterification, autocondensation, reverse Michael reactions, non-hydroxyl solvents are desirable [20].

Aiming to improve the conversion of aza-Michael addition in the presence of CALB, the reactions with ketone **1** and amine **2** were promoted, in methanol, THF, dichloromethane and *n*-hexane, by microwave irradiation (40 °C, 70 W). Under these conditions, the aza-Michael adduct **4** showed 0.3-42% conversion and imine **3** of 2-40% conversion, depending on the solvent used. The reaction in methanol gave the best result, the adduct **4** was produced at 42% conversion and imine **3** at 4%, in 3 h.

In *n*-hexane and dichloromethane, only imine **3** (c 12%) was formed in 3 h. In THF, there were low conversions of adduct **4** (0.3-6%) and imine **3** (1-3%) in 3 h. The reactions in dichloromethane and hexane, without CALB, yielded the adduct **4** and imine **3** in small amounts (c<10%) by microwave irradiation for 3 h.

The regioselectivity of the reactions depends on the conditions in which they are carried out, the type of catalyst, the solvent and the nucleophile [21]. In these studies, the reaction both in the presence and absence of CALB favored the formation of products via 1,2-addition and 1,4-addition with similar conversions.

Dhake et al. (2010) conducted aza-Michael additions between various primary and secondary amines and acrylates, in the presence of lipase, to obtain the corresponding *β*-amino esters. In this study, methyl acrylate and CALB in toluene showed the best results in the synthesis of the Michael adduct. It was observed that the activity of the enzyme increased with the temperature, up to 60 °C. However, at higher temperatures (> 80 °C), the yield of the reaction decreased to 75%, possibly due to inactivation of the enzyme [6].

All the reactions in this first part were monitored by GC-FID and GC-MS analysis, which showed the formation of the aza-Michael adduct **4** and imine **3** (Supplementary Information - SI). During the isolation of the adduct **4** by silica gel only the starting reactants, namely enone **1** and the amine **2**, were found. Thus, it was concluded that a reverse Michael reaction occurred during the purification process. The instability of these products was confirmed by the main fragments obtained in the mass spectra, which were compared with data from the Nist 5.0 library.


In the GC-MS chromatogram of the reaction, in *n*-hexane under orbital shaking for 4 days (SI-5), a signal with a retention time of 14.5 minutes was observed, which provided the spectrum of a molecular ion at *m/z* 185 $[M]^+$. The mass spectrum of imine **3** was comparable to the Nist 5.0 mass spectral database. The fragmentation of the molecular ion at *m/z* 185 provided the tropylium ion, at *m/z* 91, as the base peak. On the basis of these fragmentations it was possible to suggest the formation of imine **3** (SI-5 and SI-6).

Another signal in the chromatogram, with retention time 15.9 minutes, showed the molecular ion at *m/z* 203 that corresponds exactly to the aza-Michael adduct **4**. The mass spectrum was similar to that reported for adduct **4** in the Nist 5.0 mass spectral database, and is consistent with the literature [22, 23]. The fragmentation pattern of the molecular ion at *m/z* 203 again provided the tropylium ion (*m/z* 91) as a base peak, which confirmed the formation of adduct **4** (SI-5 and SI-6).

© 2014 Global Journals Inc. (US)

Thus, we conclude from the spectral evidence that it is possible to obtain the compounds **3** and **4** from enone **1** and amine **2**. However, due to the instability of the products, they were not isolated, but were characterized by mass spectrometry.

The next step was to evaluate the regioselectivity of the Michael reaction by using the substituted acceptor, 3-methylcyclohex-2-en-1-one **5** and the achiral donor, 1-phenylmethanamine **2**. The reactions were performed in the presence and absence of CALB in the orbital shaker, with hexane and methanol, and under microwave irradiation, with methanol, dichloromethane and hexane (Scheme 2).



Scheme 2 : Aza-Michael reactions using CALB on an orbital shaker and microwave irradiation

The reactions under orbital shaking were similar, with or without CALB, in methanol: the imine **7** was produced at 27-38% conversion in 24 h, whereas the adduct **6** was not detected under these conditions (SI-7).

In hexane and with CALB, adduct **6** was not observed after 24 h, but only imine **7** at 30% conversion (SI-8). In the absence of lipase, imine **7** was produced at less than 5%. In this case, CALB influnced greatly the product obtained by 1,2-addition. In these reactions, the methyl group at position 3 of enone **5** hinders the Michael 1,4-addition, probably by steric and electronic effects. The larger the substituent group on the ***α,β***-unsaturated ketone, the more unfavorable is the 1,4-addition [20].

Initially, good yields were expected for adduct **6** with lipase in hexane, because the enzyme acts very well in nonpolar solvents. However, ketone **5** and amine **2** have polar groups that may influence the low conversion to imine **7**.

The reactions under microwave irradiation were performed at 33 and 50 °C and 70 W. The enzymatic reaction in methanol and dichloromethane at 33 °C did not form adduct **6** and imine **7** showed conversion < 5%.


Yu et al. studied the effects of temperature on the CALB in the transesterification of (±)-2-octanol by microwave irradiation. In that study the temperature was assessed from 40 to 100 °C, and the activity of the enzyme increased with temperature up to 80 °C, while there was a fall in enzymatic activity at higher temperatures [24]. The elevated temperature in a microwave reactor increases the probability of collision between enzyme and substrate molecules, which can promote the formation of the enzyme-substrate complex and increase the rate of reaction and the final conversion [24].

Given the low conversion of reaction in methanol and dichloromethane, the temperature was raised to 50 °C and 70 W, resulting in a better performance of the reaction in methanol in the formation of the ***α,β***-unsaturated imine **7**, but with low conversion (c = 10%). The results of these experiments were similar in the enzymatic and non-enzymatic reactions (c<11%, 50 °C, 70 W), yielding only the imine **7** at low conversion. There are several factors that may influence the low performance of these reactions. Amine **2** may be interacting with other parts of the lipase, hampering the Michael addition under these conditions. Also these reactions are subject to the reverse aza-Michael pathway; thus, the instability of imine **7** was confirmed by the main fragments obtained in the mass spectra by GC-MS, compared with data from the Nist 5.0 library (SI-8).

Analysis of the chromatogram of the reaction performed in hexane by orbital shaking, showed the presence of a signal with a retention time of 15.9 minutes. Its mass spectrum showed the ion fragment at *m/z* 199, corresponding to imine **7** (SI-8 and SI-9). To confirm the formation of imine **7**, the reaction was performed in the orbital shaker with methanol. After the reaction was complete and the presence of imine **7** confirmed by GC-MS, 10 mL of distilled water was added and the reaction was maintained under shaking for one hour. After extraction, the reaction was analyzed by GC-MS and showed the reacting species **5** and **2**.

Finally, we carried out the reaction with primary amine **2** and 2,5-dimethyl-*para*-benzoquinone **8**. The experiments were conducted in hexane with CALB on an orbital shaker and under microwave irradiation for 96 h. After this time, two products with retention times of 15.0 minutes (c 43%) and 19.0 minutes (c 34%) were detected by GC-MS analysis (Scheme 3) (SI-10).

© 2014 Global Journals Inc. (US)



Scheme 3 : Aza-Michael reactions using CALB on an orbital shaker and under microwave irradiation

The reactions were conducted in hexane and microwave reactor at 60 °C (70 W) in the presence of CALB. After 5h, the reactions yielded two products with retention times of 15.0 minutes (c 21%) and 19.0 minutes (c 18%) with complete consumption of amine **2** by GC-MS analysis (SI-11).

Analysis of the mass spectrum of the compound with retention time 15.0 minutes showed the presence of a molecular ion with the base peak at *m/z* 225. There was an intense peak at *m/z* 148 (75%) corresponding to the loss of a fragment at *m/z* 77. These fragments are consistent with the structure of the compound 4-(benzylimino)-2,5-dimethylcyclohexa-2,5-dienone **9** (SI-11 and SI-12).

The compound with retention time 19.0 minutes provided a mass spectrum for a molecular ion at *m/z* 195 and base peak at *m/z* 91. Analysis of the mass spectra with reference to the Nist 5.0 library and in comparison with the literature [25] suggested the formation of *N*-benzylidene-1-phenylmethanamine **10** (SI-11 and SI-12).

The formation of **10** could occur by nucleophilic attack of amine **2** on imine **9**. The non-enzymatic reaction does not occur under microwave irradiation at 3 h. However, the reaction under microwave irradiation promoted the formation **10** in face of the reactivity of the species involved (SI-11 and SI-12).

Finally, to confirm the production of imines **9** and **10**, the reaction mixture was subjected to hydrolysis at 24 hours, then extracted with ethyl acetate and analyzed by GC-MS. After hydrolysis, products **9** and **10** were all converted to the starting reagents **2** and **8**, confirming the synthesis of unstable imines in aqueous medium (SI-13 and SI-14).

In the chromatogram of the hydrolysis reaction, no amine **2** was observed, but the corresponding *N*-benzylacetamide **11** with retention time 11.3 minutes was formed by acetylation of amine **2** during extraction with ethyl acetate. The compound was confirmed by synthesis of the authentic standard (SI-15).

IV. Conclusion

In this paper we present studies on the regioselectivity of aza-Michael addition reactions between amine **2**, and enones **1**, **5** and **8** under various experimental conditions, such as types of organic solvent, presence and absence of lipase from *Candida antarctica*, orbital shaking and microwave irradiation. All the reactions yielded products via the formation of 1,2- and 1,4-additions. Owing to the poor stability of the compounds formed, GC-MS analyzes were carried out and showed the formation of aza-Michael adducts and several imines. Reverse aza-Michael reactions and hydrolysis of imines allowed us to verify the formation of all the products. In some cases, it was also possible to observe the influence of lipase and microwave irradiation on the performance of the reactions. The compounds were confirmed by detailed analysis of their mass spectra and the corresponding ions formed.

V. Acknowledgements

SSR thanks FAPESP (2010/50529-0) for the scholarships. ALMP gratefully acknowledges CNPq and FAPESP for their financial support. TJB acknowledges the São Paulo Research Foundation for financial support (FAPESP; 2011/13993-2) and (2008/53507-7). The authors also wish to thank Novo Nordisk (Curitiba-Paraná, Brazil) for donation of the immobilized CALB enzyme (Novozym 435®). The English language was reviewed by Timothy Roberts, MSc., a native English speaker.

References Références Referencias


1. Gotor-Fernández, V.; Busto, E.; Gotor, V. *Candida antarctica* lipase B: An ideal biocatalyst for the preparation of nitrogenated organic compounds. Adv. Synth. Catal. 2006; 348: 797-812.
2. Yadav, J. S.; Reddy, B. V. S.; Basak, A. K.; Narsaiah, A. V. Aza-Michael reactions in ionic liquids. A facile synthesis of α-amino compounds. Chem. Lett. 2003; 32: 988-989.
3. Zhang, H.; Zhang, Y.; Liua, L.; Xua, H.; Wang, Y. $RuCl_3$ in poly(ethylene glycol): A highly efficient and recyclable catalyst for the conjugate addition of nitrogen and sulfur nucleophiles. Synthesis. 2005; 13: 2129–2136.
4. Trivedi, R.; Lalitha, P.; Roy, S. Vanadyl(IV) acetate: An efficient, reusable heterogenous catalyst for aza-Michael reaction under solvent-free conditions. Synth. Commun. 2008; 38: 3556-3566.
5. Rulev, A. Y. Aza-Michael reaction: achievements and prospects. Russian Chem. Rev. 2011; 80: 197-218.

© 2014 Global Journals Inc. (US)


6. Dhake, K. P.; Tambade, P. J.; Singhal, R. S.; Bhanage, B. M. Promiscuous *Candida antarctica* lipase B-catalyzed synthesis of *β*-amino esters via aza-Michael addition of amines to acrylates. Tetrahedron Lett. 2010; 51: 4455–4458.
7. Ying, A.; Zheng, M.; Xu, H.; Qiu, F.; Ge, C. Guanidine-based task-specific ionic liquids as catalysts for aza-Michael addition under solvent-free conditions. Res. Chem. Intermed. 2011; 37: 883–890.
8. Singh, S. P.; Kumar, T. V.; Chandrasekharam, M.; Giribabu, L.; Reddy, P. Y. Microwave-assisted, rapid, solvent-free aza-Michael reaction by perchloric acid impregnated on silica gel. Synth. Commun. 2009; 39: 3982–3989.
9. Wei, T-B.; Hua, M-T.; Shi, H-X.; Liu, Y.; Zhang, Y-M. Aza-Michael addition of acrylonitrile with 2-aryloxymethylbenzimidazole derivatives under microwave irradiation. J. Chem. Research. 2010; 34: 452-454.
10. Dockal, E. R.; Cass, Q. B.; Brocksom, T. J.; Brocksom, U.; Corrêa, A. G. Synth. A simple and efficient synthesis of thymoquinone and methyl *p*-benzoquinone. Commun. 1985; 15: 1033-1036.
11. Uliana, M. P.; Vieira, Y. W.; Donatoni, M. C.; Corrêa, A. G.; Brocksom, U.; Brocksom, T. J. J. Braz. Chem. Soc. Oxidation of mono-phenols to *para*-benzoquinones: a comparative study. 2008; 19: 1484-1489.
12. Ribeiro, S. S.; Raminelli, C.; Porto, A. L. M. Enzymatic resolution by CALB of organofluorine compounds under conventional condition and microwave irradiation. J. Fluorine Chem. 2013; 154: 53-59.
13. Ribeiro, S. S.; De Oliveira, J. R.; Porto, A. L. M. Lipase-catalyzed kinetic resolution of (±)-mandelonitrile under conventional condition and microwave irradiation. J. Braz. Chem. Soc. 2012; 23: 1395-1399.
14. Melgar, G. Z.; Wendler, E. P.; Santos, A. A. dos; Porto, A. L. M. First stereocontrolled acetylation of a hydroxypropargylpiperidone by lipase CALB. Tetrahedron: Asymmetry, 2010; 21: 2271–2274.
15. Ferraz, H. M. C.; Bianco, G. G.; Teixeira, C. C.; Andrade, L. H.; Porto, A. L. M. Enzymatic resolution of α-tetralols by CALB-catalyzed acetylation. Tetrahedron: Asymmetry 2007; 18: 1070–1076.
16. Souza, R. O. M. A. de; Matos, L. M. C.; Gonçalves, K. M.; Costa, I. C. R.; Nabics, I.; Leite, S. G. F.; Oestreicher, E. G.; Antunes, O. A. C. Michael additions of primary and secondary amines to acrylonitrile catalyzed by lipases. Tetrahedron Lett. 2009; 50: 2017-2018.
17. Priego, J.; Ortíz-Nava, C.; Carrillo-Morales, M.; López-Munguía, A. Escalate, J.; Castillo, E. Solvent engineering: an effective tool to direct chemoselectivity in a lipase-catalyzed Michael addition. Tetrahedron, 2009; 65: 536-539.
18. Bachu, P.; Gibson, J. S.; Sperry, J.; Brimble, M. A. The influence of microwave irradiation on lipase-catalyzed kinetic resolution of racemic secondary alcohols. Tetrahedron: Asymmetry 2007; 18: 1618-1624.
19. Rocha, L. C.; Rosset, I. G.; Luiz, R. F.; Raminelli, C.; Porto, A. L. M. Kinetic resolution of iodophenylethanols by *Candida antarctica* lipase and their application for the synthesis of chiral biphenyl compounds. Tetrahedron: Asymmetry 2010; 21: 926–929.
20. Mather, B. D.; Viswanathan, K.; Miller, K. M.; Long, T. E. Michael addition reactions in macromolecular design for emerging technologies. Prog. Polym. Sci. 2006; 31: 487-531.
21. Hoz, A. de L.; Días-Ortiz, A.; Gómez, M. V.; Mayoral, J. A.; Moreno, A.; Sánchez-Migallón, A. M.; Vazquez, E. Preparation of α- and β-substitutend alanine derivatives by α-amidoalkylation or Michael addition reactions under heterogeneous catalysis assisted by microwave irradiation. Tetrahedron 2001; 57: 5421-5428.
22. WSS: Spectral data were obtained from Wiley Subscription Services, Inc. (US).
23. Zhang, H.; Zhang, Y.; Liu, L.; Xu, H.; Wang, Y. $RuCl_3$ in poly(ethylene glycol): A highly efficient and recyclable catalyst for the conjugate addition of nitrogen and sulfur nucleophiles. Synthesis 2005; 13: 2129-2136.
24. Yu, D.; Wang, Z.; Chen, P.; Jin, L.; Vheng, Y.; Zhou, J.; Cao, S. Microwave-assisted resolution of (*R,S*)-2-octanol by enzymatic transesferification. J. Mol. Catal. B: Enzym. 2007; 48: 51-57.
25. Montalvo-González, R.; Ariza-Castolo, A. Molecular structure of di-aryl-aldimines by multinuclear magnetic resonance and X-ray diffraction. J. Mol. Structure. 2003; 655: 375-389.

© 2014 Global Journals Inc. (US)


Supplementary Information



SI-1 : Chromatograms obtained by GC-FID. Reaction of ketone **1** with amine **2** in hexane under orbital shaking (48 h, 33 °C, 133 rpm). (a) In the presence of lipase. (b) In the absence of lipase. Conditions of GC-FID analysis: DB-5 column (30 m x 0.25 mm x 0.25 µm) Ti = 80 °C, Tf = 250 °C, r = 10 °C/min, t = 22.0 min




SI- 2 : Chromatograms obtained by GC-FID. Reaction of ketone **1** with amine **2** in methanol under orbital shaking (48 h, 33 °C, 133 rpm). (b) In the presence of lipase. (b) In the absence of lipase. Conditions of GC-FID analysis: DB-5 column (30 m x 0.25 mm x 0.25 µm) Ti = 80 °C, Tf = 250 °C, r = 10 °C/min, t = 22.0 min




SI- 3 : Chromatograms obtained by GC-FID. Reaction of ketone **1** with amine **2** in dichloromethane under orbital shaking (48 h, 33 °C, 133 rpm). (a) In the presence of lipase. (b) In the absence of lipase. Conditions of GC-FID analysis: DB-5 column (30 m x 0.25 mm x 0.25 µm) Ti = 80 °C, Tf = 250 °C, r = 10 °C/min, t = 22.0 min

© 2014 Global Journals Inc. (US)



SI- 4 : Chromatograms obtained by GC-FID. Reaction of ketone **1** with amine **2** in toluene under orbital shaking (48 h, 33 °C, 133 rpm). (b) In the presence of lipase. (b) In the absence of lipase. Conditions of GC-FID analysis: DB-5 column (30 m x 0.25 mm x 0.25 µm) Ti = 80 °C, Tf = 250 °C, r = 10 °C/min, t = 22.0 min

© 2014 Global Journals Inc. (US)



SI- 5 : Chromatogram and mass spectrum obtained by GC-MS (EI, 70 eV) for the reaction of ketone **1** with amine **2** in hexane under orbital shaking for 4 days (33 °C, 133 rpm). (a) Ketone **1**. (b) Amine **2**. (c) Mass spectrum obtained from a signal with a retention time of 14.5 min. (d) Mass spectrum obtained from a signal with a retention time of 15.9 min. Conditions of GC-MS analysis: DB-5 column (30 m x 0.25 mm x 0.25 μm) Ti = 50 °C, Tf = 270 °C, r = 10 °C/min, t = 32.0 min

GC-MS (EI, 70 eV):


2-cyclohexen-1-one **1**, C_6H_8O, *m/z* (%): 68 (100), 96 (M^+, 41), 40 (21), 55 (6), 81 (0.5).

1-phenylmethanamine **2**, C_7H_9N, *m/z* (%): 106 (M^+-1, 100), 107 (M^+, 61), 79 (42), 91(12), 51 (11), 65 (6).


N-(cyclohex-2-en-1-ylidene)-1-phenylmethanamine **3**, $C_{13}H_{15}N$, *m/z* (%): 65 (15), 91 (100), 185 (M^+, 30), 157 (7).

3-(benzylamino)cyclohexanone **4** $C_{13}H_{17}NO$, *m/z* (%): 91 (100), 146 (44), 160 (32), 203 (M^+, 30), 133 (27), 106 (31), 65 (13).

© 2014 Global Journals Inc. (US)




SI-6 : Cleavage probable at the bond β to the ring, giving the tropylium ion



SI- 7 : Chromatograms obtained by GC-FID. (a) Standard of amine **2**. (b) Standard of ketone **5**. (c) Reaction of amine **2** with ketone **5** in methanol under orbital shaking in the presence of lipase (24 h, 33 °C, 133 rpm). (d) In the absence of lipase. Conditions of GC-FID analysis: DB-5 column (30 m x 0.25 mm x 0.25 μm) Ti = 80 °C, Tf = 250 °C, r = 10 °C/min, t = 22.0 min

© 2014 Global Journals Inc. (US)




Sl- 8 : Chromatogram and mass spectrum obtained by GC-MS (EI, 70 eV) for the reaction of ketone **5** with amine **2** in hexane under orbital shaking (24 h, 33 °C, 133 rpm). (a) Amine **2**. (b) Ketone **5**. (c) Mass spectrum obtained from a signal with a retention time of 15.9 min. Conditions of GC-MS analysis: DB-5 column (30 m x 0.25 mm x 0.25 μm) Ti = 50 °C, Tf = 270 °C, r = 10 °C/min, t = 32.0 min

GC-MS (EI, 70 eV):

3-methylcyclohex-2-en-1-one **5**, $C_7H_{10}O$, *m/z* (%): 82(100), 110(40, M^+), 54(36), 41(8).


1-phenylmethanamine **2** C_7H_9N, *m/z* (%): 106 (M^+-1, 100), 107 (M^+, 61), 79 (42), 91(12), 51(11), 65 (6).

N-(3-methylcyclohex-2-en-1-ylidene)-1-phenylmethanamine **7**, $C_{14}H_{17}N$, *m/z* (%): 91(100), 199 (M^+, 30), 65(17), 184(11), 79(10).




Sl-9 : Cleavage probable at the bond β to the ring, giving the tropylium ion

© 2014 Global Journals Inc. (US)



SI-10 : Chromatogram obtained by GC-MS. Reaction of amine **2** with ketone **8** in hexane under orbital shaking (96 h, 33 ^{o}C, 130 rpm). Conditions of GC-MS analysis: DB-5 column (30 m x 0.25 mm x 0.25 μm) Ti = 50 ^{o}C, Tf = 270 ^{o}C, r = 10 ^{o}C/min, t = 22.0 min

© 2014 Global Journals Inc. (US)



SI-11 : Chromatograms and mass spectrum obtained by GC-MS (EI, 70 eV). (1) Non-enzymatic reaction of amine **2** with ketone **8** in hexane under microwave irradiation (3 h, 60 °C, 70 W). (2) Enzymatic reaction of amine **2** with ketone **8** in hexane under microwave irradiation (5 h, 60 °C, 70 W). (a) Amine **2**. (b) Ketone **8**. (c) Mass spectrum obtained from a signal with a retention time of 15.0 min. (d) Mass spectrum obtained from a signal with a retention time of 19.0 min. (e) Mass spectrum obtained from a signal with a retention time of 11.3 min. Conditions of GC-MS analysis: DB-5 column (30 m x 0.25 mm x 0.25 μm) Ti = 50 °C, Tf = 270 °C, r = 10 °C/min, t = 32.0 min

© 2014 Global Journals Inc. (US)


GC-MS (EI, 70 eV):

2,5-dimethyl-*para*-benzoquinone **8**, $C_8H_8O_2$, *m/z* (%): 136(M^+, 100), 68(66), 79(28), 108(29), 40(28), 96(22), 121(3).


4-(benzylimino)-2,5-dimethylcyclohexa-2,5-dienone **9**, $C_{15}H_{15}NO$, *m/z* (%): 225(M^+, 100), 148(70), 224(52), 209(23), 77(14).

N-benzylidene-1-phenylmethanamine **10**, $C_{14}H_{13}N$, *m/z* (%): 65 (17), 91(100), 92 (37), 117 (11), 194 (40), 195 (M^+, 38).


N-benzylacetamide **11** $C_9H_{11}NO$, *m/z* (%): 106 (100), 149 (M^+, 79), 91(27), 79(18), 43(21).



SI-12 : Aza-Michael reactions using CALB on an orbital shaker and microwave irradiation




SI- 13 : Chromatograms obtained by GC-MS. (a) Enzymatic reaction of amine **2** with ketone **8** in hexane under orbital shaking in triplicate (48 h, 33 °C, 130 rpm). (b) Enzymatic reaction of amine **2** with ketone **8** in hexane after extraction into ethyl acetate and water after 24 hours on an orbital shaker (33 °C, 133 rpm). Conditions of GC-MS analysis: DB-5 column (30 m x 0.25 mm x 0.25 µm) Ti = 50 °C, Tf = 270 °C, r = 10 °C/min, t = 32.0 min



SI- 14 : Extraction of aza-Michael reaction with ethyl acetate and water after 24 h on an orbital shaker (33 °C, 133 rpm)

© 2014 Global Journals Inc. (US)



SI-15 : Acetylation of amine **2** by EtOAc

© 2014 Global Journals Inc. (US)

This page is intentionally left blank

© 2014 Global Journals Inc. (US)

Global Journal of Science Frontier Research: B
Chemistry
Volume 14 Issue 1 Version 1.0 Year 2014
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Water Quality of River Tungabhadra Due to the Discharge of Industrial Effluent at Harihar, District Davanagere, Karnataka State, India

By Basavaraj M. Kalshetty, Shobha. N & M. B. Kalashetti
Bharathier University, India

Abstract- Water is a prime mover of all activities and essential feature for all modern developments. Water is distributed in different forms, such as rain water, river water, spring water and mineral water. Rain water is considered to be the purest form, however, is associated with dissolved gases such as CO_2, SO_2, NH_3 etc., from atmosphere. Water used for Industrial development and municipal purposes, it is better to ensure the quality of water for these purposes. There is influence of Industrial waste. Sewage on the water quality, the waste products can change the water Chemistry. Water gives life to Industries, but, Industries kill the water Chemistry. The waste water which emerges out after uses from industries have no definite composition, the pollutants associated with industrial effluents such as organic matter, inorganic dissolved solids, fertilizer materials, suspended solids, heavy metals from toxic pollutants and micro organisms and also pathogens. The industrial wastes are responsible for water color, turbidity, odour, hardness, toxic elements, bacteria and micro organisms. Industrial wastes contain poisonous chemicals which are difficult to remove from its homogeneous solution state.

GJSFR-B Classification : FOR Code: 039999

WATER QUALITY OF RIVER TUNGABHADRA DUE TO THE DISCHARGE OF INDUSTRIAL EFFLUENT AT HARIHAR, DISTRICT DAVANAGERE, KARNATAKA STATE, INDIA

Strictly as per the compliance and regulations of :

© 2014. Basavaraj M. Kalshetty, Shobha. N & M. B. Kalashetti. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Water Quality of River Tungabhadra Due to the Discharge of Industrial Effluent at Harihar, District Davanagere, Karnataka State, India

Basavaraj M. Kalshetty [α], Shobha. N [σ] & M. B. Kalashetti [ρ]

Abstract- Water is a prime mover of all activities and essential feature for all modern developments. Water is distributed in different forms, such as rain water, river water, spring water and mineral water. Rain water is considered to be the purest form, however, is associated with dissolved gases such as CO_2, SO_2, NH_3 etc., from atmosphere. Water used for Industrial development and municipal purposes, it is better to ensure the quality of water for these purposes. There is influence of Industrial waste. Sewage on the water quality, the waste products can change the water Chemistry. Water gives life to Industries, but, Industries kill the water Chemistry. The waste water which emerges out after uses from industries have no definite composition, the pollutants associated with industrial effluents such as organic matter, inorganic dissolved solids, fertilizer materials, suspended solids, heavy metals from toxic pollutants and micro organisms and also pathogens. The industrial wastes are responsible for water color, turbidity, odour, hardness, toxic elements, bacteria and micro organisms. Industrial wastes contain poisonous chemicals which are difficult to remove from its homogeneous solution state.

I. Introduction

Rivers are life line of living being and constitute an integral part of both rural and urban community as a source of drinking and cooking purposes. Water used by the public must be free from disease causing bacteria; toxic chemicals, excessive amount of minerals and organic matter1. Fresh water becomes polluted due to three major reasons, excess nutrients from sewage, industries, mining and agriculture. According to recent investigations ground water contamination also increasing due to anthropogenic activities like disposal of waste, sewage, industrial waste2. Due to rapid Industrialization, Urbanization, therefore, pollution of water resources needs a serious and immediate attention through periodical check up of water quality.

Keeping in view the main objective of the present investigation is the evaluation of physico-chemical aspects of Tungabhadra River water from the selected locations at Harihar Polyfibre Industries, to specify accurately and timely information regarding the quality of river water at industrial effluent disposal point and both at upstream and downstream flow of water.

The present findings may be helpful to shape sound public policy and to implement water quality improvement programme effectively as well as efficiently.

Study area: The two Major Industries are discharging their effluents in the river Tungabhadra at Kumarpattanam near Harihar town, District Davanagere, Karnataka State.

1. Harihar Polyfibres Factory (HPF) is the largest Rayon grade pulp manufacturing unit in India (1972). Daily manufacturing capacity approximately 200 tones, generating more than 30 thousands liters of industrial effluents per day.
2. Grasilene Rayon Factory (GRF) 1977 produces about 850 tones of Grasilene fiber per month generating about 10 thousand liters of highly polluted effluents.

Both factories locating on the left bank of river Tungabhadra on NH-4 at Kumarpattanam. The area selected for studies are up stream flow of water from discharge point, discharge point and downstream flow of water from the effluent disposal point. The water quality of upstream investigated for the comparative study of effluent effect on water body. The partially treated effluent run in one canal for a distance of half km through a small village called Nalavagal, before joining the river Tungabhadra at only one location. The waste water was colored before mixing with river water, carrying lot of foam along the downstream, recently the treatment of effluents from these two factories are adapted little new technology or discharging into river without any proper treatment before discharging.

People residing in these areas utilize river water for their daily needs. Sampling locations were selected on the basis of the detailed survey of the study area and discussion with local public, the details regarding the locations is given in Table 1.

Author α: B.L.D.E'S Science College Jamkhandi, Bagalkot District, Karnataka State, India. e-mail: drkalshetty@gmail.com
Author σ: ITC Limited, ITC Life Science & Technology Centre, Peenya Industrial Area, 1st Phase, Bangalore-58. & Research Scholars, Department of Chemistry, Bharathier University, Coimbatore.
Author ρ: P.G. Studies in Department of Chemistry, Karnataka University, Dharwad.

© 2014 Global Journals Inc. (US)

Table 1: Details of sampling locations

No.	Stream	Sampling location	Distance from discharging point.
1	Upstream	Harihar side (Makanur)	0.5 km.
2	Discharge point	Factory point	50 feet
3	Down stream	Nadiharalahalli	0.5 km.
4	Down stream	Airani	5 km.

a) Experimental

All the samples were collected in clean polythene bottles with necessary precaution and brought to the research Laboratory without the addition of any preservatives and subjected to the physico-chemical analysis within 24 hours after collection, in order to assess their quality and portability water samples were analyzed using standard methods2, 3, and 4. All the chemicals used were A.R. Grade; Double distilled water was used for the preparation of various solutions. The physico-chemical parameters such as Temperature, pH, EC, TDS, DO, TH, TA, Calcium, Magnesium, Sodium, Potassium, Chloride, Nitrate and Sulphate were determined by standard procedures5. The pH and EC were carried out by using pH-meter and Electrical conductivity meter respectively. Sodium (Na+) and potassium (K+) were determined by using flame photometer; sulphate (SO4'') ion concentration was determined by using visible spectrophotometer.

II. Results and Discussion

The assessed physico-chemical parameters after the analytical determination given in Table 2 to Table 5. The values were compared with the standard values given by WHO6, ISI7 and with ICMR8 as given in Table 6. The water samples collected at upstream (spot 1) were not having any objectionable color appearance, taste or odour during the monitoring periods (August-2013, September- 2013, October- 2013 and November-2013). The samples collected at the industrial disposal point the water is found in thick dark brown coffee colored before mixing with river water, carrying lot of foam along the downstream. The water samples at the downstream from the discharge point were collected and the analyzed. The results indicate the river water under goes natural purification during its flow along the stream.

Table 2 : The average values of Physico-chemical parameters and Trace elements in the study locations:

Location: upstream (Spot 1) 0.5 km. from discharge point towards Harihar- side

Date	Temp.	pH	Ec	TDS	COD	BOD	DO	Cl-	CO3''	HCO3'	Calcium
25.08.13	26.6	7.3	298	97	24	6.4	6.0	36.8	ND	2.1	22
22.09.13	27.8	7.4	162	96	16	6.3	6.6	28.0	ND	2.2	15
26.10.13	27.5	7.3	181	88	19	3.5	6.8	26.5	0.7	2.6	26
24.11.13	26.6	7.2	149	78	10	3.1	6.9	34.0	0.6	2.8	18
Date	Mg	Na	K	SO4''	TH	Zn	Cu	Fe	Mn	Cobalt	Nickel
25.08.13	5.0	34	7	08	65	0.189	0.03	0.85	0.14	0.04	0.10
22.09.13	3.2	20	4	12	68	0.281	0.06	1.04	0.16	0.06	0.09
26.10.13	4.0	24	5	17	45	0.314	0.02	0.98	0.09	0.06	0.02
24.11.13	3.2	21	6	14	68	0.201	0.09	0.92	0.12	0.06	0.13

Table 3 : Location: Discharge point (Spot 2) 50 feet away from the factory

Date	Temp.	pH	Ec	TDS	COD	BOD	DO	Cl-	CO3''	HCO3'	Calcium
25.08.13	28.0	7.2	2896	1388	172	74	1.8	123	0.5	13.4	152
22.09.13	28.0	7.3	1099	1622	122	107	2.9	127	0.8	9.92	172
26.10.13	27.9	7.2	798	898	130	112	2.8	226	1.36	6.67	95
24.11.13	26.9	7.0	787	1012	179	105	3.8	276	1.80	12.8	87
Date	Mg	Na	K	SO4''	TH	Zn	Cu	Fe	Mn	Cobalt	Nickel
25.08.13	39	89	30	330	21	4.95	0.07	1.56	0.34	0.12	0.86
22.09.13	54	110	28	194	194	2.98	0.09	2.08	0.25	0.84	0.92
26.10.13	18	92	27	230	330	1.50	0.07	1.58	0.21	0.18	0.20
24.11.13	22	59	26	270	250	3.15	0.09	1.71	0.42	1.98	0.19

© 2014 Global Journals Inc. (US)

Table 4 : Location: Downstream (Spot 3) Near Nandihalli village 0.5 km. from the discharge point

Date	Temp.	pH	Ec	TDS	COD	BOD	DO	Cl-	CO3''	HCO3'	Calcium
25.08.13	29	7.6	1950	486	49	31	4.8	41.2	0.3	9.8	86
22.09.13	29	7.8	799	589	18	36	4.9	33.0	0.4	6.4	44
26.10.13	29	7.7	678	414	27	27.5	4.8	37.4	0.9	3.2	36
24.11.13	26.8	7.9	317	465	29	27.0	4.3	47.0	0.8	6.7	61
Date	Mg	Na	K	SO4''	TH	Zn	Cu	Fe	Mn	Cobalt	Nickel
25.08.13	28	48	27	390	99	1.96	0.06	0.95	0.30	0.14	0.38
22.09.13	27.2	55	24	129	95	2.92	0.09	1.24	0.21	0.49	0.64
26.10.13	7.2	61	21	276	178	1.96	0.06	0.31	0.14	0.12	0.09
24.11.13	5.2	60	18	165	132	0.99	0.06	1.24	0.24	1.04	0.46

Table 5 : Location: Downstream (Spot 4) Near Airani village 5 km from the discharge point

Date	Temp.	pH	Ec	TDS	COD	BOD	DO	Cl-	CO3''	HCO3'	Calcium
25.08.13	29.1	7.9	1546	461	47	27.0	5.6	54.5	0.2	6.8	91.0
22.09.13	29.0	8.3	825	612	14	28.5	5.8	31.1	0.1	3.7	51.4
26.10.13	29.1	7.6	640	220	23	20.1	6.2	54.2	0.5	2.7	41.0
24.11.13	26.4	8.2	402	312	18	31.0	6.6	51.0	0.5	4.2	54.0
Date	Mg	Na	K	SO4''	TH	Zn	Cu	Fe	Mn	Cobalt	Nickel
25.08.13	23.3	47	25	280	65	0.98	0.07	0.86	0.21	0.09	0.40
22.09.13	16.3	54	19	101	89	1.86	0.06	1.36	0.14	0.62	0.32
26.10.13	5.8	31	18	176	92	0.85	0.06	0.27	0.17	0.09	0.08
24.11.13	4.9	26	14	94	122	0.82	0.03	0.94	0.18	0.59	0.24

a) Temperature

All physiological activities and life processes of aquatic organisms are generally influenced by water temperature9, it is an important parameter because effects bio-chemical reactions in aquatic organisms. A raise in temperature of water leads to the speeding up of chemical reactions in the water body. Temperature reduces the solubility of gases and amplifies the tastes and odours. In the present investigation the temperature range was found in between 26.9oC to 28.30oC at the industrial effluent discharge point. The temperature of upstream water found in between 26.6oC to 27.8oC where as downstream water temperature at spot S3 and S4 were in between 26.4oC to 29.1oC.

Table 6 : Prescribed legal limits and guidelines

Parameter	WHO	ISI	ICMR
pH	7.0 – 8.5	6.5-8.5	6.5-9.2
EC	1.4 ds/m	1400µs/cm	250
TDS	500	500	1500-3000
BOD	5.0	5.0	---
COD	10.0	10.0	---
DO	5.0	5.0	5.0
Cl'	250	250	1000
Calcium	75-100	45-75	200
Magnesium	30-150	30	50
Sodium	200	200	---
Potassium	1.4	1.4	---
Sulphate	150-250	150-200	400
TH	300	300	600
TA	120	200-600	200-600
Zinc	0.75	---	---
Copper	0.60	---	---
Iron	<1	0.3	---
Manganese	2.0	0.3	---
Nitrate	40-100	40-100	40-100
Fluoride	1.0-1.5	1.0-1.5	1.0-1.5
Color	5 HU	5 HU	5 HU
Odour	Unobjectionable	Unobjectionable	Unobjectionable
Turbidity	5 NTU	5 NTU	5 NTU

© 2014 Global Journals Inc. (US)

b) Taste and Odour

Disagreeable odour and taste in water may be because of the presence of vegetation, inorganic constituents and organic matters, the discharge of waste water and industrial effluents in water bodies. WHO and BIS limit: unobjectionable or agreeable due to aesthetic condition. But, at the present studies water samples were found to be odourless, whereas the water chemistry was spoiled at the discharge disposal point. Downstream water under gone the natural purification possessing tasteless and odourless characters.

c) Turbidity

Turbidity in water is caused by suspended matter finely divided organic and inorganic matter, soluble colored organic compounds are responsible for the developing the turbidity in water. Plankton and other microscopic organisms make the water turbid. Turbidity was found to be O NTU. WHO and BIS limit: 5-10 NTU; aesthetic consideration and harmful bacteria may be associated with particles. In the present study the water sample at industrial effluent disposal point was found to be in coloured due to industrial waste and finely divided organic and inorganic matter.

d) pH (Power of Hydrogen)

The pH of natural water is important index of hydrogen ion activity and it is resulting value of the acid - base interaction of a number of mineral and organic components in water. pH is an important ecological factor and is a term used and universally to express the intensity of the acid and alkaline condition of the water samples. Most of the water samples were slightly alkaline due to the presence of carbonates (CO_3'') and bicarbonates (HCO_3'). pH-value determines the equilibrium between free CO_2, HCO_3' and CO_3''. The present study reveals that the pH of upstream water found in between 7.2 to 7.4, where as in downstream water was found to be in between 7.6 to 8.3 indicates more alkaline in nature and may reflect contamination of OH- ions by strong bases such as sodium hydroxide and calcium hydroxide10. At the disposal point the pH of water was ranged from 7.0 to 7.3 (WHO and BIS limit: 6.5 to 8.5; health reflect problem, may affect mucous membrane and problem in water supply system).

e) Electric Conductivity(EC)

Electric conductivity is caused due to presence of electrolytes which dissociate in to cations and anions, It is a measure of water capacity to convey electric current. It is an indicator of the degree of mineralization of water. The EC is correlated with total dissolved solids11. It signifies the amount of total dissolved salts. Presence of high amount of dissolved inorganic substances in ionized form.

In the present investigation the EC-values of water sample at the industrial effluent point during monitoring periods ranged in between 789 to 2896 Micromhos/cm, and indicate the presence of some ionic matter such as Ca,Mg,Cl,SO4,CO3,HCO3 and some trace elements.

The EC-values of upstream and downstream water samples were in the legal limit, most of the water sample content substantial amounts of dissolved CO_2, HCO_3' and hydroxides, these constituents are the results of dissolution of minerals in the soil and atmosphere12.

f) Density, Surface Tension and Viscosity

Density of the water samples at the disposal point was 0.9982 gram/cm3, while surface Tension was 71.812 dyne/cm. Viscosity of water sample was calculated as 0.0091 poise/cm/sec. which was in the comparable ranges.

g) Total dissolved solids (TDS)

TDS is an index of solid present in dissolved form and estimated as being 0.5 to 0.9 times the conductivity in ms/cm. It indicates the salinity behavior of ground water. Water containing more than 500 ppm of TDS is not considered desirable for drinking water supplies, but in unavoidable case 1500 ppm is also allowed13 for domestic use. In the present investigation the TDS values of upstream water samples at various monitoring months found in between 78 ppm to 97 ppm. The downstream water samples are also found in the normal prescribed legal ranges ie., in between 129 ppm to 589 ppm, where as the TDS values of water sample at the industrial effluent point was ranged in between 787 ppm to 2896 ppm, more than the proscribed value, this indicates that the factory effluents contains large amount of dissolved solids, this salinity behavior of water sample due to the dumping of chemicals both organic and inorganic matter in to the river water. (WHO and BIS limit: 500 ppm -2000 ppm; palatability decrease and may cause gastro intestinal irritation).

h) Dissolved Oxygen (DO)

Oxygen is a regulator of metabolic processes of plants and animals. The depletion of oxygen level is due to high amount of organic wastes and industrial wastes. In the present investigation water samples at the industrial effluent discharge point their DO values found to be in between 1.8 ppm to 3.8 ppm during the monitoring months. While the upstream water showed the DO values in between 6.0 to 6.9 ppm and the downstream water showed the DO values in between 4.3 ppm to 6.6 ppm this indicates the river water gets natural purification. The lower value of DO at the effluent discharge point water may be due to inclusion of high inorganic and organic load in the water which leads to oxygen depletion.

i) Total Hardness (TH)

The total hardness may be due to the presence of divalent cations like Ca, Mg found abundant in water. The total hardness is sum of the calcium and

© 2014 Global Journals Inc. (US)

Magnesium concentrations, both expressed as Calcium carbonates in ppm values. Water is conventionally classified as hard or soft from the following classification 50-100 ppm (Soft), 100-250 ppm (Moderately hard), 250 – 350 ppm (Hard). Hardness is the property of water which prevents the later formation with soap and increases the boiling points of water14.

In the present investigation the TH of the upstream water samples ranged in between 45 ppm to 68 ppm. Similarly the TH values of downstream water found in between 65 ppm to 178 ppm during the monitoring periods. Whereas the water samples at industrial effluent point showed TH range was from 194 ppm to 330 ppm. Thus the water samples were found to be moderately hard and thus need to be treated before using it for the domestic uses.

j) Calcium and Magnesium Hardness

The water samples collected at the industrial effluent discharge point contain the calcium hardness from 87 ppm to 172 ppm these values were close resembling and were within the permissible limit. Magnesium hardness values from 18 ppm 54 ppm and were within the permissible limit of 30 to 150 ppm (WHO). Jain et al15 and other reported that the high concentration of hardness (150 to 300 ppm) may cause heart disease and kidney problems.

k) Alkalinity

Alkalinity of water is acid neutralizing capacity of the water to predestinated pH. Alkalinity in water is mainly due to CO3'' HCO3' and OH- content. Borates, phosphates, silicates or other bases if present also contribute for alkalinity. (WHO and BIS limit: 200 to 600 ppm; taste become unpleasant). In the present study the water samples collected at disposal point the total methyl orange alkalinity was found in between the range was within the prescribed limits.

l) Biological Oxygen Demand(BOD)

Biological oxygen demands are a measure of the quantity of oxygen used by microorganisms (aerobic bacteria) in the oxidation of organic matter. In the present work the BOD values of water samples during the monitoring periods at the industrial effluent disposal point was found in between 74 ppm to 112 ppm, this indicates the effluent runoff carries chemical wastes during the polyfibre synthesis. This is due to the discharge of untreated or partially treated industrial effluent directly in to the river. There are makeable variation in the BOD content along the downstream from the discharge point during the study period which was ranged in between 20.1 ppm to 36 ppm. The upstream water showed low BOD values 3.1 ppm to 6.4 ppm. The decrease in BOD and increase in DO from effluent discharge point was due to natural self purification.

m) Chemical Oxygen Demand (COD)

COD determines the oxygen required for the chemical oxidation of organic matter. COD values conveyed the amount of dissolved oxidizable organic matter including non-biodegradable matter present in it. The COD value in the sample effluent was found to be 122 ppm to 179 ppm; the high COD value determines the organic load. The COD values found within the 10 ppm to 24 ppm found within the permissible limit.

n) Chloride (Cl)

The chloride concentration serves as an indicator of pollution by sewage, industrial effluents. Chloride occurs in all ground waters widely in varying concentration. Excessive chloride in potable water is not particularly harmful. Chloride in excess (more than 250 ppm) imparts a salty taste to water. People accustomed to higher chloride in water are subjected to laxative effects16. In the present investigation the chloride values ranged from 26.5 ppm to 36.8 ppm from the upstream water samples, and chloride found in between 31.08 ppm to 54.5 ppm along the downstream water. Whereas the chloride content of the water samples collected at the effluent discharge point ranged in between 123 ppm to 276 ppm, slight more than the permissible legal range. The tolerance of chloride by human being varies with climate, exertion and loss through perspiration. (WHO and BIS limit: 250 to 1000 ppm decreases potability, salty taste and corrosion).

o) Sodium (Na)

Sodium levels in ground waters vary widely; depends upon geological formation. In surface water generally the sodium concentration ranges in between 1 and 300 ppm depending upon the geographical area. Excessive intake of sodium chloride causes vomiting. (Guideline values: 200 ppm; based on taste threshold and health consideration). In the present investigation the sodium concentration of the water samples collected from upstream water,at the industrial effluent disposal point and water samples of downstream were found in between the ranges of 20 ppm to 34 ppm, 59 ppm to 110 ppm and 26 ppm to 54 ppm respectively, this indicates the more concentration of sodium in the industrial effluent point due to the chemical combination of compounds leads to change in the quality of water.

p) Potassium(K)

The quality of water in the natural process most commonly found the concentrations of both sodium and potassium. The concentration of potassium is quite lower than sodium. It remains most common in solution without undergoing any precipitation. The potassium concentration of upstream river water sample found to be very low i.e., less than the prescribed limit. Whereas the potassium concentration in water sample collected at discharge point found in between 26 ppm to 30 ppm also less than legal range. The concentration of

© 2014 Global Journals Inc. (US)

potassium falls with the downstream due to the natural purification.

q) Acidity

Acidity is a measure of the effects of combination of compounds and conditions in water. It is the power of water to neutralize OH- and is expressed in terms of CaCO3. Water attains acidity from industrial effluents, municipal waste (drainage) and from humic acid. No phenolphthalein acidity and methyl orange acidity was found in the studied water samples.

r) Pathogens

The most common danger associated with drinking water is contamination by sewage, by other waste or by human or animal excreta. In general this is determined by conducting test for total coli form counts, fecal coli form counts. The pathogen observed was salmonella species which is gram negative and short rod shape. (BIS limit: 10 CFU/ 100 ml and absent; causes illness and fatal for life).

s) Sulphate (SO4)

In the present investigation sulphate concentration of upstream water sample was recorded and ranged in between 8 ppm to 17 ppm. In downstream water was ranged in between 94 ppm to 280 ppm. The SO4 content in the water sample collected at the discharge point found in between 194 ppm to 330 ppm. All the values found within the permissible limits. It is known that the sulphate concentration in water samples around 1000 ppm; it has Laxative effect and causes gastrointestinal irritation17. Excess Na2SO4 in water causes cathartic action. High concentration of Na and MgSO4 is associated with respiratory illness.

t) Iron (Fe)

Iron is an important essential element to human body which is present in hemoglobin and myoglobin. When iron concentration exceeds permissible limit in drinking water it gives stringent taste to water. According to WHO standards, limit of Fe concentration in drinking water is 0.3 mg/L, exceeding which causes toxicity. In the present investigation the iron content of the water samples analyzed was found to be more than the permissible limits. These values vary from 1.56 ppm to 2.08 ppm and indicate the effluent consists with more iron content because of using chemical compounds in industrial process.

u) Copper (Cu)

Copper is essential element of human body, but excessive large doses may lead to mucosal irritation, corrosion, wide spread capillary damage, hepatic and renal damage and also to central nervous system. Copper below 20 µ gram/L may found in drinking water. Due to the industrial effluent concentration of copper in water may increase. The concentration of copper found in natural waters is not known to have adverse effects on humans. Though copper in excess of 1.0 mg/L may impart some taste to water18. Toxicity of copper to aquatic life is dependent on the alkalinity of water. At lower alkalinity, copper is more toxic to aquatic life. In the present investigation the concentration of copper recorded a maximum 0.09 ppm at industrial effluent disposal point.

v) Zinc (Zn)

Zinc is an essential element and it is necessary for the functioning of various enzyme systems, deficiency and low intake of Zinc of which results in retardation of growth, immaturity and anemia, condition known as “Zinc deficiency syndrome”. Symptoms of Zn toxicity in humans include vomiting, dehydration, abdominal pain, dizziness and lack of muscular coordination. Zinc imparts undesirable bitter astringent taste to water at levels above 5.0 m gram/L. zinc deficiency in human body may result in infantilism, impaired wound healing and several other diseases. Pollution from industrial sources to a great extent is responsible for high concentration of zinc in water. The concentration of zinc in the study area (at discharge point) ranges from 1.50 ppm to 4.95 ppm which below the prescribed limits.

w) Manganese (Mn)

Excess Manganese in a diet prevents the use of Iron in the regeneration of blood hemoglobin. Large doses of Manganese cause apathy, irritability, headaches, insomnia, and weakness of the legs.

Psychological symptoms many also develop including in impulsive acts, absent-mindedness, hallucinations, aggressiveness and unaccountable laugher. Finally a condition similar to Parkinson’s disease may develop19. Manganese Pollution from HPF industrial effluent discharging of chemical waste at disposal site was analyzed, the concentration of Manganese in the study area ranged from 0.21 to 0.42 mg/L [BIS Limits: 0.30 mg/L]. Hence, Manganese-values exceed this limit at some monitory period. Thus, the quality of water is suspect able for drinking and other domestic Purposes. The other elements such as Cobalt and Nickel are also important and essential to human body.

III. Conclusion

- Data reveals that the river water polluted to some extent at the industrial effluent discharge point, as indicated by a very high value of COD and other activities. The data also indicates that the water bodies total hardness were sound to be beyond the permissible limits as prescribed by WHO and other standards.
- Qualitatively the water in the studied area is not potable, hard and chemical mixed at the discharge

© 2014 Global Journals Inc. (US)

point. Hence, the water at this area is may be used for irrigation but not for domestic and drinking purposes.

- From the analysis it may be concluded that proper environment management plan may be adopted to control the release of effluent. Hence, it is suggested to exercise all the necessary precaution before the water is used for domestic and irrigation. Otherwise it may lead to much adverse health effect.
- To improve or to keep the quality of water at the effluent discharge point, the industrial waste should be treated properly before disposal into the river stream. Hence, there should be continuous monitoring of the pollution level.
- The polluted water due to the disposal of industrial effluent, used for irrigation which would not improve the soil fertility but also reduces the enrichment of nutrients present in the soils.
- The Figures 1 to 3 indicate the comparative studies in the variation of the physicochemical parameters and presence of trace elements in the water samples at various selecting points. The result implies the water sample at industrial effluent discharge point found more polluted rather than upstream, the water in the downstream itself gets natural purification.

IV. Acknowledgement

The authors are thankful to Dr. N. Gopukumar, Deputy Secretary, UGC, and South Western Regional Office Bangalore for their help and encouragement by funding the financial assistance in term of Minor Research Project. We strongly acknowledge Dr. C.N.R.Rao FRS, Chairman VGST Bangalore, and Dr. Ananth Raj Executive secretary VGST group Bangalore for providing financial support in terms of Major Projects. the authors are very much thankful of our college management and principal of our college for providing the research laboratory facilities and encouragement.




Figure 1 : Comparative study of physicochemical parameters at Upstream (Series 1), Discharge point (Series 2) and Downstream (Series 3) water samples

© 2014 Global Journals Inc. (US)




Figure 2 : Comparative study of physicochemical parameters at Upstream (Series 1), Discharge point (Series 2) and Downstream (Series 3) water samples




Figure 3 : Comparative study of Elements at Upstream (Series 1), Discharge point (Series 2) and Downstream (Series 3) water samples

References Références Referencias

1. B. N. Yadav, B.N. Sigh, Physico-Chemical studies on Pollution Potential River Ganga at Kanpur, ACTA ciencia India 25 (C) (2): 285–290(2009).
2. APHA, standard methods for the examination of water and waste water, 18th Edition: American Public Health Association, Washington D.C (1985).
3. Manivasakam N, Physico chemical Examination of water, sewage and Industrial effluents 5th Edition Pragati Prakashan, meerut (2005).
4. Kotaiab B and Kumarswamy N. Environmental Engineering Laboratory manual, 1st Ed chaotar publishing House Pvt, Ltd. India.
5. B. M. Kalshetty, R. C. Sheth, R. S. Gani, S. S. Karabasannavar and M. B. Kalashetti "Water Quality Assessment of Water Quality in and Around Jamkhandi City, Bagalkot District, Karnataka State, India" GLOBAL JOURNAL OF SCIENCE FRONTIER RESEARCH CHEMISTRY, Volume 13, Issue 4, Version 1.0, Year 2013. Global Journals Inc. (USA).
6. WHO, International standards for Drinking water, Geneva, (1971).
7. ISI, International Standards of water quality for Drinking water is: 10500, 1983.
8. ICMR, Manual standards of water Quality for Drinking water supplies, 2nd ED, Indian council of medical research, New Delhi Pp.21 (1973).
9. H. Murhekar Gopal Krushna, Res, J. chem. Sci, 1 (4), 117, 2011. B. M. Kalshetty, R. C. Sheth, R. S. Gani, S. S. Karabasannvar and M. B. Kalashetti " Assessment of water Quality in and around Jamkhandi city, Bagalkot District, Karnataka Global Journal of Science Frontier Research V0l.13, issue 4, version 1.0, year 2013 , Global Journals Inc.(USA).

© 2014 Global Journals Inc. (US)

10. D. Langmuir, Aqueoan Environment chemistry. Prentise Hall, Inc, New Jersey, 1997.
11. S. Arya, V. Kumar, Minaskhi, A. Dhaka, int. Multidiscipline. Res. J. 1(7), 11,20111
12. Nagaraju, S.Sursh, K.Killham and K. Hudson Edward, Turkish J-Eng. Env. Sci, 30, 203, 2006.
13. Shrinivas Rao B. and venkateswarlu P, Indian J. Environ Prot, 20(3), 1617, 2000.
14. Trivedy R.K. and Good P.K. Chemical and Biological methods of water Pollution studies environmental publication, Karad, 1986.
15. C. K. Jain, K.K.S. Bhatia, Y. Vijay, Ind. J. Envi, Health, 39(3), 182, 1982.
16. Sudhir Dahiya and Amarjeet Kour. J. Env. Poll. 6(4), 281, 1991.
17. H. murhekar Gopal Krushna, Res, J. chem. Sci, 1(4), 117, 2011.
18. E. Merian(Ed), metal and their compounds in the Environment: Occurrence, Analysis and Biological Relevance, UCH, weinheim, NewYork (1991).
19. Nitin Gupta S.M. Nafees, M.K. Jain and Kalpanar "Study on heavy metal contamination in GW at outer Skirts of kota cents , Rajastan, media" int J. C. Lem. S. S. i, 9(1), 2011, 285-306.

© 2014 Global Journals Inc. (US)

This page is intentionally left blank

© 2014 Global Journals Inc. (US)

Global Journal of Science Frontier Research: B
Chemistry
Volume 14 Issue 1 Version 1.0 Year 2014
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Synthesis, Characterization and Antimicrobial Activity of Azetidin-2-One based Phenyl Sulfonyl Pyrazoline Derivatives

By Shailesh H. Shah, Pankaj S. Patel, Mitesh Prajapati, Bharat Vaghari & Jayesh Parmar

Patel Jdkd Science College, India

Abstract- A new series of 4-(4-Chlorophenyl)-3-chloro-1-{4-[5-(Substitutedphenyl)-1-(phenylsulfonyl)-4,5-dihydro-pyrazol-3- yl]phenyl}azetidin-2-one are synthesized by reacting 3-chloro-1-{4-[5-(Substitutedphenyl)-4,5-dihydro-pyrazol-3-yl]phenyl}-4-(4-Chlorophenyl) azetidin-2-one with Benzene sulfonyl chloride in presence of pyridine. All these compounds were characterized by means of their IR, 1H NMR, Spectroscopic data and were tested for their antibacterial and antifungal activities by broth dilution method.

Keywords: *chalcones, phenyl sulfonyl-pyrazolines, azetidin-2-one, antimicrobial activity.*

GJSFR-B Classification : *FOR Code: 030306p*



Strictly as per the compliance and regulations of :

© 2014. Shailesh H. Shah, Pankaj S. Patel, Mitesh Prajapati, Bharat Vaghari & Jayesh Parmar. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Synthesis, Characterization and Antimicrobial Activity of Azetidin-2-One based Phenyl Sulfonyl Pyrazoline Derivatives

Shailesh H. Shah [α], Pankaj S. Patel [σ], Mitesh Prajapati [ρ], Bharat Vaghari [ω] & Jayesh Parmar [¥]

Abstract- **A new series of 4-(4-Chlorophenyl)-3-chloro-1-{4-[5-(Substituted phenyl)-1-(phenylsulfonyl) -4,5-dihydro-pyrazol-3-yl]phenyl}azetidin-2-one are synthesized by reacting 3-chloro-1-{4-[5-(Substituted phenyl)-4,5-dihydro-pyrazol-3-yl]phenyl}-4-(4-Chlorophenyl)azetidin-2-one with Benzene sulfonyl chloride in presence of pyridine. All these compounds were characterized by means of their IR, 1H NMR, Spectroscopic data and were tested for their antibacterial and antifungal activities by broth dilution method.**

Keywords: chalcones, phenyl sulfonyl-pyrazolines, azetidin-2-one, antimicrobial activity.

I. Introduction

Heterocyclic compounds are the well-known class of compounds for its Biological applications. Keeping in view the importance of these Biological activities, it was considered of interest to synthesize some new phenyl sulfonyl derivatives of pyrazoles.

Pyrazolines are heterocyclic compounds which posses wide range of interesting biological activities such as anti-inflammatory [1], insecticidal [2], anti-tubercular [3], antitumor [4], tranquilizing [5], immunosuppressive [6], diuretic [7], anticonvulsant [8], antifungal [9], antidepressant activities [10-11], antibacterial activities [12], molluscidal [13]. In the present communication, we report the reaction of 3-chloro-1-{4-[5-(Substituted phenyl)-4, 5-dihydro-pyrazol-3-yl] phenyl}-4-(4-Chlorophenyl) azetidin-2-one with Benzene sulfonyl chloride in presence of pyridine to form phenyl sulfonyl Pyrazoline (5a-j). The structures of the various synthesized compounds were assigned on the basis of IR, 1H-NMR spectral data and elemental analysis. These compounds were also screened for their antimicrobial activity.


II. Experimental

The IR spectra were recorded on IR affinity-1, DRS-8000A, Shimadzu, Ptc. Ltd., Japan spectrophotometer. The 1H-NMR was recorded in DMSO on Bruker Advance II 400 MHz spectrometer using TMS as an internal standard. Melting points were determined in open capillary tubes and are uncorrected. The purity of the compounds was checked by TLC-using Silica gel-G (Merck). Column chromatography was performed on silica gel. All the compounds were tested for their antibacterial and antifungal activities by broth dilution method

Author α: Department of Chemistry, Patel JBR Arts, Patel AMR Commerce & Patel JDKD Science College, Borsad.
e-mail: shailchem@yahoo.com
Author σ: Department of Chemistry, Sheth L.H. Science College, Mansa, Gujarat. e-mail: pspatel_mansa@yahoo.co.in
Authors ρ ω ¥: Students of Patel JBR Arts, Patel AMR Commerce & Patel JDKD Science College, Borsad, Gujarat-India.

© 2014 Global Journals Inc. (US)

Reaction Scheme



a) Preparation of 1-(4-{[(4-Chlorophenyl) methylene] amino} phenyl) ethanone (1)

A mixture of 4-Chloro Benzaldehyde (0.01M), 1-(4-aminophenyl) ethanone (0.01M) and methanol (30ml) was heated for about 5 min. in a beaker (250 ml) to get a clear solution. The solution was kept overnight at room temperature to get the respective crude solid which was recrystallized from ethanol to obtain the pure crystals of 1-(4-{[(4-chlorophenyl) methylene] amino} phenyl) ethanone respectively. The yield of the product was 75% and the product melts at 1200C. Found: C(69.88%) H(4.65%) N(5.41%) , Calcd. for C15H12ClNO: C(69.91%) H(4.69%) N(5.43%). IR, cm-1: 3084(=C-H), 2922(-C-H), 1678(>C=O), 1628(>C=N-), 1595 (>C=C<), 1408(-CH3, bend), 1301(-C-N<), 1240(-C-CO-C-), 738(-C-Cl). 1H-NMR (DMSO, d, ppm): 2.5785 (3H, s, COCH3), 6.5144-7.7992 (8H, m, Ar-H), 8.803 (1H, s, -CH=N-).

b) Preparation of 1-(4-acetylphenyl)-3-chloro-4-(4-Chlorophenyl) azetidin-2-one (2)

In a 100ml Round bottom flask 1-(4-{[(4-Chloro phenyl) methylene] amino} phenyl) ethanone (0.01M) in

© 2014 Global Journals Inc. (US)

70ml benzene was taken. Chloro acetyl chloride (0.01M) was added at room temperature with constant stirring and triethylamine 1ml was added and the reaction mixture was refluxed for 7 hours. After the completion of reaction, solvent was removed by vacuum distillation. The solid was filtered, dried and recrystallized from toluene. The yield of the product was 60% and the product melts at 1080C. Found: C(61.07%) H(3.88%) N(4.17%), Calcd. for C17H13Cl2NO2: C(61.10%) H(3.92%) N(4.19%). IR, cm-1: 3041(=C-H), 2921(-C-H), 1712(>C=O), 1548(>C=C<), 1365(-CH3, bend), 1292(-C-N<), 1197(-C-CO-C-), 642(-C-Cl). 1H-NMR (DMSO, d, ppm): 2.5550 (3H, s, COCH3), 4.8102 (1H, d, >CH-Ar), 5.4594 (1H, d, >CH-Cl), 7.3170-8.0618 (8H, m, Ar-H).

c) *Preparation of 3-chloro-1-{4-[3-(Substituted phenyl) prop-2-enoyl] phenyl}-4-(4-Chlorophenyl) azetidin-2-one (3a-j)*

To the solution of 1-(4-acetylphenyl)-3-chloro-4-(4-Chloro phenyl) azetidin-2-one (0.01M) in absolute ethanol (50 ml), substituted Benzaldehyde (0.01M) and 2% NaOH were added and refluxed for 10 hours. After refluxing the reaction mixture was concentrated, cooled, filtered and neutralized with dil. HCl. The solid residue thus obtained was crystallized by absolute ethanol. IR(3d), cm-1: 3043(=C-H), 1722(>C=O), 1624 (>C=C<), 1451(-N=O), 1286(-C-N<), 1232 (-C-O-), 684(-C-Cl). 1H-NMR (3g-DMSO, d, ppm): 4.8757 (1H, d, >CH-Ar), 5.4224 (1H, d, >CH-Cl), 6.3621-8.5674 (12H, m, Ar-H), 7.9978 (2H, d, -CH=CH-), 9.9660 (1H, s, Ar-OH).

d) *Preparation3-chloro-1-{4-[5-(Substituted phenyl)-4,5-dihydro-pyrazol-3-yl]phenyl}-4-(4-Chlorophenyl)azetidin-2-one.(4a-j)*











A mixture of 3-chloro-1-{4-[3-(Substituted phenyl) prop-2-enoyl] phenyl}-4-(4-Chlorophenyl) azetidin-2-one (0.01M) and 99% hydrazine hydrate (0.015M) in ethanol (50ml) refluxed gently for 3 hours. Then the mixture was concentrated and allowed to cool. The resulting solid was filtered, washed with ethanol and recrystallized from ethanol to give a pale brown solid. IR(4e), cm-1: 3041 (=C-H), 2931(-C-H), 1728(>C=O), 1643(>C=N-), 1552 (>C=C<), 1452(-CH2, bend), 1313(-C-N<),1290 (-N-N), 663 (-C-Cl-),. 1H-NMR (4b-DMSO, δ, ppm): 3.61 (2H, d, CH2-of Pyrazol), 4.33 (1H, t, >CH-Ar of Pyrazol), 4.80 (1H, d, >CH-Ar of Azetidine), 5.32 (1H, d, >CH-Cl of Azetidine), 6.56-7.92 (13H, m, Ar-H), 9.65 (1H, s, Ar-OH).

e) *Preparation of 4-(4-Chlorophenyl)-3-chloro-1-{4-[5-(Substituted phenyl)-1-(phenyl sulfonyl)-4, 5 -dihydro-pyrazol-3-yl] phenyl} azetidin-2-one (5a-j)*

A solution of 3-chloro-1-{4-[5-(Substituted phenyl)-4, 5-dihydro-pyrazol-3-yl] phenyl}-4-(4-Chlorophenyl) azetidin-2-one (0.001M) in dry pyridine (25ml) cooled in an ice-bath and to it benzene sulfonyl chloride (0.0011M) was added. The mixture was stirred for 1 hour at room temperature and was then treated with cold dilute HCl (2N). The resulting solid was filtered, washed with water, and recrystallized from absolute ethanol IR(5f), cm-1: 3040(=C-H), 1730(>C=O), 1631(>C=N-), 1529 (>C=C<), 1452(-CH2,bend), 1375 (CH3,bend), 1313 (-C-N-), 1230(-N-N), 696 (-C-Cl-), 1133 (>S=O). 1H-NMR (5e-DMSO, δ, ppm): 3.9 (2H, d, CH2- of Pyrazol), 4.3 (1H, t, >CH-Ar of Pyrazol), 4.8 (1H, d, >CH-Ar of Azetidine), 5.6 (1H, d, >CH-Cl of Azetidine), 6.9-8.0 (17H, m, Ar-H).

© 2014 Global Journals Inc. (US)

Table 1 : Physical constant of 4-(4-Chlorophenyl)-3-chloro-1-{4-[5-(Substituted phenyl)-1-(phenyl sulfonyl)-4, 5 -dihydro-pyrazol-3-yl] phenyl} azetidin-2- one

Compd	R	M.F.	Yield %	M.P. °C	Elemental Analysis		
					% C Found (Calcd)	% N Found (Calcd)	% H Found (Calcd)
5a		$C_{30}H_{22}Cl_3N_3O_3S$	66	65	58.93 (58.98)	6.84 (6.88)	3.61 (3.63)
5b		$C_{30}H_{23}Cl_2N_3O_4S$	65	82	60.76 (60.81)	7.03 (7.09)	3.88 (3.91)
5c		$C_{32}H_{27}Cl_2N_3O_5S$	63	79	60.33 (60.38)	6.56 (6.60)	4.22 (4.28)
5d		$C_{30}H_{22}Cl_2N_4O_5S$	67	100	57.93 (57.98)	8.98 (9.01)	3.54 (3.57)
5e		$C_{30}H_{22}Cl_3N_3O_3S$	62	78	58.92 (58.98)	6.84 (6.88)	3.59 (3.63)
5f		$C_{34}H_{32}Cl_2N_4O_3S$	68	80	63.02 (63.06)	8.61 (8.65)	4.92 (4.98)
5g		$C_{30}H_{23}Cl_2N_3O_4S$	70	108	60.78 (60.81)	7.04 (7.09)	3.88 (3.91)
5h		$C_{32}H_{28}Cl_2N_4O_3S$	64	145	62.00 (62.03)	9.01 (9.04)	4.51 (4.56)
5i		$C_{30}H_{23}Cl_2N_3O_3S$	71	138	62.46 (62.50)	7.25 (7.29)	3.98 (4.02)
5j		$C_{31}H_{25}Cl_2N_3O_5S$	69	75	59.78 (59.81)	6.71 (6.75)	4.01 (4.05)

III. Results and Discussion

a) Antimicrobial activity

The MICs of synthesized compounds were carried out by broth micro dilution method as described by Rattan (2000). It is one of the non automated in vitro bacterial susceptibility tests. This classic method yields a quantitative result for the amount of antimicrobial agents that is needed to inhibit growth of specific microorganisms.

The invitro antimicrobial activity of test compounds were assessed against 24 hr cultures of several selected bacteria and fungi. The bacteria used were E. coli, S. aureus, P. aeruginosa, and S. pyogenus; the fungi used were C. albicans, A. Niger, and A. clavatus.

The antimicrobial activity was performed by broth dilution method in DMSO. Gentamycin, Ampicilin, Chloramphenicol, Ciprofloxacin, Norfloxacin, Nystatin

© 2014 Global Journals Inc. (US)

and Greseofulvin were used as standard for the evaluation of antibacterial and antifungal activities respectively. The activity was reported by Minimal Inhibition Concentration. The results are summarized in Table-2

Table 2 : Antimicrobial activity 4-(4-Chlorophenyl)-3-chloro-1-{4-[5-(Substituted phenyl)-1-(phenyl sulfonyl)-4, 5 -dihydro-pyrazol-3-yl] phenyl} azetidin-2-one

SR. NO.	COMP. NO.	R	ANTIBACTERIAL ACTIVITY MINIMAL INHIBITION CONCENTRATION				ANTIFUNGAL ACTIVITY MINIMAL INHIBITION CONCENTRATION		
			E.COLI	P.AERUGINOSA	S.AUREUS	S.PYOGENUS	C.ALBICANS	A.NIGER	A.CLAVATUS
			MTCC 443	MTCC 1688	MTCC 96	MTCC 442	MTCC 227	MTCC 282	MTCC 1323
1	5a	-2-Cl	200	250	200	62.5	1000	>1000	>1000
2	5b	-2-OH	200	250	125	250	1000	1000	1000
3	5c	-3-OCH_3, -4-OCH_3	175	225	200	125	800	800	800
4	5d	-3-NO_2	200	200	225	200	1000	1000	**>1000**
5	5e	-4-Cl	100	125	100	250	>1000	>1000	>1000
6	5f	-4-$N(C_2H_5)_2$	125	200	62.5	100	1000	1000	1000
7	5g	-4-OH	200	200	175	250	800	1000	800
8	5h	-4-$N(CH_3)_2$	225	225	150	200	>1000	800	700
9	5i	-H	175	225	200	150	1000	1000	1000
10	5j	-3-OCH_3, -4-OH	100	125	100	250	500	>1000	1000

Table 3 : Antibacterial Activity: Minimal Inhibition Concentration (The Standard Drugs)

DRUG	E.COLI	P.AERUGINOSA	S.AUREUS	S.PYOGENUS
-	MTCC 443	MTCC 1688	MTCC 96	MTCC 442
(MICROGRAMME/ML)				
GENTAMYCIN	0.05	1	0.25	0.5
AMPICILLIN	100	--	250	100
CHLORAMPHENICOL	50	50	50	50
CIPROFLOXACIN	25	25	50	50
NORFLOXACIN	10	10	10	10

Table 4 : Antifungal Activity: Minimal Inhibition Concentration (The Standard Drugs)

DRUG	C.ALBICANS	A.NIGER	A.CLAVATUS
-	MTCC 227	MTCC 282	MTCC 1323
(MICROGRAMME/ML)			
NYSTATIN	100	100	100
GRESEOFULVIN	500	100	100

Biological screening result of activities 4-(4-Chlorophenyl)-3-chloro-1-{4-[5-(Substituted phenyl)-1-(phenyl sulfonyl)-4, 5 -dihydro-pyrazol-3-yl] phenyl} azetidin-2-one based derivatives shows that compound 5e & 5j have shown better activity against E. coli. Compound 5f show good to very good activity against S. pyogenus, while rest of all compound possessed good activity against S.aureus in the range of 62.5-225 μg/ml. Compound 5j is found to be good antifungal activity against C. albicans, against standard drugs Greseofulvin. While rest of all derivatives are poor against A. Niger, and A.clavatus.

IV. Conclusion

The Main focus of this research work was to synthesize, characterize and evaluate antimicrobial activities of the newly synthesized phenyl sulfonyl Pyrazoline derivatives, structures of synthesized compounds were confirmed and characterized with the help of analytical data's such as IR and 1H-NMR. In summary, we have described the synthesis and antimicrobial activity of some new 4-(4-Chlorophenyl)-3-chloro-1-{4-[5-(Substituted phenyl)-1-(phenyl sulfonyl) -4, 5 -dihydro-pyrazol-3-yl] phenyl} azetidin-2-one MIC values revealed that amongst newly synthesized

© 2014 Global Journals Inc. (US)

compound having Methoxy-Hydroxide type linkage has shown good activity against the bacterial strains.

V. Acknowledgement

Authors are thankful to UGC. Ganeshkhind, Pune for Teacher Research Fellowship F.No. 47-180/12 (WRO) The authors are thankful to the Principal and Management of Patel JB Rudelwala Arts, Patel AM Rudelwala Commerce & Patel JDK Davolwala Science College, Borsad for providing laboratory facilities, SAIF, Chandigarh for NMR Spectra and Loyola Research Center- Xavier's College, Ahmedabad for IR spectra and Micro-care laboratory, Surat, Gujarat, India for biological activity.

References Références Referencias

1. S. Sridhar., SC Dinda., Y Rajendra Prasad, Indian J. Chem. Sci. 8(4), 2010, 2697-2707.
2. SS Kristophar and MS David. Pestic biochem and Physiol, 81, 2005, 136.
3. VH Babu., SK Manna., KK Srinivas and GV Bhat. Indian J. Heterocycle Chem. 13, 2004, 253, Chem Absr, 141, 2004, 314227b.
4. EC Taylor., H Patel and H Kumar. Tetrahedron, 48, 1992, 8089.
5. H Brudecer., R Richle and R Ruegg. (Hoffman-la-Roche, Inc.) U.S. 3, 822, 283 (Cl. 260-310R, C07d) 02 Jul (1974) Appl. 206-691'11Oct (1972) 9, Chem. Abstd. Pp-81, 105495 (1974).
6. JG Lombardino and IG Otterness. J.Med.Chem. 24, 830 (1981).
7. Z Brzozowski, Z Kaminski and S Angielski. Acta Pol.Pharma, 36(6), 645 (1979), Chem. Abstr., 93, 204525e (1980).
8. AVK Srivatava and A Kumar. Arzneim Foresch, 52, 2002, 787, Chem.Abstr. 138, 2003, 3537584.
9. SS Korgaokar, PH Patil, MT Shah and HH Parekh, Indian J. Pharma Sci., 58, 1996, 222.
10. O Ruhogia, Z Oxdemir, V Calis, B Gumuses and AA Bilgin, Arzneim Foresch, 55, 2005,431.
11. R MallikarjunaRao, J Sreeramulu, LK Ravindranath, G NagarajaReddy, K Hanumanthurayudu, GN Reddy, A Jayaraju, P. Madhusudhan, Journal of Chemical and Pharmaceutical Research, 4(1):272-278 (2012).
12. PV Badadh, NM Chavan, PG Mandhane, Indian J. of Chem., 50B, June 2011, pp 879-884.
13. NB Colthup, LH Daly and SE Wiberly, Introduction to Infrared and Raman Spectroscopy, Academic Press, New York (1964).

© 2014 Global Journals Inc. (US)

Global Journal of Science Frontier Research: B
Chemistry
Volume 14 Issue 1 Version 1.0 Year 2014
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Elimination of Methylene Blue Dye with Natural Adsorbent « Banana Peels Powder »

By Faisal Moubarak, Rachid Atmani, Ibtissam Maghri, M'hamed Elkouali, Mohammed Talbi, Mona Latifa Bouamrani, Mohammed Salouhi & Abdelkbir Kenz

University Hassan II, Morocco

Abstract- The investigation focused on the adsorption of the blue methylene on banana peels powder.

The studies led to determine contacting time (40 minutes for 90% elimination of substrate methylene blue). However, this contacting time can vary with dye concentration. The optimization of parameter such as the temperature, agitation and initial concentration of dye has showed a sensitive improvement of the capacity and rate adsorption. Moreover, the adsorption of this compound is well described by both Langmuir and Freundlich isotherm models.

Keywords: *adsorption, blue methylene, dye, isotherm models, banana peels.*

GJSFR-B Classification : *FOR Code: 259999*



Strictly as per the compliance and regulations of :

© 2014. Faisal Moubarak, Rachid Atmani, Ibtissam Maghri, M'hamed Elkouali, Mohammed Talbi, Mona Latifa Bouamrani, Mohammed Salouhi & Abdelkbir Kenz. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Elimination of Methylene Blue Dye with Natural Adsorbent « Banana Peels Powder »

Faisal Moubarak [α], Rachid Atmani [σ], Ibtissam Maghri [ρ], M'hamed Elkouali [ω], Mohammed Talbi [¥], Mona Latifa Bouamrani [§], Mohammed Salouhi [χ] & Abdelkbir Kenz [ν]

Abstract- The investigation focused on the adsorption of the blue methylene on banana peels powder.

The studies led to determine contacting time (40 minutes for 90% elimination of substrate methylene blue). However, this contacting time can vary with dye concentration. The optimization of parameter such as the temperature, agitation and initial concentration of dye has showed a sensitive improvement of the capacity and rate adsorption. Moreover, the adsorption of this compound is well described by both Langmuir and Freundlich isotherm models.

Keywords: adsorption, blue methylene, dye, isotherm models, banana peels.

I. Introduction

Intensive use of colors in everyday life has created a lot of problems both in the environment and in food [1]. It is important to mention that the water pollution is due to discharges of textile industries, and in feed, the toxicity is due to incorporation of several synthetic dyes [2] .

To address this situation, several decontamination methods have been developed, we cite as examples the adsorption [3,4], ion exchange [5,6], coagulation - flocculation [7] etc. In this work, we present the results for the adsorption blue methylene in banana peels powder. We determined successively contact time and the influence of parameters such as particle size, the initial concentration of substrate, the mass of the adsorbent and the temperature. These studies are needed to better understand the mechanisms governing the adsorption process.

II. Theoretical and Experimental Part

The adsorbent used in this work, banana peels powder was first washed with distilled water to remove impurities and then dried in air for 15 days , then ground in mortar and analyzed by size in order to obtain two different types of fractions, the first fraction is characterized by a diameter less than 80 microns (d < 80 microns), and the second fraction has a diameter of between 80 microns and 2 mm (80 microns < d < 2mm) .

The measurements and spectral evolution of the optical density of the solutions of methylene blue at different reaction times were followed by UV/visible spectrometry (ShimadzuUV mini - 1240)

The experiments were conducted in "batch method" (100 ml Erlenmeyer flask) at an ambient temperature (22°C). It should be noted that the temperature control was made using a digital thermometer.

Moreover, to ensure good dispersion of solid particles Banana peels powder, we adopted the value of 1g/l (0.1g/100 ml) for solid/liquid ratio.

To investigate the adsorption power of our support, we applied models Langmuir and Freundlich. Due to their simplicity, the kinetic models the most commonly used are those of Langmuir and Freundlich.

a) Adsorption Isotherms

The adsorption data from experiments were fitted with:

i. *Langmuir isotherm [8]*

Langmuir adsorption isotherm sheds no light on the mechanistic aspects of adsorption. It provides information on uptake capabilities and also reflects the usual equilibrium process behaviour The Langmuir non linear equation is:

$$q_e = q_m \cdot \frac{b.C_e}{1 + b.C_e} \quad (1)$$

Where q_e is the amount of methylene blue sorbed per unit weight of biomass at equilibrium, C_e is the residual equilibrium Dye, q_m is the maximum possible amount of dye adsorbed per unit weight of biomass and b is the equilibrium constant related to the affinity of the binding sites for the methylene blue, lower is b more is the affinity of methylene blue to biomass.

Low values of parameter b indicate that banana peel powder have high affinity for methylene blue. The table 1 presents linear equations of Langmuir model.

The essential characteristic of a Langmuir isotherm can be expressed as a dimensionless constant, defined as the separation factor:

Authors α σ ρ ω ¥ § ν: Faculty of Sciences Ben M'Sik, Avenue Cdt Driss Elharti, BP: 20702 , Casablanca, Morocco.
e-mails: moubarak46@hotmail.fr, atmanirachid12@gmail.com, maghri.ibtissam@gmail.com, m.elkouali@gmail.com, maarifcentre@yahoo.fr, a.kenz@voila.fr, mlbouamrani@gmail.com
Author χ: Laboratory of bio systemes science and ingenierie, EMI, Rabat Agdal, Morocco. e-mail: m.salouhi@gmail.com

© 2014 Global Journals Inc. (US)

$$R_L = \frac{1}{1 + b.C_0} (2)$$

b) Freundlich isotherm [9]

The Freundlich non linear equation is:

$$q_e = K_F C^n e (3)$$

Where, K_F and n are constants indicating adsorption capacity and adsorption intensity, respectively. The constants were obtained from the plots of the linearized equations:

$$\log q_e = \log K_F + n.\log C_e (4)$$

Another use of the results is to plot the variation of the distribution coefficient K_d as a function of q_e in logarithmic scale:

$$\log K_d = (\frac{1}{n}).\log K_F . + [\frac{(n-1)}{n})].(\log q_e) (5)$$

III. Results and Discussions

a) Optimization of methylene blue adsorption

i. *Determination of equilibrium time*

The study of the adsorption of methylene blue on banana peels powder (PBP), clearly implies the determination of contact time, time which corresponds to the adsorption equilibrium or a state of saturation of the support the substrate . In this case, the experimental procedure is simple " batch method" which involves 10 mg / l of methylene blue with 1g / l of powdered banana peels . Analysis by UV/visible allows us to determine the residual concentrations of the substrate at different reaction times.

Thus, the determination of the equilibrium time, allowed the establishment of adsorption isotherms that are essential for the calculation of the maximum adsorption capacity and to identify the type of adsorption to occur either in mono or multilayers. The results obtained at the end of these experiments showed that the contact time required for the elimination of 90 % of methylene blue is 40 minutes. Furthermore, the extension of the time to (90 minutes) does not lead to a significant improvement of the percentage removal. This justifies the inclusion of this contact time for other adsorption experiments (Figure 1).This result was obtained by the ratio :

$$q = V.\frac{C_0 - C_t}{m} (6)$$

q: Adsorption capacity of the support (mg/g).
C_0: Initial concentration of dye (mg/l).
C_t: Concentration of dye (mg/l) at time t of the adsorption process .
V: Volume of solution (substrate).
m: Mass of support (g).

Other factors such as the particle size, the initial concentration of the substrate, the mass of the adsorbent and the temperature may affect the adsorption capacity [10]. It finds its application in various adsorption models used.

b) Effect of particle size

To study the influence of the particle size of the adsorbent We choose two types of fraction, the first fraction having a particle size less than 80 microns, and the second fraction is between 80 microns and 2 mm, results is shown in fig 2, it is found that the adsorption is fast and relatively large for the fine particles (d <80 microns), this could be explained by the fact that the extent of adsorption depends on the external surfaces of the particles, More the particle size is small over the exchange surfaces are provided important promoting high speed transfer of dye to the adsorbent. On considering this, we will continue work with fine particle (d <80 microns).

c) Effect of mass of adsorbent

The influence of the mass of the adsorbent was studied in the range 0.1- 0.3 g. The figure 3 shows the variation of the adsorption amount depending on the time chosen for different masses ; 0.1 g, 0.2 g and 0.3 g, the mass 0.1g provides removal rate which can reach 96%. This effect, we chose the mass 0.1 g, as an optimal weight for the rest of our work.

d) Effect of initial concentration

Figure 4 shows the influence of the initial concentration of methylene blue on the dye adsorption capacity (the mass of the support being fixed), it shows that the adsorption capacity is better for the high initial concentration of methylene blue.

e) Effect of temperature

Figure 5 shows the influence of temperature on the adsorption of the dye. When we use different temperatures to 40 minutes of stirring, the adsorption capacity of powdered banana peels over time increases, the values obtained were Q = 8.8991 mg / g at 22°C.

The experimental results show that this parameter does not affect the adsorption retention of methylene blue on the support of the study process.

f) Adsorption Isotherm

Adsorption isotherms play an important role in determining the maximum capacity and in determining the type of adsorption front is produire.Elles are obtained first by the knowledge of the time and then the graphical presentation of Qe = f (Ce) and Qe These are respectively the amount of adsorbed dye per g of adsorbent and the equilibrium concentration of this colorant.les experimental results obtained show that the S-type is isothermal, which classification coorspond

© 2014 Global Journals Inc. (US)

Gilles[11]. This indicates the adsorption of growth when the concentration of the adsorbate augmante (FIG. 6)

The shape obtained shows that the adsorption of the dye process could occur in a monolayer.

g) Langmuir isotherm

The values of equilibrium relation parameter, RL were calculated. As shown in Table 1, RL values lie between 0 and 1 which indicate favourable sorption isotherm for methylene blue. Low values of parameter b indicate that banana peels powder have high affinity for methylene blue.

h) Freundlich isotherm

The values of equilibrium relation parameter, n were calculated for two Freundlich model's, the results is shown in table 1, n values is high than 1 so the methylene blue adsorption was not well fitted with Freundlich model and The low values of maximum adsorption capacities obtained from the Freundlich model, confirm that the molecule of Methylene Blue is not strongly adsorbed inside the pores because of its size.

The main parameters characterizing each model are summarized in Table 1. The results shown in Figures 7 and 8 show that the adsorption of Methylene blue on banana peels powder (PBP) follow the two linear models Freundlich and Langmuir.

Table1 : Valeur des paramétres d'adsorption Qm,K et Kf du BM sur les Pelures de Bananes en Poudre

Models	Equation	Q_m (mg/g)	$K_{(l.mg^{-1})}$	K_f(mg/g)	R
Langmuir	$Q_e = \frac{Q_m\, KC_e}{1+KC_e}$ $\frac{1}{Q_e} = \frac{1+KC_e}{Q_m KC_e} = \frac{1}{Q_m KC_e} + \frac{1}{Q_m}$	111,111	0,310	-	0,972
Freundlich	$Q_e = KC_e^n$ $\ln(Q_e) = \ln K_f + n \ln(C_e)$	-	-	25,559	0,943




Fig. 1 : Adsorption kinetic of BM on banana peel powder, [BM]=20 mg/l ; m =1g/l ; T=22°C and d<80 μm

© 2014 Global Journals Inc. (US)




Fig. 2 : Effect of Granulometry banana peel powder, m=1g/l and T=22°C




Fig. 3 : Effect of banana peel powder mass. [BM]=20 mg/l ; T=22°C and d<80 μm




Fig. 4 : Effect of methylene blue initial Concentration, m =1g/l; T=22°C and d< 80μm

© 2014 Global Journals Inc. (US)




Fig. 5 : Effect of Temperature, [BM]= 20 mg/l. m 1g/l and< 80μm




Fig. 6 : Isotherm of methylene blue adsorption. C0=20 mg/l; m =1g/l; T=22 °C and d <80 μm




Fig. 7 : Langmuir linear Model of Methylene blue adsorption,T=22°C; d<80 μm

© 2014 Global Journals Inc. (US)




Fig. 8 : Freundlich Model of Methylene blue adsorption, T=22°C; d<80μm

IV. Conclusion

The results obtained in this work have to mount that :

- Methylene blue (20 mg/l) is adsorbed relatively well on this medium with a contact time of 40 minutes which corresponds to a removal rate of 90 %. Giving it a relatively large affinity with respect to peels Banana Powder (PBP).
- The effect of particle size, the mass of the adsorbent and the adsorbate concentration had a positive influence on the capacity and kinetics of retention Methylene Blue on our support study. Models of Langmuir - Freundlich and correctly describes the adsorption process.

References Références Referencias

1. T. sauer G. Gesconeto HJ.J and Moreira,R.F P.M. (2002), 149, 147-154.
2. N. Daneshver, D. Salari, A.R andkhateee, J. Photochem, 2003, Photobiol A :Chemistry, 157, 111-116.
3. G.S. Gupta,G. Prasad and V.N. Singh, 1988, J.Environ.Sci.Health,123, 205.
4. M. Mazet, O. Dussort, M. Roger and Dussoubes-Marmier, 1990, revue des sciences de l'eau, 3, 129.
5. F. Perineaux, J. Molinier and M.R. Ladisch, 1988, Enzym. Microb. Tech., 10, 632.
6. Y. Yang, C. Ladisch and M.R. Ladisch, 1988, Enzym. Microb. Tech., 10, 632.
7. Z. Lisheng and B. Dobias, 1992, Watertreatment, 7, 221-232.
8. I. Langmuir,1918, the adsorption of gases on plane surfaces of glass, mica and platinum, Journal of American Chemical Society, 40, P 1361-1403.
9. H. Freundlich,1909, Kapillarchemie. Akademische Verlagsgesesellschaft, Leipzig, Synthesis and catalysis, 4, 259-269.
10. E. L Slejko and M. Dekker, 1985, adsorption technologie; a step by step by step approch solutions to process evolutions and application. Edited M. Dekker, Newyork
11. C.H. Gilles, M. MC Evant, S.W. Nakhawas and D.J.J. Smith, 1960, Chem. Soc., 3973.

© 2014 Global Journals Inc. (US)

Global Journals Inc. (US) Guidelines Handbook 2014

www.GlobalJournals.org

FELLOWS

FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (FARSS)

Global Journals Incorporate (USA) is accredited by Open Association of Research Society (OARS), U.S.A and in turn, awards "FARSS" title to individuals. The 'FARSS' title is accorded to a selected professional after the approval of the Editor-in-Chief/Editorial Board Members/Dean.



- The "FARSS" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSS or William Walldroff, M.S., FARSS.

FARSS accrediting is an honor. It authenticates your research activities. After recognition as FARSB, you can add 'FARSS' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, and Visiting Card etc.

The following benefits can be availed by you only for next three years from the date of certification:

FARSS designated members are entitled to avail a 40% discount while publishing their research papers (of a single author) with Global Journals Incorporation (USA), if the same is accepted by Editorial Board/Peer Reviewers. If you are a main author or co-author in case of multiple authors, you will be entitled to avail discount of 10%.

Once FARSB title is accorded, the Fellow is authorized to organize a symposium/seminar/conference on behalf of Global Journal Incorporation (USA). The Fellow can also participate in conference/seminar/symposium organized by another institution as representative of Global Journal. In both the cases, it is mandatory for him to discuss with us and obtain our consent.





You may join as member of the Editorial Board of Global Journals Incorporation (USA) after successful completion of three years as Fellow and as Peer Reviewer. In addition, it is also desirable that you should organize seminar/symposium/conference at least once.

We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



© Copyright by Global Journals Inc.(US) | Guidelines Handbook

The FARSS can go through standards of OARS. You can also play vital role if you have any suggestions so that proper amendment can take place to improve the same for the benefit of entire research community.



As FARSS, you will be given a renowned, secure and free professional email address with 100 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.



The FARSS will be eligible for a free application of standardization of their researches. Standardization of research will be subject to acceptability within stipulated norms as the next step after publishing in a journal. We shall depute a team of specialized research professionals who will render their services for elevating your researches to next higher level, which is worldwide open standardization.



The FARSS member can apply for grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.
Once you are designated as FARSS, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria. After certification of all your credentials by OARS, they will be published on your Fellow Profile link on website https://associationofresearch.org which will be helpful to upgrade the dignity.

The FARSS members can avail the benefits of free research podcasting in Global Research Radio with their research documents. After publishing the work, (including published elsewhere worldwide with proper authorization) you can upload your research paper with your recorded voice or you can utilize chargeable services of our professional RJs to record your paper in their voice on request.



The FARSS member also entitled to get the benefits of free research podcasting of their research documents through video clips. We can also streamline your conference videos and display your slides/ online slides and online research video clips at reasonable charges, on request.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook



The FARSS is eligible to earn from sales proceeds of his/her researches/reference/review Books or literature, while publishing with Global Journals. The FARSS can decide whether he/she would like to publish his/her research in a closed manner. In this case, whenever readers purchase that individual research paper for reading, maximum 60% of its profit earned as royalty by Global Journals, will be credited to his/her bank account. The entire entitled amount will be credited to his/her bank account exceeding limit of minimum fixed balance. There is no minimum time limit for collection. The FARSS member can decide its price and we can help in making the right decision.

The FARSS member is eligible to join as a paid peer reviewer at Global Journals Incorporation (USA) and can get remuneration of 15% of author fees, taken from the author of a respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account.



MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (MARSS)

The ' MARSS ' title is accorded to a selected professional after the approval of the Editor-in-Chief / Editorial Board Members/Dean.



The "MARSS" is a dignified ornament which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., MARSS or William Walldroff, M.S., MARSS.

MARSS accrediting is an honor. It authenticates your research activities. After becoming MARSS, you can add 'MARSS' title with your name as you use this recognition as additional suffix to your status. This will definitely enhance and add more value and repute to your name. You may use it on your professional Counseling Materials such as CV, Resume, Visiting Card and Name Plate etc.

The following benefitscan be availed by you only for next three years from the date of certification.



MARSS designated members are entitled to avail a 25% discount while publishing their research papers (of a single author) in Global Journals Inc., if the same is accepted by our Editorial Board and Peer Reviewers. If you are a main author or co-author of a group of authors, you will get discount of 10%.

As MARSS, you will be given a renowned, secure and free professional email address with 30 GB of space e.g. johnhall@globaljournals.org. This will include Webmail, Spam Assassin, Email Forwarders,Auto-Responders, Email Delivery Route tracing, etc.



© Copyright by Global Journals Inc.(US) | Guidelines Handbook



We shall provide you intimation regarding launching of e-version of journal of your stream time to time.This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.

The MARSS member can apply for approval, grading and certification of standards of their educational and Institutional Degrees to Open Association of Research, Society U.S.A.





Once you are designated as MARSS, you may send us a scanned copy of all of your credentials. OARS will verify, grade and certify them. This will be based on your academic records, quality of research papers published by you, and some more criteria.

It is mandatory to read all terms and conditions carefully.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

AUXILIARY MEMBERSHIPS

Institutional Fellow of Global Journals Incorporation (USA)-OARS (USA)

Global Journals Incorporation (USA) is accredited by Open Association of Research Society, U.S.A (OARS) and in turn, affiliates research institutions as "Institutional Fellow of Open Association of Research Society" (IFOARS).



The "FARSC" is a dignified title which is accorded to a person's name viz. Dr. John E. Hall, Ph.D., FARSC or William Walldroff, M.S., FARSC.

The IFOARS institution is entitled to form a Board comprised of one Chairperson and three to five board members preferably from different streams. The Board will be recognized as "Institutional Board of Open Association of Research Society"-(IBOARS).

The Institute will be entitled to following benefits:

The IBOARS can initially review research papers of their institute and recommend them to publish with respective journal of Global Journals. It can also review the papers of other institutions after obtaining our consent. The second review will be done by peer reviewer of Global Journals Incorporation (USA) The Board is at liberty to appoint a peer reviewer with the approval of chairperson after consulting us.

The author fees of such paper may be waived off up to 40%.

The Global Journals Incorporation (USA) at its discretion can also refer double blind peer reviewed paper at their end to the board for the verification and to get recommendation for final stage of acceptance of publication.





The IBOARS can organize symposium/seminar/conference in their country on behalf of Global Journals Incorporation (USA)-OARS (USA). The terms and conditions can be discussed separately.

The Board can also play vital role by exploring and giving valuable suggestions regarding the Standards of "Open Association of Research Society, U.S.A (OARS)" so that proper amendment can take place for the benefit of entire research community. We shall provide details of particular standard only on receipt of request from the Board.



The board members can also join us as Individual Fellow with 40% discount on total fees applicable to Individual Fellow. They will be entitled to avail all the benefits as declared. Please visit Individual Fellow-sub menu of GlobalJournals.org to have more relevant details.

© Copyright by Global Journals Inc.(US) | Guidelines Handbook

We shall provide you intimation regarding launching of e-version of journal of your stream time to time. This may be utilized in your library for the enrichment of knowledge of your students as well as it can also be helpful for the concerned faculty members.



After nomination of your institution as "Institutional Fellow" and constantly functioning successfully for one year, we can consider giving recognition to your institute to function as Regional/Zonal office on our behalf.
The board can also take up the additional allied activities for betterment after our consultation.

The following entitlements are applicable to individual Fellows:



Open Association of Research Society, U.S.A (OARS) By-laws states that an individual Fellow may use the designations as applicable, or the corresponding initials. The Credentials of individual Fellow and Associate designations signify that the individual has gained knowledge of the fundamental concepts. One is magnanimous and proficient in an expertise course covering the professional code of conduct, and follows recognized standards of practice.

Open Association of Research Society (US)/ Global Journals Incorporation (USA), as described in Corporate Statements, are educational, research publishing and professional membership organizations. Achieving our individual Fellow or Associate status is based mainly on meeting stated educational research requirements.



Disbursement of 40% Royalty earned through Global Journals : Researcher = 50%, Peer Reviewer = 37.50%, Institution = 12.50% E.g. Out of 40%, the 20% benefit should be passed on to researcher, 15 % benefit towards remuneration should be given to a reviewer and remaining 5% is to be retained by the institution.

We shall provide print version of 12 issues of any three journals [as per your requirement] out of our 38 journals worth $ 2376 USD.

Other:

The individual Fellow and Associate designations accredited by Open Association of Research Society (US) credentials signify guarantees following achievements:

- The professional accredited with Fellow honor, is entitled to various benefits viz. name, fame, honor, regular flow of income, secured bright future, social status etc.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

- In addition to above, if one is single author, then entitled to 40% discount on publishing research paper and can get 10%discount if one is co-author or main author among group of authors.
- The Fellow can organize symposium/seminar/conference on behalf of Global Journals Incorporation (USA) and he/she can also attend the same organized by other institutes on behalf of Global Journals.
- The Fellow can become member of Editorial Board Member after completing 3yrs.
- The Fellow can earn 60% of sales proceeds from the sale of reference/review books/literature/publishing of research paper.
- Fellow can also join as paid peer reviewer and earn 15% remuneration of author charges and can also get an opportunity to join as member of the Editorial Board of Global Journals Incorporation (USA)
- • This individual has learned the basic methods of applying those concepts and techniques to common challenging situations. This individual has further demonstrated an in–depth understanding of the application of suitable techniques to a particular area of research practice.

Note :

"

- In future, if the board feels the necessity to change any board member, the same can be done with the consent of the chairperson along with anyone board member without our approval.
- In case, the chairperson needs to be replaced then consent of 2/3rd board members are required and they are also required to jointly pass the resolution copy of which should be sent to us. In such case, it will be compulsory to obtain our approval before replacement.
- In case of "Difference of Opinion [if any]" among the Board members, our decision will be final and binding to everyone.

"

Process of submission of Research Paper

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (*.DOC,*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission.Online Submission: There are three ways to submit your paper:

(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.

(II) Choose corresponding Journal.

(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.

(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.

(C) If these two are not conveninet , and then email the paper directly to dean@globaljournals.org.

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.

© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Preferred Author Guidelines

MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11'"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

You can use your own standard format also.
Author Guidelines:

1. General,

2. Ethical Guidelines,

3. Submission of Manuscripts,

4. Manuscript's Category,

5. Structure and Format of Manuscript,

6. After Acceptance.

1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global

Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.

2) Drafting the paper and revising it critically regarding important academic content.

3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.

Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.



To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

5.STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also.Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

Papers: These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

(a)Title should be relevant and commensurate with the theme of the paper.

(b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.

(c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.

(d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.

(e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.

(f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;

(g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.

(h) Brief Acknowledgements.

(i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.

The Editorial Board reserves the right to make literary corrections and to make suggestions to improve briefness.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

Format

Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than 1.4 × 10-3 m3, or 4 mm somewhat than 4 × 10-3 m. Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

Abstract, used in Original Papers and Reviews:

Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art.A few tips for deciding as strategically as possible about keyword search:

- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

Acknowledgements: Please make these as concise as possible.

References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

Tables, Figures and Figure Legends

Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.

Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.

Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.

Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.

6. AFTER ACCEPTANCE

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

6.1 Proof Corrections

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

www.adobe.com/products/acrobat/readstep2.html. This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at dean@globaljournals.org within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

6.3 Author Services

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

6.4 Author Material Archive Policy

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

6.5 Offprint and Extra Copies

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: editor@globaljournals.org .



Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

1. Choosing the topic: In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

2. Evaluators are human: First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

3. Think Like Evaluators: If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

4. Make blueprints of paper: The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

5. Ask your Guides: If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

6. Use of computer is recommended: As you are doing research in the field of Computer Science, then this point is quite obvious.

7. Use right software: Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

8. Use the Internet for help: An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

9. Use and get big pictures: Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

10. Bookmarks are useful: When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

11. Revise what you wrote: When you write anything, always read it, summarize it and then finalize it.

12. Make all efforts: Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

13. Have backups: When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

14. Produce good diagrams of your own: Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

15. Use of direct quotes: When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

16. Use proper verb tense: Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

17. Never use online paper: If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

18. Pick a good study spot: To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

19. Know what you know: Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

20. Use good quality grammar: Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

21. Arrangement of information: Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

22. Never start in last minute: Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

23. Multitasking in research is not good: Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

24. Never copy others' work: Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

25. Take proper rest and food: No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

26. Go for seminars: Attend seminars if the topic is relevant to your research area. Utilize all your resources.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

27. Refresh your mind after intervals: Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

28. Make colleagues: Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

29. Think technically: Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

30. Think and then print: When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

31. Adding unnecessary information: Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

32. Never oversimplify everything: To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

33. Report concluded results: Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

34. After conclusion: Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium though which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

Informal Guidelines of Research Paper Writing

Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.

Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

General style:

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

· Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

· Use standard writing style including articles ("a", "the," etc.)

· Keep on paying attention on the research topic of the paper

· Use paragraphs to split each significant point (excluding for the abstract)

· Align the primary line of each section

· Present your points in sound order

· Use present tense to report well accepted

· Use past tense to describe specific results

· Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives

· Shun use of extra pictures - include only those figures essential to presenting results

Title Page:

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript--must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for briefness. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

Approach:

- Single section, and succinct
- As a outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an conceptual must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.

- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

Procedures (Methods and Materials):

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

Methods:

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

Approach:

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

What to keep away from

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

Results:

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently.You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.

Administration Rules Listed Before Submitting Your Research Paper to Global Journals Inc. (US)

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

Segment Draft and Final Research Paper: You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptive of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.

- Do not give permission to anyone else to "PROOFREAD" your manuscript.

- Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.



© Copyright by Global Journals Inc.(US)| Guidelines Handbook

Criterion for Grading a Research Paper (Compilation) by Global Journals Inc. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

Topics	Grades		
	A-B	**C-D**	**E-F**
Abstract	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form Above 200 words	No specific data with ambiguous information Above 250 words
Introduction	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
Methods and Procedures	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
Result	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
Discussion	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
References	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring

INDEX

A

Aeruginosa · 46
Alcoholysis · 13

B

Benzylamino · 11, 23

F

Freundlich · 50, 54, 55, 58,

G

Greseofulvin · 47

M

Molluscidal · 42
Monochromator · 4

N

Norfloxacin · 46
Nystatin · 46

P

Phenolphthalein · 37
Pyrazoline · 42, 43, 44, 46, 47, 49

S

Spherulite · 1

T

Triethylamine · 44
Trifluoroacetic · 3
Tungabhadra · 31, 33, 34, 35, 36, 37, 38, 39, 40, 41



Global Journal of Science Frontier Research

Visit us on the Web at www.GlobalJournals.org | www.JournalofScience.org
or email us at helpdesk@globaljournals.org

ISSN 9755896



© Global Journals