Online ISSN: 2249-4626
Print ISSN: 0975-5896

Global Journal of Science Frontier Research: F

Mathematics and Decision Sciences

Highlights

- Kinematic Wave Theory
- Manifolds Admitting Special
- Coupled Oscillator Systems
- Optimal Discounted Portfolio

Discovering Thoughts, Inventing Future

VOLUME 13 ISSUE 7 VERSION 1.0

© 2001-2013 by Global Journal of Science Frontier Research , USA

Global Journal of Science Frontier Research: F
Mathematics & Decision Sciences

Global Journal of Science Frontier Research: F
Mathematics & Decision Sciences
Volume 13 Issue 7 (Ver. 1.0)

Open Association of Research Society

© Global Journal of Science Frontier Research .2013.

All rights reserved.

This is a special issue published in version 1.0 of "Global Journal of Science Frontier Research." By Global Journals Inc.

All articles are open access articles distributed under "Global Journal of Science Frontier Research"

Reading License, which permits restricted use. Entire contents are copyright by of "Global Journal of Science Frontier Research" unless otherwise noted on specific articles.

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopy, recording, or any information storage and retrieval system, without written permission.

The opinions and statements made in this book are those of the authors concerned. Ultraculture has not verified and neither confirms nor denies any of the foregoing and no warranty or fitness is implied.

Engage with the contents herein at your own risk.

The use of this journal, and the terms and conditions for our providing information, is governed by our Disclaimer, Terms and Conditions and Privacy Policy given on our website http://globaljournals.us/terms-and-condition/menu-id-1463/

By referring / using / reading / any type of association / referencing this journal, this signifies and you acknowledge that you have read them and that you accept and will be bound by the terms thereof.

All information, journals, this journal, activities undertaken, materials, services and our website, terms and conditions, privacy policy, and this journal is subject to change anytime without any prior notice.

Incorporation No.: 0423089
License No.: 42125/022010/1186
Registration No.: 430374
Import-Export Code: 1109007027
Employer Identification Number (EIN):
USA Tax ID: 98-0673427

Global Journals Inc.

*(A Delaware USA Incorporation with "Good Standing"; **Reg. Number: 0423089**)*

Sponsors: Open Association of Research Society
Open Scientific Standards

Publisher's Headquarters office

Global Journals Inc., Headquarters Corporate Office, Cambridge Office Center, II Canal Park, Floor No. 5th, ***Cambridge (Massachusetts)***, Pin: MA 02141 United States
USA Toll Free: +001-888-839-7392
USA Toll Free Fax: +001-888-839-7392

Offset Typesetting

Open Association of Research Society, Marsh Road, Rainham, Essex, London RM13 8EU United Kingdom.

Packaging & Continental Dispatching

Global Journals, India

Find a correspondence nodal officer near you

To find nodal officer of your country, please email us at *local@globaljournals.org*

eContacts

Press Inquiries: *press@globaljournals.org*
Investor Inquiries: *investers@globaljournals.org*
Technical Support: *technology@globaljournals.org*
Media & Releases: *media@globaljournals.org*

Pricing (Including by Air Parcel Charges):

For Authors:
22 USD (B/W) & 50 USD (Color)
Yearly Subscription (Personal & Institutional):
200 USD (B/W) & 250 USD (Color)

Editorial Board Members (HON.)

John A. Hamilton,"Drew" Jr.,
Ph.D., Professor, Management
Computer Science and Software Engineering
Director, Information Assurance Laboratory
Auburn University

Dr. Henry Hexmoor
IEEE senior member since 2004
Ph.D. Computer Science, University at Buffalo
Department of Computer Science
Southern Illinois University at Carbondale

Dr. Osman Balci, Professor
Department of Computer Science
Virginia Tech, Virginia University
Ph.D.and M.S.Syracuse University, Syracuse, New York
M.S. and B.S. Bogazici University, Istanbul, Turkey

Yogita Bajpai
M.Sc. (Computer Science), FICCT
U.S.A.Email:
yogita@computerresearch.org

Dr. T. David A. Forbes
Associate Professor and Range Nutritionist
Ph.D. Edinburgh University - Animal Nutrition
M.S. Aberdeen University - Animal Nutrition
B.A. University of Dublin- Zoology

Dr. Wenying Feng
Professor, Department of Computing & Information Systems
Department of Mathematics
Trent University, Peterborough,
ON Canada K9J 7B8

Dr. Thomas Wischgoll
Computer Science and Engineering,
Wright State University, Dayton, Ohio
B.S., M.S., Ph.D.
(University of Kaiserslautern)

Dr. Abdurrahman Arslanyilmaz
Computer Science & Information Systems Department
Youngstown State University
Ph.D., Texas A&M University
University of Missouri, Columbia
Gazi University, Turkey

Dr. Xiaohong He
Professor of International Business
University of Quinnipiac
BS, Jilin Institute of Technology; MA, MS, PhD,. (University of Texas-Dallas)

Burcin Becerik-Gerber
University of Southern California
Ph.D. in Civil Engineering
DDes from Harvard University
M.S. from University of California, Berkeley & Istanbul University

Dr. Bart Lambrecht
Director of Research in Accounting and FinanceProfessor of Finance
Lancaster University Management School
BA (Antwerp); MPhil, MA, PhD (Cambridge)

Dr. Carlos García Pont
Associate Professor of Marketing
IESE Business School, University of Navarra
Doctor of Philosophy (Management), Massachusetts Institute of Technology (MIT)
Master in Business Administration, IESE, University of Navarra
Degree in Industrial Engineering, Universitat Politècnica de Catalunya

Dr. Fotini Labropulu
Mathematics - Luther College
University of ReginaPh.D., M.Sc. in Mathematics
B.A. (Honors) in Mathematics
University of Windso

Dr. Lynn Lim
Reader in Business and Marketing
Roehampton University, London
BCom, PGDip, MBA (Distinction), PhD, FHEA

Dr. Mihaly Mezei
ASSOCIATE PROFESSOR
Department of Structural and Chemical Biology, Mount Sinai School of Medical Center
Ph.D., Etvs Lornd University
Postdoctoral Training,
New York University

Dr. Söhnke M. Bartram
Department of Accounting and FinanceLancaster University Management SchoolPh.D. (WHU Koblenz)
MBA/BBA (University of Saarbrücken)

Dr. Miguel Angel Ariño
Professor of Decision Sciences
IESE Business School
Barcelona, Spain (Universidad de Navarra)
CEIBS (China Europe International Business School).
Beijing, Shanghai and Shenzhen
Ph.D. in Mathematics
University of Barcelona
BA in Mathematics (Licenciatura)
University of Barcelona

Philip G. Moscoso
Technology and Operations Management
IESE Business School, University of Navarra
Ph.D in Industrial Engineering and Management, ETH Zurich
M.Sc. in Chemical Engineering, ETH Zurich

Dr. Sanjay Dixit, M.D.
Director, EP Laboratories, Philadelphia VA Medical Center
Cardiovascular Medicine - Cardiac Arrhythmia
Univ of Penn School of Medicine

Dr. Han-Xiang Deng
MD., Ph.D
Associate Professor and Research Department Division of Neuromuscular Medicine
Davee Department of Neurology and Clinical NeuroscienceNorthwestern University Feinberg School of Medicine

Dr. Pina C. Sanelli
Associate Professor of Public Health
Weill Cornell Medical College
Associate Attending Radiologist
NewYork-Presbyterian Hospital
MRI, MRA, CT, and CTA
Neuroradiology and Diagnostic Radiology
M.D., State University of New York at Buffalo,School of Medicine and Biomedical Sciences

Dr. Roberto Sanchez
Associate Professor
Department of Structural and Chemical Biology
Mount Sinai School of Medicine
Ph.D., The Rockefeller University

Dr. Wen-Yih Sun
Professor of Earth and Atmospheric SciencesPurdue University Director National Center for Typhoon and Flooding Research, Taiwan University Chair Professor Department of Atmospheric Sciences, National Central University, Chung-Li, TaiwanUniversity Chair Professor Institute of Environmental Engineering, National Chiao Tung University, Hsin-chu, Taiwan.Ph.D., MS The University of Chicago, Geophysical Sciences
BS National Taiwan University, Atmospheric Sciences
Associate Professor of Radiology

Dr. Michael R. Rudnick
M.D., FACP
Associate Professor of Medicine
Chief, Renal Electrolyte and Hypertension Division (PMC)
Penn Medicine, University of Pennsylvania
Presbyterian Medical Center, Philadelphia
Nephrology and Internal Medicine
Certified by the American Board of Internal Medicine

Dr. Bassey Benjamin Esu
B.Sc. Marketing; MBA Marketing; Ph.D Marketing
Lecturer, Department of Marketing, University of Calabar
Tourism Consultant, Cross River State Tourism Development Department
Co-ordinator , Sustainable Tourism Initiative, Calabar, Nigeria

Dr. Aziz M. Barbar, Ph.D.
IEEE Senior Member
Chairperson, Department of Computer Science
AUST - American University of Science & Technology
Alfred Naccash Avenue – Ashrafieh

President Editor (HON.)

Dr. George Perry, (Neuroscientist)
Dean and Professor, College of Sciences
Denham Harman Research Award (American Aging Association)
ISI Highly Cited Researcher, Iberoamerican Molecular Biology Organization
AAAS Fellow, Correspondent Member of Spanish Royal Academy of Sciences
University of Texas at San Antonio
Postdoctoral Fellow (Department of Cell Biology)
Baylor College of Medicine
Houston, Texas, United States

Chief Author (HON.)

Dr. R.K. Dixit
M.Sc., Ph.D., FICCT
Chief Author, India
Email: authorind@computerresearch.org

Dean & Editor-in-Chief (HON.)

Vivek Dubey(HON.)
MS (Industrial Engineering),
MS (Mechanical Engineering)
University of Wisconsin, FICCT
Editor-in-Chief, USA
editorusa@computerresearch.org

Sangita Dixit
M.Sc., FICCT
Dean & Chancellor (Asia Pacific)
deanind@computerresearch.org

Suyash Dixit
(B.E., Computer Science Engineering), FICCTT
President, Web Administration and
Development , CEO at IOSRD
COO at GAOR & OSS

Er. Suyog Dixit
(M. Tech), BE (HONS. in CSE), FICCT
SAP Certified Consultant
CEO at IOSRD, GAOR & OSS
Technical Dean, Global Journals Inc. (US)
Website: www.suyogdixit.com
Email:suyog@suyogdixit.com

Pritesh Rajvaidya
(MS) Computer Science Department
California State University
BE (Computer Science), FICCT
Technical Dean, USA
Email: pritesh@computerresearch.org

Luis Galárraga
J!Research Project Leader
Saarbrücken, Germany

Contents of the Volume

i. Copyright Notice
ii. Editorial Board Members
iii. Chief Author and Dean
iv. Table of Contents
v. From the Chief Editor's Desk
vi. Research and Review Papers

1. Lorentzian Para Sasakian Manifolds Admitting Special Semi Symmetric Recurrent Metric Connection. *1-6*
2. Global Attractor of the Nonlinearfour Order Wave Equations. *7-16*
3. On Semi-Open Sets and Semi-Separability. *17-20*
4. Lie Algebraic Approach and Complex Invariant Coupled Oscillator Systems. *21-27*
5. Sensitivity Analysis for the Physiographic Parameters of Overland as well as Cannel Flow Elements Based on Kinematic Wave (Kw) Theory. *29-42*
6. Optimal Discounted Portfolio, Expected Wealth and Strategic Consumption for a Defined Contribution Pension Scheme. *43-57*
7. On the Right Perfect Ring. *59-63*

vii. Auxiliary Memberships
viii. Process of Submission of Research Paper
ix. Preferred Author Guidelines
x. Index

Global Journal of Science Frontier Research
Mathematics and Decision Sciences
Volume 13 Issue 7 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Lorentzian Para Sasakian Manifolds Admitting Special Semi Symmetric Recurrent Metric Connection

By Sunil Kumar Srivastava & R. P. Kushwaha
D.D.U Gorakhpur University, India

Abstract - Several author as Agashe and Chafle [1], Sengupta, De. U.C, Binh [4] and many other introduced semi symmetric non metric connection in different way. In this paper we have studied LP sasakian manifold with special semi-symmetric recurrent metric connection [2] and discuss it exientance in LP sasakian manifold. In section 3 we establish the relation between the Riemannian connection and special semi-symmetric recurrent metric connection on LP sasakian manifold [4]. The section 4 deals with ξ–conformaly flat and ϕ concircularly flat of n dimensional LP sasakian manifold and we proved that ξ–conformaly flatness with special semi-symmetric recurrent metric connection and Riemannian manifold coincide.

Keywords : lorentzian para saskian manifold, semi symmetric recurrent metric connection, ξ–conformaly flat, ϕ concircularly flat.

GJSFR-F Classification : MSC 2000: 53C25

LORENTZIAN PARA SASAKIAN MANIFOLDS ADMITTING SPECIAL SEMI SYMMETRIC RECURRENT METRIC CONNECTION

Strictly as per the compliance and regulations of :

© 2013. Sunil Kumar Srivastava & R. P. Kushwaha. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Global Journal of Science Frontier Research (F) Volume XIII Issue VII Version I Year 2013

Lorentzian Para Sasakian Manifolds Admitting Special Semi Symmetric Recurrent Metric Connection

Sunil Kumar Srivastava [α] & R. P. Kushwaha [σ]

Abstract - Several author as Agashe and Chafle [1], Sengupta, De. U.C, Binh [4] and many other introduced semi symmetric non metric connection in different way. In this paper we have studied LP sasakian manifold with special semi-symmetric recurrent metric connection [2] and discuss it exientance in LP sasakian manifold. In section 3 we establish the relation between the Riemannian connection and special semi-symmetric recurrent metric connection on LP sasakian manifold [4]. The section 4 deals with ξ–conformaly flat and ∅ concircularly flat of n dimensional LP sasakian manifold and we proved that ξ–conformaly flatness with special semi-symmetric recurrent metric connection and Riemannian manifold coincide.

Keywords : lorentzian para saskian manifold, semi symmetric recurrent metric connection, ξ–conformaly flat, ∅ concircularly flat.

I. Introduction

An n-dimensional differentiable manifold M^n is a lorentzian para-sasakian manifold if it admits tensor field of type (1,1), a contravariant vector field ξ, a covariant vector field η and a lorentzian metric g satisfying;

$$\emptyset^2 X = X + \eta(x)\xi \tag{1.1}$$

$$\eta(\xi) = -1 \tag{1.2}$$

$$g(\emptyset X, \emptyset Y) = g(X,Y) + \eta(X)\eta(Y) \tag{1.3}$$

$$g(X,\xi) = \eta(X) \tag{1.4}$$

$$(D_X \emptyset)(Y) = g(X,Y)\xi + \eta(Y)X + 2\eta(X)\eta(Y)\xi \tag{1.5}$$

$$D_X \xi = \emptyset X \tag{1.6}$$

For arbitrary vector field *and* Y, whrere D denotes the operator of covariant differentiation with respect to lorentzian metric g [5].

Author α : Deptt. of Science & Humanities Columbia Institute of Engg. & Technology Raipur India.
Author σ : Deptt. of Mathematics & Statistics D.D.U Gorakhpur University, Gorakhpur India. E-mail : Sunilk537@gmail.com

Ref

1. Agashe, N. S and Chafle, M. R: A semi symmetric non metric connection on a Riema-nnian Manifold. Indian. J. Pure appl. Math. 399-409, 1992.

© 2013 Global Journals Inc. (US)

In a LP sasakian manifold with structure $(\emptyset, \xi, \eta, g)$ the following relation hold.

$$\text{(a)} \quad \emptyset(\xi) = 0 \qquad \text{(b)} \quad \eta(\emptyset X) = 0 \qquad \text{rank}\emptyset = n - 1 \tag{1.7}$$

Let us *put* $F(X,Y) = g(\emptyset X, Y)$ (1.8)

The tensor field F is symmetric (0,2) tensor field ie $F(X,Y) = F(Y,X)$ (1.9)

And
$$(D_X\eta)(Y) = F(X,Y) = g(\emptyset X, Y) \tag{1.10}$$

Also in a LP sasakian manifold the following relation holds;

$$g(R(X,Y)Z,\xi) = \eta(R(X,Y)Z) = g(Y,Z)\eta(X) - g(X,Z)\eta(Y) \tag{1.11}$$

And
$$S(Y,\xi) = (n-1)\eta(X) \tag{1.12}$$

For any vector field X, Y, Z where $R(X,Y)Z$ is the Riemannian curvarture tensor and S is the Ricci tensor.

Let (M^n, g) be an LP sasakian manifold with Levi-Civita connection D. we define a linear connection $\bar{D}$ on M^n by

$$\bar{D}_X Y = D_X Y - \eta(X)Y \tag{1.13}$$

Where η is 1-form associated with vector field ξ on M^n, given by

$$g(X,\xi) = \eta(X) \tag{1.14}$$

Using (1.13) the torsoin tensor $\bar{T}$ of M^n with respect to connection $\bar{D}$ is given by

$$\bar{T}(X,Y) = \bar{D}_X Y - \bar{D}_Y X - [X,Y] = \eta(Y)X - \eta(X)Y \tag{1.15}$$

A linear connection satisfying (1.15) is called semi symmetric connection.
Further from (1.13), we have

$$(\bar{D}_X g)(Y,Z) = 2\eta(X)g(Y,Z) \tag{1.16}$$

A linear connection satisfying (1.16) is called semi symmetric recurrent metric connectionThe word special is used to distinguish it from other connection.

II. Existence of Special Semi-Symmetric Recurrent Metric Connection

Let $\bar{D}$ be a linear connection in M^n, given by

$$\bar{D}_X Y = D_X Y + H(X,Y) \tag{2.1}$$

Where H is a tensor of type (1,2).

Now, we determine the tensor field H such that $\bar{D}$ satifies (1.15) and (1.16).

Notes

© 2013 Global Journals Inc. (US)

Notes

From (2.1), we have

$$\bar{T}(X,Y) = H(X,Y) - H(Y,X) \tag{2.2}$$

Let $G(X,Y,Z) = (\bar{D}_X g)(Y,Z)$ then (2.3)

$$g(H(X,Y),Z) + g(H(X,Z),Y) - -G(X,Y,Z) \tag{2.4}$$

From (2.1),(2.2) and (2.4), we have

$$g(\bar{T}(X,Y),Z) + g(\bar{T}(Z,X),Y) + g(\bar{T}(Z,Y),X)$$

$$= g(H(X,Y),Z) - g(H(Y,X),Z) + g(H(Z,X),Y) - g(H(X,Z),Y) + g(H(Z,Y),X) - g(H(Y,Z),X)$$

$$= 2g(H(X,Y),Z) + 2\eta(X)g(Y,Z) - 2\eta(Z)g(X,Y) + 2\eta(Y)g(X,Z$$

Or

$$H(X,Y) = \frac{1}{2}\{\bar{T}(X,Y) + \bar{T}'(X,Y) + \bar{T}'(Y,X)\} - \eta(X)Y - \eta(Y)X + g(X,Y)\xi \tag{2.5}$$

Where

$$g(\bar{T}'(X,Y),Z) = g(\bar{T}(Z,X),Y) \tag{2.6}$$

Using (1.5), (2.6) we get

$$\bar{T}'(X,Y) = \eta(X)Y - g(X,Y)\xi \tag{2.7}$$

Then in view of (1.15), (2.5) and (2.7),we get

$$H(X,Y) = -\eta(X)Y$$

This implies

$$\bar{D}_X Y = D_X Y - \eta(X)Y$$

Conversely, a connection $\bar{D}$ given by (1.13), satisfies (1.15) and (1.16) show that $\bar{D}$ is a special semi symmetric recurrent metric connection.
So we state the following theorem.

Theorem 2.1: let (M^n, g) be an LP sasakian manifold with lorentzian para contact metric structure $(\emptyset, \xi, \eta, g)$ admits a special semi-symmetric connection which is given by

$$\bar{D}_X Y = D_X Y - \eta(X)Y$$

III. Curvature Tensor of M^n with Respect to Special Semi –Symmetric Recurrent Metric Metric Connection $\bar{D}$

The Curvature tensor of M^n with respect to special semi - symmetric recurrent metric metric connection $\bar{D}$ is given by

$$\bar{R}(X,Y,Z) = \bar{D}_X\bar{D}_Y Z - \bar{D}_X\bar{D}_Y Z - \bar{D}_{[X,Y]}Z$$

Using (1.13) and (1.10) in above we have

$$\bar{R}(X,Y,Z) = R(X,Y,Z) \tag{3.1}$$

Hence we conculude.

© 2013 Global Journals Inc. (US)

Proposition 3.1: The Curvature tensor of M^n with respect to special semi-symmetric recurrent metric metric connection $\bar{D}$ coincide with the curvature tensor of connection D of Riemannian manifold.
Taking the inner product of (3.1) with W, we have

$$\bar{R}(X,Y,Z,W) = R(X,Y,Z,W) \tag{3.2}$$

Where $\bar{R}(X,Y,Z,W) = g(\bar{R}(X,Y,Z),$

From (3.2), we have

$$\bar{R}(X,Y,Z,W) = -R(Y,X,Z,W) \tag{3.3}$$

$$\bar{R}(X,Y,Z,W) = -R(X,Y,W,Z) \tag{3.4}$$

Combining above two relation, we have

$$\bar{R}(X,Y,Z,W) = R(Y,X,W,Z) \tag{3.5}$$

We also have,

$$\bar{R}(X,Y,Z) + \bar{R}(Y,Z,X) + \bar{R}(Z,X,Y) = 0 \tag{3.6}$$

This is the Bianchi first identity for $\bar{D}$.

Hence we conclude that the curvature tensor of M^n with respect to special semi - symmetric recurrent metric metric connection $\bar{D}$ satifies the first Bianchi identy.
Contracting (3.2) over X and W, we obtain

$$\bar{S}(Y,Z) = S(Y,YZ) \tag{3.7}$$

Where $\bar{S}$ and S denote the Ricci tensor of the connection $\bar{D}$ and D respectively.

From (3.7) we obtain a relation between the scalar curvature of M^n with respect to the Riemannian connection and special semi-symmetric recurrent metric connection which is given by

$$\bar{r} = r \tag{3.8}$$

So we have following

Proposition 3.2: Forn dimensional LP sasakian manifold with special semi symmetric recurrent metric connection $\bar{D}$

(1) The curvature tensor $\bar{R}$ is given by (3.1)

(2) Ricci tensor $\bar{S}$ is given by (3.7)

(3) $\bar{r} = r$

Notes

© 2013 Global Journals Inc. (US)

IV. Concircular Curvature Tensor of Lp Sasakian Manifold with Respect to Special Semi-Symmtric Recurrent Metric Connection

Analogous to the definition of concircularcurvarure tensor in a Riemannian manifold we define concircular curvature tensor with respect to the special semi symmetric recurrent metric connection $\bar{D}$ as

$$\bar{C}(X,Y,Z) = \bar{R}(X,Y,Z) - \frac{r}{n(n-1)}\{g(Y,Z)X - g(X,Z)Y\} \tag{4.1}$$

Using (3.1) and (3.7) in (4.1), we have

$$\bar{C}(X,Y,Z) = C(X,Y,Z) \tag{4.2}$$

So we have

Proposition 4.1: $\bar{C}(X,Y,Z) = C(X,Y,Z)$ that is manifold coincide with Riemannian Manifold.

The notion of an ξ-conformaly flat contact manifold was given by Zhen, Cabrezizo and Fermander [3]. In an analogous we define an ξ-conformaly flat n - dimensional LP sasakian manifold.

Defination 5.2: An n-dimensional LP sasakian manifold is called ξ-conformaly flat if the condition $\bar{C}(X,Y)\xi = 0$ holds on M^n.
From (4.2) it is clear that $\bar{C}(X,Y)\xi = C(X,Y)\xi$
So we have the following theorem.

Theorem 4.1: In an n-dimensional LP Sasakian manifold, an ξ–conformaly flatness with respect to special semi-symmetric recurrent metric connection and Riemannian connection coincide.

Defination 4.3: an n dimensional LP Sasakian manifold satisfying the condition

$$\emptyset^2\bar{C}(\emptyset X,\emptyset Y)\emptyset Z = 0 \tag{4.3}$$

Is called $\emptyset -$ concircularly flat.

Letus suppose that M^n be n dimensional $\emptyset -$ concircularly flat LP sasakian manifold with respect to special semi-symmetric recurrent metric connection. It can easily be seen that $\emptyset^2\bar{C}(\emptyset X,\emptyset Y)\emptyset Z = 0$ if and only if

$$g(\bar{C}(\emptyset X,\emptyset Y)\emptyset Z,\emptyset W) = 0 \tag{4.4}$$

For all X,Y, Z, Won T(M).

Using (4.1), $\emptyset -$ concircularly flat means

$$g(\bar{R}(\emptyset X,\emptyset Y)\emptyset Z,\emptyset W) = \frac{r}{n(n-1)}\{g(\emptyset Y,\emptyset Z)g(\emptyset X,\emptyset W) - g(\emptyset X,\emptyset Z)g(\emptyset Y,\emptyset W)\} \tag{4.5}$$

Ref

3. Zhen, G, Cabrezizo J. L, Fernander, L.M and Fernander: on ξ conformally flat contact metric manifold. Indian J. Pure. appl. math 28, 725-734, 1997.

© 2013 Global Journals Inc. (US)

Let $\{e_1, e_2 \ldots., \xi\}$ be a local orthogonal basis of the vector in M^n using the fact that $\emptyset e_1, \emptyset e_2 \ldots. \xi\}$ is also a local orthogonal basis, putting $X = W = e_i$ in (4.5) and summanig with respect to i, we have

$$S(\emptyset Y, \emptyset Z) = \frac{r}{n(n-1)}\{g(\emptyset Y, \emptyset Z)\} \tag{4.6}$$

Putting $Y = \emptyset Y, Z = \emptyset Z$ in (4.6) and using the fact S is symmetric, we have

$$g(\bar{R}(\emptyset X, \emptyset Y)\emptyset Z, \emptyset W) = 0$$

Hence we have

Theorem 4.2: an n-dimensional LP sasakian manifold is $\emptyset -$ concircularly flat with respect to special semi-symmetric recurrent metric connectionand manifold coincide with Riemannian Manifold.

References Références Referencias

1. Agashe, N. S and Chafle, M. R: A semi symmetric non metric connection on a Riema-nnian Manifold. Indian. J. Pure appl. Math. 399-409, 1992.
2. Liang,Y: On semi symmetric recurrent metric connection. Tensor N. S 55, 107-112, 1994.
3. Zhen, G, Cabrezizo J. L, Fernander, L.M and Fernander: on ξ conformally flat contact metric manifold. Indian J. Pure. appl. math 28, 725-734, 1997.
4. Dsa L, De, U.C, Singh, R.N and Pandey M. K: Lorentzian manifold admittin a type of semi symmetric non metric connection. Tensor N. S 70, 70-85, 2008.
5. Mosmuto, M: on Lorentizian para contact manifold. Bull of Yamagata Univ. Nat. Sci1 2 (2), 151-156, 1989.

Notes

© 2013 Global Journals Inc. (US)

Global Journal of Science Frontier Research
Mathematics and Decision Sciences
Volume 13 Issue 7 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Global Attractor of the Nonlinearfour Order Wave Equations

By Ling Zhao
Sichuan Normal University, China

Abstract - Global attractor of the nonlinear four order wave equations is consideredin this paper. Firstly, it is proved that this system possesses an absorbing set in $L^2(\Omega) \times H^2(\Omega)$. Secondly, it is obtained that the nonlinear four order wave equations have a global attractor in $L^2(\Omega) \times H^2(\Omega)$ by using $C-$condition.

Keywords : *global attractor, $C-$condition, four order wave equations.*

GJSFR-F Classification : *MSC 2010: 35B40, 35B41, 35G30, 35L75*

GLOBAL ATTRACTOR OF THE NONLINEARFOUR ORDER WAVE EQUATIONS

Strictly as per the compliance and regulations of :

© 2013. Ling Zhao. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Global Journal of Science Frontier Research (F) Volume XIII Issue VII Version I Year 2013

Global Attractor of the Nonlinearfour Order Wave Equations

Ling Zhao

Abstract - Global attractor of the nonlinear four order wave equations is consideredin this paper. Firstly, it is proved that this system possesses an absorbing set in $L^2(\Omega) \times H^2(\Omega)$. Secondly, it is obtained that the nonlinear four order wave equations have a global attractor in $L^2(\Omega) \times H^2(\Omega)$ by using $C-$ condition.

Keywords : *global attractor, $C-$ condition, four order wave equations.*

I. Introduction

Nonlinear four order wave equations can be applied in mechanics of elastic constructions, and have a long history([1]-[11]). But few of the studies consider attractor of the nonlinear four order wave equations. In this paper we study attractor of the following nonlinear four order wave equations

$$u_{tt} + \triangle^2 u + \alpha u_t = f(u), \quad x \in \Omega, \quad t > 0, \tag{1.1}$$

$$u = \triangle u = 0, \quad x \in \partial\Omega, \quad t > 0, \tag{1.2}$$

$$u(x,0) = \varphi(x), \quad u_t(x,0) = \psi(x), \quad x \in \Omega, \tag{1.3}$$

where $\alpha > 0$, Δ is the Laplacian operator, Ω denotes an open bounded set of $R^n (n = 1, 2, 3)$ with smooth boundary $\partial\Omega$, and $f(s) \in C^1(R)$ satisfies the following conditions

$$F(s) = \int_\Omega f(s)dx \leq c, \tag{1.4}$$

$$sf(s) - F(s) \leq c, \tag{1.5}$$

and there is a $0 \leq r < \infty$, such that

$$\lim_{s\to\infty} \frac{f'(s)}{s^r} = 0. \tag{1.6}$$

Author : *College of Mathematics and Software Science, Sichuan Normal University, Chengdu, Sichuan 610066, China.*
E-mail : zhaoling609@163.com

Ref

[1] Eisley JG: Nonlinear vibration of beams and rectangular plates. Z.Angew. Math. Phys. 15, 167-175 (1964)

© 2013 Global Journals Inc. (US)

As far as the theory of infinite-dimensional dynamical system is concerned, we refer to [12]-[22]. In the study of infinite dimensional dynamical system, the long-time behavior of the solution to equations is an important issue. The long-time behavior of the solution to equations can be shown by the global attractor with the finite-dimensional characteristics. Some authors have already studied the existence of the global attractor for some evolution equations. The global attractor strictly defined as ω-limit set of ball, which under additional assumptions is nonempty, compact, and invariant ([14],[16]). We investigate global attractor of the equations (1.1)-(1.3) in this article.

The paper is organized as follows. In Section 2 we recall preliminary results. In Section 3, we obtain global attractor of the equations.

II. Preliminaries

Let X and X_1 be two Banach spaces, $X_1 \subset X$ a compact and dense inclusion. Consider the abstract nonlinear evolution equation defined on X, given by

$$\begin{cases} \frac{du}{dt} = Lu + G(u), \\ \\ u(x,0) = u_0. \end{cases} \tag{2.1}$$

where $u(t)$ is an unknown function, $L : X_1 \to X$ a linear operator, and $G : X_1 \to X$ a nonlinear operator.

A family of operators $S(t) : X \to X (t \geq 0)$ is called a semigroup generated by (2.1) if it satisfies the following properties:

(1) $S(t) : X \to X$ is a continuous map for any $t \geq 0$,

(2) $S(0) = id : X \to X$ is the identity,

(3) $S(t+s) = S(t) \cdot S(s), \forall t, s \geq 0$. Then, the solution of (2.1) can be expressed as

$$u(t, u_0) = S(t)u_0.$$

Next, we introduce the concepts and definitions of invariant sets, global attractors, and ω-limit sets for the semigroup $S(t)$.

Definition 2.1 Let $S(t)$ be a semigroup defined on X. A set $\Sigma \subset X$ is called an invariant set of $S(t)$ if $S(t)\Sigma = \Sigma, \forall t \geq 0$. An invariant set Σ is an attractor of $S(t)$ if Σ is compact, and there exists a neighborhood $U \subset X$ of Σ such that for any $u_0 \in U$,

Ref
[12] Babin AV, Vishik MI: Attractors of Evolution Equations, North-Holland, Amsterdam, 1992.

© 2013 Global Journals Inc. (US)

$$\inf_{v\in\Sigma}\|S(t)u_0 - v\|_X \to 0, \quad \text{as} \quad t\to\infty.$$

In this case, we say that Σ attracts U. Especially, if Σ attracts any bounded set of X, Σ is called a global attractor of $S(t)$ in X.

For a set $D\subset X$, we define the ω-limit set of D as follows

$$\omega(D) = \bigcap_{s\geq 0}\overline{\bigcup_{t\geq s} S(t)D},$$

where the closure is taken in the X-norm. Lemma 2.2 is the classical existence theorem of global attractor by Temam [16].

Lemma 2.2 Let $S(t) : X \to X$ be the semigroup generated by (2.1). Assume the following conditions hold

(1) $S(t)$ has a bounded absorbing set $B\subset X$, i.e., for any bounded set $A\subset X$ there exists a time $t_A\geq 0$ such that $S(t)u_0\in B, \forall u_0\in A$ and $t>t_A$;

(2) $S(t)$ is uniformly compact, i.e., for any bounded set $U\subset X$ and some $T>0$ sufficiently large, the set $\overline{\bigcup_{t\geq T} S(t)U}$ is compact in X.

Then the ω-limit set $\mathcal{A}=\omega(B)$ of B is a global attractor of (2.1), and $\mathcal{A}$ is connected providing B is connected.

Definition 2.3[15] We say that $S(t) : X\to X$ satisfies $C-$condition, if for any bounded set $B\subset X$ and $\varepsilon>0$, there exist $t_B>0$ and a finite dimensional subspace $X_1\subset X$, such that $\{PS(t)B\}$ is bounded, and

$$\|(I-P)S(t)u\|_X < \varepsilon, \quad \forall t\geq t_B \ \text{ and } \ u\in B,$$

where $P : X\to X_1$ is a projection.

Lemma 2.4[15] Let $S(t) : X\to X(t\geq 0)$ be a dynamical systems. If the following condition are satisfied

(1) there exists a bounded absorbing set $B\subset X$,

(2) S(t) satisfies $C-$condition,

then $S(t)$ has a global attractor in X.

We introduce the spaces

$$H = L^2(\Omega), \quad V = \{u\in H^2 | u|_{\partial\Omega} = \triangle u|_{\partial\Omega} = 0\}, \quad E = H\times V.$$

Ref

[16] Temam R: Infinite-Dimensional Dynamical Systems in Mechanics and physics. Applied Mathematical Science, Vol. 68, Springer-Verlag, New York, 2nd., 1997

© 2013 Global Journals Inc. (US)

The denotations $(\cdot,\cdot), \|\cdot\|$ are the inner product and norm in H respectively, i.e., for any $u, v \in H$, $(u,v) = \int_\Omega uvdx$, $\|u\| = (\int_\Omega |u|^2 dx)^{\frac{1}{2}}$. The norm in V is $\|u\|_V = \|\triangle u\|$ and we have $\|\triangle u\|^2 \geq \beta_1 \|u\|^2$, where β_1 is the first eigenvalue of Δu with boundary condition (1.2).

Lemma 2.5 Assume the nonlinear term $f : R \to R$ satisfies (1.6). Then the nonlinear operator $f : V \to H$ is compact.

Proof. Let $\{u_n\}$ be a bounded sequence in V. By the Soblev embedding theorem, $\{u_n\}$ is bounded in $L^k(\Omega)$ for any $1 \leq k < \infty$, and has a convergent subsequence in $L^2(\Omega)$. Without loss of generality, we assume $\{u_n\}$ converges to u_0 in H. It is sufficient to prove that $\{f(u_n)\}$ converges to $f(u_0)$ in H.

From (1.6), for any $\eta > 0$, we have

$$|f'(s)|^2 \leq \eta |s|^{2r} + c_\eta,$$

where $c_\eta \to 0$ if $\eta \to 0$.

Then, for some $0 \leq \varepsilon \leq 1$, we obtain

$$\begin{aligned}
\int_\Omega |f(u_n) - f(u_0)|^2 dx &= \int_\Omega |f'(u_0 + (1-\varepsilon)(u_n - u_0))|^2 |u_n - u_0|^2 dx \\
&\leq \eta \int_\Omega |u_0 + (1-\varepsilon)(u_n - u_0)|^{2r} |u_n - u_0|^2 dx + c_\eta \int_\Omega |u_n - u_0|^2 dx \\
&= \eta \int_\Omega |\varepsilon u_0 + (1-\varepsilon) u_n)|^{2r} |u_n - u_0|^2 dx + c_\eta \int_\Omega |u_n - u_0|^2 dx \\
&\leq 4^r \eta \int_\Omega (|u_0|^{2r} + |u_n|^{2r}) |u_n - u_0|^2 dx + c_\eta \int_\Omega |u_n - u_0|^2 dx \\
&\leq 4^r \eta [(\int_\Omega |u_0|^{2rp} dx)^{\frac{1}{p}} + (\int_\Omega |u_n|^{2rp} dx)^{\frac{1}{p}}](\int_\Omega |u_n - u_0|^{2q} dx)^{\frac{1}{q}} + c_\eta \int_\Omega |u_n - u_0|^2 dx
\end{aligned}$$

where $p > 0, q > 0$, and $\frac{1}{p} + \frac{1}{q} = 1$.

Let $n \to \infty$ and $\eta \to 0$. We have

$$\int_\Omega |f(u_n) - f(u_0)|^2 dx = 0.$$

Then the nonlinear operator $f : V \to H$ is compact.

III. Existence of Global Attractor

Theorem 3.1 Let $\varphi \in H$, $\psi \in V$, and f satisfy the conditions (1.4) and (1.5). Suppose that the problem (1.1)-(1.3) has a unique weak solution and $S(t), t > 0$, defined by $S(t)(\varphi, \psi) =$

Notes

© 2013 Global Journals Inc. (US)

$(u(t), u_t(t))$, is the semigroup generated by the problem (1.1)-(1.3). Then $S(t)$ has a bounded absorbing ball.

Proof. Take the inner product of (1.1) in H with $v = u_t + \theta u$, $0 < \theta \leq \theta_0$, and θ_0 will be chosen later. we obtain

Notes

$$(u_{tt}, v) + (\triangle^2 u, v) + \alpha(u_t, v) = (f, u).$$

Using the condition (1.4), we have

$$\frac{1}{2}\frac{d}{dt}(\|v\|^2 + \|\triangle u\|^2 - 2\int_\Omega F(u)dx) + (\alpha - \theta)\|v\|^2 + \theta\|\triangle u\|^2$$

$$+(\theta^2 - \alpha\theta)(u, v) - \theta\int_\Omega f(u)udx = 0. \tag{3.1}$$

Using the Hölder inequality, Young inequality and the condition (1.5), we get

$$(\alpha - \theta)\|v\|^2 + \theta\|\triangle u\|^2 + (\theta^2 - \alpha\theta)(u, v) - \theta\int_\Omega f(u)udx$$

$$\geq (\alpha - \theta)\|v\|^2 + \theta\|\triangle u\|^2 + (\theta^2 - \alpha\theta)\|u\|\|v\| - \theta\int_\Omega (F(u) + c)dx$$

$$\geq (\alpha - \theta)\|v\|^2 + \theta\|\triangle u\|^2 - \frac{\alpha\theta}{\beta_1^{\frac{1}{2}}}\|\triangle u\|\|v\| - \theta\int_\Omega (F(u) + c)dx$$

$$\geq (\alpha - \theta)\|v\|^2 + \theta\|\triangle u\|^2 - \frac{\theta}{2}\|\triangle u\|^2 - \theta\frac{\alpha^2}{2\beta_1}\|v\|^2 - \theta\int_\Omega (F(u) + c)dx$$

$$\geq [\alpha - \theta(1 + \frac{\alpha^2}{2\beta_1})]\|v\|^2 + \frac{\theta}{2}\|\triangle u\|^2 - \theta\int_\Omega (F(u) + c)dx.$$

Choose θ_0, such that $\theta_0(1 + \frac{\alpha^2}{2\beta_1}) = \frac{\alpha}{2}$. Hence, $\alpha - \theta(1 + \frac{\alpha^2}{2\beta_1}) \geq \frac{\alpha}{2} \geq \theta$.

From (3.1), we have

$$\frac{1}{2}\frac{d}{dt}(\|v\|^2 + \|\triangle u\|^2 - 2\int_\Omega F(u)dx) + \frac{\alpha}{2}\|v\|^2 + \frac{\theta}{2}\|\triangle u\|^2 - \theta\int_\Omega (F(u) + c)dx \leq 0.$$

Then, it follows that

$$\frac{d}{dt}[\|v\|^2 + \|\triangle u\|^2 + 2\int_\Omega (c - F(u))dx] + \theta[\|v\|^2 + \|\triangle u\|^2 + 2\int_\Omega (c - F(u))dx]$$

$$\leq 4c\theta|\Omega|, \tag{3.2}$$

where $|\Omega|$ is measure of the Ω.

Let $y(t) = \|v\|^2 + \|\triangle u\|^2 + 2\int_\Omega (c - F(u))dx$. Then $y(t) \geq 0$ and (3.2) can be read as

$$\frac{dy}{dt} + \theta y \leq 4c\theta|\Omega|.$$

© 2013 Global Journals Inc. (US)

Using the Gronwall inequality, we have

$$y(t) \leq y(0)e^{-\theta t} + 4c|\Omega|(1 - e^{-\theta t}), \quad t \geq 0.$$

For any bounded set B of E, $(\varphi, \psi) \in B$ and $\int_\Omega [c - F(\varphi)]dx$ is bounded. Then

$$R(B) = \sup_{(\varphi,\psi)\in B} y(0) = \sup_{(\varphi,\psi)\in B} \{\|\varphi\|_V^2 + \|\psi + \theta\varphi\| + 2\int_\Omega (c - F(\varphi))dx\} < \infty,$$

and

$$\lim_{t\to\infty} \sup_{(\varphi,\psi)\in B} y(t) \leq 4c|\Omega| = \mu_0^2.$$

Let $\mu_1 > \mu_0$ be fixed, and there is a $t_0 = t_0(R(B), \mu_1) = \frac{1}{\theta} \ln \frac{R(B)}{\mu_1^2 - \mu_0^2}$ such that for any $t \geq t_0$, we have $y(t) \leq \mu_1^2$. Then

$$\begin{aligned}
\|u(t)\|_V^2 + \|u_t(t)\|^2 &\leq \|u(t)\|_V^2 + 2\|u_t(t) + \theta u(t)\|^2 + 2\theta^2\|u(t)\|^2 \\
&\leq \|u(t)\|_V^2 + 2\|u_t(t) + \theta u(t)\|^2 + \frac{2\theta^2}{\beta_1}\|u(t)\|_V^2 \\
&\leq 2(1 + \frac{\theta^2}{\beta_1})(\|u(t)\|_V^2 + \|u_t(t) + \theta u(t)\|^2) \\
&\leq 2(1 + \frac{\theta^2}{\beta_1})y(t) \leq 2(1 + \frac{\theta^2}{\beta_1})\mu_1^2.
\end{aligned}$$

Let $\rho_0^2 = 2(1 + \frac{\theta^2}{\beta_1})\mu_1^2$. Then For all $t \geq t_0$, we have

$$\|u\|_V^2 + \|u_t\|^2 \leq \rho_0^2, \tag{3.3}$$

which implies that the ball of E, $B_0 = B_E(0, \rho_0)$, centered at 0 of radius ρ_0, is an absorbing set.

Theorem 3.2 Assume $f : R \to R$ satisfies (1.4)-(1.6). Then the semigroup $S(t), t \geq 0$ associated with problem (1.1)-(1.3) possesses a global attractor.

Proof. The eigenvalue equation

$$\begin{cases} \Delta^2 u = \beta u, \\ u|_{\partial\Omega} = \triangle u|_{\partial\Omega} = 0, \end{cases} \tag{3.4}$$

has eigenvalues $\beta_1, \beta_2, \cdots, \beta_k, \cdots$and eigenvector $\{e_k | k = 1, 2, 3, \cdots\}$, and $0 < \beta_1 \leq \beta_2 \leq \cdots \leq \beta_k \geq \cdots$. If $k \to \infty$, $\beta_k \to \infty$. $\{e_k | k = 1, 2, 3, \cdots\}$ constitutes an orthogonal base of H.

For $\forall u \in H$, we have

$$u = \sum_{k=1}^{\infty} u_k e_k, \quad \|u\|_{L^2}^2 = \sum_{k=1}^{\infty} u_k^2,$$

Notes

© 2013 Global Journals Inc. (US)

Notes

Introduce subspace $E_1 = span\{e_1, e_2, \cdots, e_k\} \subset H$. Let E_2 be an orthogonal subspace of $E_1 \subset H$.

For $\forall (u_t, u) \in E$, we find

$$u_t = u_{1t} + u_{2t}, \quad u_{1t} = P_m u_t, \quad u_{2t} = (I - P_m)u_t,$$

and

$$u = u_1 + u_2, \quad u_1 = P_m u \quad u_2 = (I - P_m)u,$$

where $P_m : H \to E_1$ be the orthogonal projection.

Let $P_i : E \to E_i \times E_i$ be the orthogonal projection. Thanks to Definition 2.3, we will prove that for any bounded set $B \subset E$ and $\varepsilon > 0$, there exists $t_* > 0$ such that

$$\|P_1 S(t)B\|_H \leq M, \quad \forall t > t_*, \quad M \quad \text{is a constant}, \tag{3.5}$$

$$\|P_2 S(t)B\|_H \leq \varepsilon, \quad \forall t > t_*, \quad \varphi \in B. \tag{3.6}$$

From Theorem 3.1, $S(t)$ has an absorbing set B_M. Then for any bounded set $B \subset E$, there exists $t_0 > 0$ such that $S(t)B \subset B_M$, $\forall t > t_0$, which imply (3.5).

From Lemma 2.5, $f : V \to H$ is compact, then for any $\varepsilon > 0$, there exists some $m \in N$ such that

$$\|(I - P_m)f\| \leq \varepsilon.$$

Multiply (1.1) by $v_2 = u_{2t} + \theta u_2$ $(0 < \theta \leq \theta_1)$, and integrate over Ω. We obtain

$$\int_\Omega (u_{tt} + \triangle^2 u + \alpha u_t) v_2 dx = \int_\Omega f v_2 dx.$$

Then, we have

$$\frac{1}{2}\frac{d}{dt}(\|v_2\|^2 + \|\triangle u_2\|^2) + (\alpha - \theta)\|v_2\|^2 + \theta\|\triangle u_2\|^2$$

$$+(\theta^2 - \alpha\theta)(u_2, v_2) \leq (f, v_2) \leq \|(I - P_m)f\|\|v_2\| \leq \varepsilon\|v_2\| \leq \frac{\alpha}{4}\|v_2\|^2 + \frac{\varepsilon^2}{\alpha}. \tag{3.7}$$

Using the Hölder inequality and Young inequality, we get

$$(\alpha - \theta)\|v_2\|^2 + \theta\|\triangle u_2\|^2 + (\theta^2 - \alpha\theta)(u_2, v_2)$$

$$\geq (\alpha - \theta)\|v_2\|^2 + \theta\|\triangle u_2\|^2 + (\theta^2 - \alpha\theta)\|u_2\|\|v_2\|$$

$$\geq (\alpha - \theta)\|v_2\|^2 + \theta\|\triangle u_2\|^2 - \frac{\alpha\theta}{\beta_{m+1}^{\frac{1}{2}}}\|\triangle u_2\|\|v\|$$

© 2013 Global Journals Inc. (US)

$$\geq (\alpha-\theta)\|v_2\|^2+\theta\|\triangle u_2\|^2-\frac{\theta}{2}\|\triangle u_2\|^2-\theta\frac{\alpha^2}{2\beta_{m+1}}\|v_2\|^2$$

$$\geq [\alpha-\theta(1+\frac{\alpha^2}{2\beta_{m+1}})]\|v_2\|^2+\frac{\theta}{2}\|\triangle u_2\|^2.$$

Choose θ_1, such that $\theta_1(1+\frac{\alpha^2}{2\beta_1})=\frac{\alpha}{2}$. Hence, $\alpha-\theta(1+\frac{\alpha^2}{2\beta_1})\geq\frac{\alpha}{2}\geq\theta$.

From (3.7), we have

$$\frac{1}{2}\frac{d}{dt}(\|v_2\|^2+\|\triangle u_2\|^2)+\frac{\alpha}{4}\|v_2\|^2+\frac{\theta}{2}\|\triangle u_2\|^2\leq\frac{\varepsilon^2}{\alpha}.$$

Then, it follows that

$$\frac{d}{dt}(\|v_2\|^2+\|\triangle u_2\|^2)+\theta(\|v\|^2+\|\triangle u\|^2)\leq\frac{2\varepsilon^2}{\alpha}. \tag{3.8}$$

Using the Gronwall inequality, we have

$$\|v_2(t)\|^2+\|\triangle u_2(t)\|^2\leq(\|v_2(0)\|^2+\|\triangle u_2(0)\|^2)e^{-\theta t}+\frac{2\varepsilon^2}{\alpha}(1-e^{-\theta(t-t_0)}),\quad t\geq t_0.$$

where t_0 is given in the proof of Theorem 3.1. Then

$$\|v_2(0)\|^2+\|\triangle u_2(0)\|^2\leq\rho_0^2.$$

Let $t_1-t_0=\frac{1}{\theta}\ln\frac{\rho_0^2\alpha}{\varepsilon^2}$. Hence

$$\|v_2(t)\|^2+\|\triangle u_2(t)\|^2\leq\frac{3\varepsilon^2}{\alpha},\quad t\geq t_1.$$

Then

$$\|u_2(t)\|_V^2+\|u_{2t}(t)\|^2\leq\|u_2(t)\|_V^2+2\|u_{2t}(t)+\theta u_2(t)\|^2+2\theta^2\|u_2(t)\|^2$$

$$\leq\|u_2(t)\|_V^2+2\|u_{2t}(t)+\theta u_2(t)\|^2+\frac{2\theta^2}{\beta_{m+1}}\|u_2(t)\|_V^2$$

$$\leq 2(1+\frac{\theta^2}{\beta_{m+1}})(\|u_2(t)\|_V^2+\|u_{2t}(t)+\theta u_2(t)\|^2)$$

$$\leq 2(1+\frac{\theta^2}{\beta_{m+1}})(\|v_2(t)\|^2+\|\triangle u_2(t)\|^2)\leq(1+\frac{\theta^2}{\beta_{m+1}})\frac{6\varepsilon^2}{\alpha}.$$

Let $C=\frac{6}{\alpha}(1+\frac{\theta^2}{\beta_{m+1}})$. Then for all $t\geq t_1$, we have

$$\|u\|_V^2+\|u_t\|^2\leq C\varepsilon^2, \tag{3.9}$$

which implies (3.6).

From Lemma 2.4, the equations(1.1)-(1.3) possess a global attractor.

Notes

© 2013 Global Journals Inc. (US)

References Références Referencias

Notes

[1] Eisley JG: Nonlinear vibration of beams and rectangular plates. Z.Angew. Math. Phys. 15, 167-175 (1964)

[2] Medeiros LA: On a new class of nonlinear wave equations. J. Math. Anal. Appl. 69, 252-262 (1979)

[3] Fitzgibbon WE: Global existence and boundedness of solutions to the extensible beam equation. SIAM J. Math. Anal. 13, 739-745 (1982)

[4] Kouemou Patcheu S: On a global solution and asymptotic behaviour for the generalized damped extensible beam equation. J. Diff. Equa. 135, 299-314 (1997)

[5] Weinstock R: Calculus of variations with applications to physics and engineering. McGraw-Hill Book Company Inc., New York-Toronto-London (1952)

[6] Agliardi R, Cicognani M: Operators of p-evolution with nonregular coefficients in the time variable. J. Diff. Equa. 202, 143-157 (2004)

[7] Agliardi R, Zanghirati L: Cauchy problem for nonlinear p-evolution equations. Bull. Sci. Math., 133(4), 406-418 (2009)

[8] Pausader B: Scattering and the Levandosky-Strauss conjecture for fourth-order nonlinear wave equations. J. Diff. Equa., 241: 237-278 (2007)

[9] Tataru D, Tucsnak M: On the Cauchy problem for the full von Karman system. NoDEA Nonlinear Differential Equations Appl. 4(3): 325-340 (1997)

[10] Koch H, Stahel A: Global existence of classical solutions to the dynamics of von Karman equations. Math. Methods Appl. Sci. 16: 581-586 (1993)

[11] Favini A, Horn M, Lasiecka I, Tataru, D: Global existence, uniqueness and regularity of solutions to a von Karman equation with nonlinear boundary dissipation. Diff. Inte. Equa. 9: 267-294 (1996)

© 2013 Global Journals Inc. (US)

[12] Babin AV, Vishik MI: Attractors of Evolution Equations, North-Holland, Amsterdam, 1992.

[13] Ghidaglia JM: A note on the strong convergence towards attractors for damped forced Kdv equations. J. Diff. Eqn., 110: 356-359(1994)

[14] Hale JK Asymptotic Behavior of Dissipative Systems, in: Math. Surveys Monographs, Vol. 25, American Mathematical Society, Providence, RI, 1987.

[15] Ma QY, Wang SH, Zhong CK: Necessary and sufficient conditions for the existence of global attractor for semigroups and applications. Indiana Univ. Math. J., 51(6): 1541-1570(2002)

[16] Temam R: Infinite-Dimensional Dynamical Systems in Mechanics and physics. Applied Mathematical Science, Vol. 68, Springer-Verlag, New York, 2nd., 1997

[17] Carvalho AN, Cholewa JW: Attractors for strongly damped wave equations with critical nonlinearities. Pac. J. Math., 207: 287-310(2002)

[18] Massatt P: Longtime behaviour of strongly damped nonlinear wave equations. J. Diff. Eqn., 48: 334-349 (1983)

[19] Fan X, Zhou S: The Inflated Attractors of Non-autonomous Strongly Damped Wave Equations. Acta Math. Appl. Sinica, English Series. 20(4): 547-556(2004)

[20] Belleri V, Pata, V: Attractors for semilinear strongly damped wave equation on R^3. Discrete Contin. Dynam. Systems, 7: 719-735(2001)

[21] Zhang YH, Zhong CK: Existence of global attractors for a nonlinear wave equation. Appl. Math. Lett, 18:77-84(2005)

[22] Carvalho AN, Cholewa JW: Attractors for strongly damped wave equations with critical nonlinearities. Pacific J. Math., 207: 287-310(2002)

Notes

© 2013 Global Journals Inc. (US)

Global Journal of Science Frontier Research
Mathematics and Decision Sciences
Volume 13 Issue 7 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

On Semi-Open Sets and Semi-Separability

By V. Srinivasa kumar
JNTU University, India

Abstract - In this paper, we introduce the concepts of semi-limit and semi-separability. We prove that separability and semi-separability are equivalent and also prove a few interesting results in this connection.

Keywords : semi-open, semi-closed, semi-closure, semi-neighborhood, semi-limit, semise-parability, sepa-rability, semi-topology.

GJSFR-F Classification : AMS Code: 54A10, 54A20, 54C08

ON SEMI-OPEN SETS AND SEMI-SEPARABILITY

Strictly as per the compliance and regulations of :

© 2013. V. Srinivasa kumar. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Global Journal of Science Frontier Research (F) Volume XIII Issue VII Version I Year 2013

On Semi-Open Sets and Semi-Separability

V. Srinivasa kumar

Abstract - In this paper, we introduce the concepts of semi-limit and semi-separability. We prove that separability and semi-separability are equivalent and also prove a few interesting results in this connection.

Keywords : semi-open, semi-closed, semi-closure, semi-neighborhood, semi-limit, semiseparability, separability, semi-topology.

I. Introduction

In 1963, Norman Levin introduced the concept of semi-open sets in his paper [2]. It has drawn the attention of various authors including Crossley, Hildebrand and Dorsett and they have probed deeply into this area and developed many interesting concepts like semi-closed sets, semi-compactness etc. In this present paper, we introduce the concepts of semi-limit and semi-separability and prove that semi-separability is equivalent to separability. Also we construct a topology using semi-open sets and we call this topology a semi-topology.

In what follows (X,T) stands for a topological space. The symbols $cl(\ )$ and $Int(\ )$ denote the closure and interior in a topological space respectively.

a) *Preliminaries*

1.1 Definition: Let A be a subset of X. A is said to be

(i) semi-open in (X,T) if $A \subseteq cl(Int(A))$.

(ii) semi-closed if $X - A$ is semi-open in (X,T).

(iii) semi-neighborhood of a point $x \in X$ if $x \in A$ and A is semi-open in (X,T).

1.2 Definition: The semi-closure of a set A in (X,T) denoted by $scl(A)$, is the intersection of all semi-closed supersets of A.

1.3 Definition: A point $x \in X$ is said to be a semi-limit point of a set A in (X,T), if every semi-neighborhood of x contains a point of A different from x in X.

Author : Assistant Professor, Department of Mathematics, JNTUH College of Engineering, JNTU, Hyderabad-500085, A.P., India.
E-mail : srinu_vajha@yahoo.co.in

Ref

2. Levine, N. *Semi-open sets and semi-continuity in topological space*, Amer. Math Monthly, 70, pp. 36-41, 1963.

© 2013 Global Journals Inc. (US)

b) Semi-Limit and Semi-Topology

2.1 Definition: Let $x \in X$ and let $\{ x_\lambda / \lambda \in \Delta \}$ be a net in (X,T). We say that x is a semi-limit of $\{x_\lambda / \lambda \in \Delta\}$ and we write $x = s.\lim_{\lambda \in \Delta} x_\lambda$ if for every semi-neighborhood A of x in X there exists a $\lambda_A \in \Delta$ such that $x_\lambda \in A \quad \forall \lambda \geq \lambda_A$.

2.2 Proposition: For $A \subset X$ and $x \in X$, the following are equivalent.

(i) x is a semi-limit point of A

(ii) there exists a net $\{x_\lambda / \lambda \in \Delta\}$ in A such that $x = s.\lim_{\lambda \in \Delta} x_\lambda$

(iii) $x \in scl(A)$

2.3 Remark: Let $S(T)$ be the collection of all semi-open sets in (X,T). The set $S(T)$ clearly contains T and is closed under arbitrary unions. However, being not closed under finite intersections, $S(T)$ is not a topology on X. However, if $A \in T$ and $B \in S(T)$ then $A \bigcap B \in S(T)$.

2.4 Definition: We define $S_0(T) = \{A \in S(T) / A \bigcap B \in S(T) \ \ \forall B \in S(T)\}$ and $S_{00}(T) = \{A \in S(S_0) / A \bigcap B \in S(S_0) \ \ \forall B \in S(S_0)\}$ where $S(S_0)$ is the collection of all semi-open sets in the topological space $(X, S_0(T))$.

2.5 Proposition:

(a) $S_0(T)$ and $S_{00}(T)$ are topologies on X.

(b) $T \subseteq S_0(T) \subseteq S(T)$.

(c) $S(S_0) \subseteq S(T)$.

(d) $S_0(T) = S_{00}(T)$.

2.6 Remark: We call the topology $S_0(T)$, a semi-topology on X.

2.7 Notation: We denote the closure of a subset A of X in the topological space $(X, S_o(T))$ by the symbol $cl_0(A)$ and interior of A by $Int_o(A)$.

2.8 Proposition: For $A \subseteq X$, $scl(A) \subseteq cl_0(A) \subseteq cl(A)$.

c) Semi-Separability

3.1 Definition: (X,T) is said to be separable if there exists a countable subset A of X such that $cl(A) = X$.

Notes

© 2013 Global Journals Inc. (US)

Notes

3.2 Definition: (X,T) is said to be semi-separable if there exists a countable subset A of X such that $scl(A)=X$.

3.3 Proposition: (X,T) is separable if and only if it is semi-separable.

Proof: Suppose that (X,T) is separable.

$\Rightarrow$ there exists a countable subset A of X such that $cl\,(A)=X$.

Let $x\in X$ and G be a semi-neighborhood of x in (X,T)

$\Rightarrow$ $x\in G$ and G is semi-open in (X,T)

$\Rightarrow$ there exists $O\in T$ such that $O\subseteq G\subseteq cl\,(O)$.

Assume that $G\bigcap A=\phi$ $\Rightarrow A\subseteq X-G\subseteq X-O$

$\Rightarrow cl\,(A)\subseteq X-O$ $\Rightarrow X=X-O$

$\Rightarrow O=\phi$ $\Rightarrow G=\phi$ which is a contradiction.

Hence $G\bigcap A\neq\phi$.

Thus each semi-neighborhood of x in (X,T) intersects A

$\Rightarrow$ $x\in s\,cl(A)$. Hence $s\,cl(A)=X$ $\Rightarrow$ X is semi-separable.

The converse follows from the definitions 3.1, 3.2 and the proposition 2.8.

3.4 Proposition: $(X,S_0(T))$ is semi-separable $\Leftrightarrow$ (X,T) is semi-separable.

Proof: Suppose that $(X,S_0(T))$ is semi-separable

$\Rightarrow$ there exists a countable subset A of X such that $cl_0(A)=X$

$\Rightarrow cl^{(}A)=X$ $\Rightarrow (X,T)$ is separable and hence it is semi-separable.

Conversely suppose that (X,T) is separable $\Rightarrow (X,T)$ is semi-separable

$\Rightarrow$ there exists a countable subset A of X such that $scl(A)=X$

$\Rightarrow$ $cl_0(A)=X$ $\Rightarrow$ $(X,S_0(T))$ is separable and hence it is semi-separable.

II. Acknowledgements

I sincerely thank my Professor, Dr. ***I. Ramabhadrasarma*** for his guidance and encouragement in making this paper.

© 2013 Global Journals Inc. (US)

References Références Referencias

1. **Crossley, S., and Hildebrand, S.** *Semi-Closure,* Texas J.Sci., 22, pp. 99-112, 1970.
2. **Levine, N.** *Semi-open sets and semi-continuity in topological space,* Amer. Math Monthly, 70, pp. 36-41, 1963.
3. **Ramabhadra Sarma, I., and V. Srinivasa kumar** *On Semi-Open Sets and Semi-Continuity*, Journal of Advanced Studies in Topology, Vol. 3, No. 3, pp. 06-102, 2012.

Notes

© 2013 Global Journals Inc. (US)

Global Journal of Science Frontier Research
Mathematics and Decision Sciences
Volume 13 Issue 7 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Lie Algebraic Approach and Complex Invariant Coupled Oscillator Systems

By Jasvinderpal Singh Virdi
Panjab University, India

Abstract - In classical mechanics, the system of coupled harmonic oscillators is shown to possess the symmetry applicable toa six-dimensional space in complex coordinates, two-dimensional phase space consisting of two position and twomomentum variables. In search into the features of a dynamical system, with the possibility of its complex invariant,we explore this dynamical systems. Dynamical algebraic approach is used to study two-dimensional complex systems(coupled oscillator system) on the extended complex phase plane (ECPS). Scope and importance of invariants in theanalysis of complex trajectories for dynamical systems is discussed.

Keywords : *complex hamiltonian, complex invariant.*

GJSFR-F Classification : *AMS Classification: 37K10, 03D15, 34C45*

LIE ALGEBRAIC APPROACH AND COMPLEX INVARIANT COUPLED OSCILLATOR SYSTEMS

Strictly as per the compliance and regulations of :

© 2013. Jasvinderpal Singh Virdi. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Global Journal of Science Frontier Research (F) Volume XIII Issue VII Version I Year 2013

Lie Algebraic Approach and Complex Invariant Coupled Oscillator Systems

Jasvinderpal Singh Virdi

Abstract - In classical mechanics, the system of coupled harmonic oscillators is shown to possess the symmetry applicable toa six-dimensional space in complex coordinates, two-dimensional phase space consisting of two position and twomomentum variables. In search into the features of a dynamical system, with the possibility of its complex invariant,we explore this dynamical systems. Dynamical algebraic approach is used to study two-dimensional complex systems(coupled oscillator system) on the extended complex phase plane (ECPS). Scope and importance of invariants in theanalysis of complex trajectories for dynamical systems is discussed.

Keywords : *complex hamiltonian, complex invariant.*

I. Introduction

The coupled oscillator provides many soluble models in different branches of physics because of its mathematical simplicity. It stays with us in many different forms because it provides the mathematical basis for many soluble models in physics, including the Lee model in quantum field theory [1], the Bogoliubov transformation in superconductivity, relativistic models of elementary particles [2]. In physics coupled harmonic oscillator system in two-dimensions [3], in the Bells inequality experiments employing coupled harmonic oscillators [4]. This also has been used for description of motion of a charged particle in a magnetic field [5, 6] or . They are also studied in context of electrical circuits with time-varying capacitors and inductors, particularly with reference to their memory property, has become of considerable interest in recent years [7]. Blasone *et al* [8] studied momentum-dependent terms in the Hamiltonian structure in the context of the so-called holographic principle and in the treatment of quantum gravity as a dissipative and deterministic system. Hamiltonian for such system is given by

$$H = \frac{1}{2}[\alpha_1 p_x^2 + \alpha_2 p_y^2 + \beta_1 x^2 + \beta_2 y^2] + \alpha_3(p_x y + x p_y). \tag{1}$$

Invariants for above Hamiltonian provided they exist and can be computed, and even the complex invariants [9] exist, are a very useful tool to understand the theoretical structure of this dynamical systems. Since invariants of real Hamiltonian systems have been played a vital role in understanding the underlying dynamics of the systems and so we expect that the complex invariants can also be helpful in exploring some deep insights into features of complex dynamical systems. In the past, complex invariants have been discussed in context of understanding fermion masses and quark mixing, and CP-conserving two-Higgs-doublet model scalar potentials in the Particle physics [10, 11].
In this paper, we construct a complex invariants corresponding coupled oscillators based on the ECPS approach in complex domain [12]. Recently, with a view to explore some role of invariants for complex systems, Kaushal et al. [13] found invariants for some one dimensional systems within the framework of an ECPS. Some quantum mechanical studies within the ECPS are also reported [14]. But most of such studies are restricted in one dimension only. Such studies in higher dimensions is desirable from the intrinsic mathematical interest, to check the validity of various methods/theories and to find solutions of some realistic physical problems. With this motivation, recently we generalized the ECPS in two dimensions and studied the coupled oscillator.

Author : *Department of Physics, Panjab University, Chandigarh-160014. E-mail : jpsvirdi@gmail.com*

Ref

1. T. D. Lee, Some special examples in renormalizable field theory, Phys. Rev. **95**, 1329 (1954).

© 2013 Global Journals Inc. (US)

We make use of the Lie algebraic approach to derive at least one invariant for the TD versions of above coupled oscillator system. In fact, the Lie algebraic approach (ref therein [15]) commands several advantages over the rationalization method, particularly for the TD systems, not only in terms of the closure property of the Poisson bracket algebra of phase space functions but also for its straightforward extension to the corresponding quantum system.

a) *Lie algebraic approach*

If we define

$$x = x_1 + ip_3; \quad y = x_2 + ip_4; \quad p_x = p_1 + ix_3; \quad p_y = p_2 + ix_4. \tag{2}$$

then, a two dimensional real phase space (x, y, p_x, p_y, t), may be transformed into the corresponding extended complex phase plane (ECPS) $(x_1, p_3, x_2, p_4, p_1, x_3, p_2, x_4, t)$ The above transformations add four additional degrees of freedom, (x_3, x_4, p_3, p_4, t), which can make mathematical analysis of a problem a bit more involved. But nevertheless, these type of transformations are used in many studies [12, 13, 14, 13]. From eq.(2) one can easily obtain one can easily obtain

$$\frac{\partial}{\partial x} = \frac{\partial}{\partial x_1} - i\frac{\partial}{\partial p_3}; \quad \frac{\partial}{\partial y} = \frac{\partial}{\partial x_2} - i\frac{\partial}{\partial p_4}; \quad \frac{\partial}{\partial p_x} = \frac{\partial}{\partial p_1} - i\frac{\partial}{\partial x_3}; \quad \frac{\partial}{\partial p_y} = \frac{\partial}{\partial p_2} - i\frac{\partial}{\partial x_4}. \tag{3}$$

Now consider a complex phase space function $I(x, y, p_x, p_y, t)$ as

$$I = I_1(x_1, p_3, x_2, p_4, p_1, x_3, p_2, x_4, t) + iI_2(x_1, p_3, x_2, p_4, p_1, x_3, p_2, x_4, t). \tag{4}$$

Further, the invariance of I implying

$$\frac{dI}{dt} = \frac{\partial I}{\partial t} + [I, H] = 0, \tag{5}$$

where [., .] is the Poisson bracket, which in view of the definition, eq.(2), turns out to be

$$\begin{aligned}[I, H]_{(x,p)} &= [I, H]_{(x_1,p_1)} - i[I, H]_{(x_1,x_3)} - i[I, H]_{(p_3,p_1)} - [I, H]_{(p_3,x_3)} \\ &+ [I, H]_{(x_2,p_2)} - i[I, H]_{(x_2,x_4)} - i[I, H]_{(p_4,p_2)} - [I, H]_{(p_4,x_4)}.\end{aligned} \tag{6}$$

b) *Example*

Consider a coupled harmonic oscillator systems in two-dimensions, whose Hamiltonian is given by (1). Using (2), the above Hamiltonian (see [12] for detail method) can be expressed as

$$\begin{aligned}H &= \frac{\alpha_1}{2}p_1^2 - \frac{\alpha_1}{2}x_3^2 + i\alpha_1 p_1 x_3 + \frac{\alpha_1}{2}p_2^2 - \frac{\alpha_1}{2}x_4^2 + i\alpha_1 p_2 x_4 + \frac{\alpha_2}{2}x_1^2 - \frac{\alpha_2}{2}p_3^2 + i\alpha_2 p_3 x_1 p_4 \\ &+ \frac{\alpha_2}{2}x_2^2 - \frac{\alpha_2}{2}p_4^2 + i\alpha_2 x_2 + \alpha_3(p_1 x_2 + ip_1 p_4 + ix_3 x_2 - x_3 p_4 + x_1 p_2 + ix_1 x_4 + ip_2 p_3 - p_3 x_4) \\ &= \sum_{m=1}^{20} h_m(t)\Gamma_m(x_1, p_3, x_2, p_4, p_1, x_3, p_2, x_4),\end{aligned} \tag{7}$$

and the various Γ's and h(t)'s for the above complex H are given as

$$\begin{aligned}&\Gamma_1 = \frac{p_1^2}{2}; \ \Gamma_2 = \frac{x_3^2}{2}; \ \Gamma_3 = p_1 x_3; \ \Gamma_4 = \frac{p_2^2}{2}; \ \Gamma_5 = \frac{x_4^2}{2}; \ \Gamma_6 = p_2 x_4; \ \Gamma_7 = \frac{x_1^2}{2}; \ \Gamma_8 = \frac{p_3^2}{2}; \\ &\Gamma_9 = x_1 p_3; \ \Gamma_{10} = \frac{x_2^2}{2}; \ \Gamma_{11} = x_2 p_4; \ \Gamma_{12} = \frac{p_4^2}{2}; \ \Gamma_{13} = p_1 x_2; \ \Gamma_{14} = p_1 p_4; \ \Gamma_{15} = x_3 x_2; \\ &\Gamma_{16} = x_3 p_4, \ \ \Gamma_{17} = x_1 p_2; \ \ \Gamma_{18} = x_1 x_4; \ \ \Gamma_{19} = p_3 p_2; \ \ \Gamma_{20} = p_3 x_4.\end{aligned} \tag{8}$$

With

$$\begin{aligned}&h_1 = h_2 = \alpha_1; \ h_3 = i\alpha_1; \ h_4 = h_5 = \alpha_2; \ h_6 = i\alpha_2; \ h_7 = h_8 = \beta_1; \ h_9 = i\beta_1; \ h_{10} = h_{11} = \beta_2 \\ &h_{12} = i\beta_2; \ h_{14} = h_{15} = i\alpha_3; \ h_{13} = \alpha_3; \ h_{16} = -\alpha_3; \ h_{17} = \alpha_3; \ h_{18} = h_{19} = i\alpha_3; \ h_{20} = -\alpha_3.\end{aligned} \tag{9}$$

Ref

14. R. S. Kaushal, On the quantum mechanics of complex Hamiltonian systems in one dimension, J. Phys. A: Math. Gen. **34** (2001) L709-721; R. S. Kaushal and Parthasarathi, Quantum mechanics of complex Hamiltonian systems in one dimension, J. Phys. A **35**, (2002) 8743-8754.

© 2013 Global Journals Inc. (US)

The dynamical algebra in this case is not closed. To find closure property for the above system, we have to add sixteen more phase space functions (Γ_l)'s. The additional (Γ_l)'s are as follow

$$\Gamma_{21} = p_1p_3;\ \Gamma_{22} = p_1x_1;\ \Gamma_{23} = x_2p_1;\ \Gamma_{24} = p_1p_4,\ \Gamma_{25} = p_3x_3;\ \Gamma_{26} = x_1x_3;\ \Gamma_{27} = x_2x_3;\ \Gamma_{28} = x_3p_4;$$
$$\Gamma_{29} = x_2p_2;\ \Gamma_{30} = p_4p_2;\ \Gamma_{31} = p_2p_3;\ \Gamma_{32} = p_2x_1,\ \Gamma_{33} = x_2x_4;\ \Gamma_{34} = x_4p_4;\ \Gamma_{35} = x_1x_4;\ \Gamma_{36} = p_3x_4, \tag{10}$$

with corresponding $h_l(t) = 0$. Now in the light of Poisson bracket (6) for complex systems, we get large number (288 no. of) nonvanishing Poisson brackets (for more detail see [12]). Therefore, their use (5) yields the following set of PDEs in λ's as described in (**??**) section two:

$$\dot{\lambda}_1 = 4\alpha_1(i\lambda_{21} - \lambda_{22}) - 4\alpha_3(\lambda_{23} - i\lambda_{24}), \tag{11}$$

$$\dot{\lambda}_2 = -4\alpha_1(\lambda_{25} + i\lambda_{26}) - 4\alpha_3(i\lambda_{27} + \lambda_{28}), \tag{12}$$

$$\dot{\lambda}_3 = -2\alpha_1(\lambda_{21} + i\lambda_{22} + i\lambda_{25} - \lambda_{26}) - 2\alpha_3(i\lambda_{23} + \lambda_{24} + \lambda_{27} - i\lambda_{28}), \tag{13}$$

$$\dot{\lambda}_4 = -4\alpha_2(\lambda_{29} + i\lambda_{30}) + 4\alpha_3(i\lambda_{31} - \lambda_{32}), \tag{14}$$

$$\dot{\lambda}_5 = -4\alpha_2(\lambda_{34} + i\lambda_{33}) - 4\alpha_3(\lambda_{36} + i\lambda_{35}), \tag{15}$$

$$\dot{\lambda}_6 = -2\alpha_2(i\lambda_{29} + \lambda_{30} - i\lambda_{34} + \lambda_{33}) - 2\alpha_3(\lambda_{31} + i\lambda_{32} + \lambda_{35} - i\lambda_{36}), \tag{16}$$

$$\dot{\lambda}_7 = -4\beta_1(-\lambda_{22} + i\lambda_{26}) + 4\alpha_3(\lambda_{32} - i\lambda_{35}), \tag{17}$$

$$\dot{\lambda}_8 = 4\beta_1(i\lambda_{21} + \lambda_{25}) + 4\alpha_3(\lambda_{36} + i\lambda_{31}), \tag{18}$$

$$\dot{\lambda}_9 = 2\beta_1(\lambda_{21} + i\lambda_{22} - i\lambda_{25} + \lambda_{26}) + 2\alpha_3(\lambda_{31} + i\lambda_{32} + \lambda_{35} - i\lambda_{36}), \tag{19}$$

$$\dot{\lambda}_{10} = 4\alpha_3(\lambda_{23} - i\lambda_{27}) - 4\beta_2(\lambda_{29} + \lambda_{33}), \tag{20}$$

$$\dot{\lambda}_{11} = 2\alpha_3(i\lambda_{23} + \lambda_{24} + \lambda_{27} - i\lambda_{28}) - 2\alpha_3(\lambda_{29} + i\lambda_{30} + \lambda_{34} - i\lambda_{33}), \tag{21}$$

$$\dot{\lambda}_{12} = 4\alpha_3(i\lambda_{24} + \lambda_{28}) - 4\beta_2(\lambda_{30} + \lambda_{34}), \tag{22}$$

$$\dot{\lambda}_{13} = 2\alpha_3(\lambda_{22} - i\lambda_{33} - i\lambda_{26} + \lambda_{29}) + 2\beta_1(i\lambda_{23} - i\lambda_{27}) + 2\beta_2(\lambda_{32} - i\lambda_{35}), \tag{23}$$

$$\dot{\lambda}_{14} = 2\alpha_3(\lambda_{26} + \lambda_{30} - i\lambda_{34} + i\lambda_{22}) + 2\beta_1(\lambda_{24} - i\lambda_{28}) + \beta_2(-\lambda_{32} + i\lambda_{35}), \tag{24}$$

$$\dot{\lambda}_{15} = 2\alpha_3(\lambda_{21} - i\lambda_{25} + \lambda_{33} + i\lambda_{29}) + 2\beta_1(i\lambda_{23} + \lambda_{27}) + \beta_2(\lambda_{31} - \lambda_{36}), \tag{25}$$

$$\dot{\lambda}_{16} = 2\beta_2(i\lambda_{21} + \lambda_{25} + i\lambda_{30} + i\lambda_{34}) + 2\beta_1(i\lambda_{24} + \lambda_{28}) - \beta_2(\lambda_{31} + i\lambda_{36}), \tag{26}$$

$$\dot{\lambda}_{17} = 2\alpha_3(i\lambda_{21} - \lambda_{22} - \lambda_{29} + i\lambda_{30}) + 2\alpha_1(i\lambda_{31} - \lambda_{32}) - 2\alpha_2(i\lambda_{23} + \lambda_{24}), \tag{27}$$

$$\dot{\lambda}_{18} = -2\alpha_3(\lambda_{21} + i\lambda_{22} + i\lambda_{29} + \lambda_{30}) + 2\alpha_1(-\lambda_{31} + i\lambda_{32}) + \alpha_2(-i\lambda_{23} + \lambda_{24}), \tag{28}$$

Ref

12. J.S. Virdi, and S.C. Mishra, Exact complex integrals in two dimensions for shifted harmonic oscillators, Pramana J. Phys. **79** (2012) 19-40; J.S. Virdi, F. Chand, C.N. Kumar, and S.C. Mishra, Complex dynamical invariants for two-dimensional complex potentials, Pramana J. Phys. **79**, (2012) 173-183.

© 2013 Global Journals Inc. (US)

$$\dot{\lambda}_{19} = 2\alpha_3(i\lambda_{25} - \lambda_{26} - i\lambda_{29} - \lambda_{30}) - 2\alpha_1(\lambda_{31} + i\lambda_{32}) + 2\alpha_2(-\lambda_{27} + i\lambda_{28}), \tag{29}$$

$$\dot{\lambda}_{20} = -2\alpha_3(\lambda_{25} + i\lambda_{26} + i\lambda_{29} + \lambda_{30}) - 2\alpha_1(\lambda_{31} + i\lambda_{32}) + 2\alpha_2(-i\lambda_{27} - \lambda_{28},) \tag{30}$$

$$\dot{\lambda}_{21} = 2\beta_1(i\lambda_1 + \lambda_3) + 2\alpha_1(-\lambda_9 + i\lambda_8) + 2\alpha_3(-i\lambda_{16} - \lambda_{15}) + 2\alpha_3(i\lambda_{17} + \lambda_{18}), \tag{31}$$

$$\dot{\lambda}_{22} = 2\beta_1(\lambda_1 - i\lambda_3) + 2\alpha_1(i\lambda_9 + \lambda_7) + 2\alpha_3(i\lambda_{14} - \lambda_{13}) + 2\alpha_3(\lambda_{17} - i\lambda_{18}), \tag{32}$$

$$\dot{\lambda}_{23} = 2\beta_2(i\lambda_{17} - i\lambda_{18}) + 2\alpha_1(-\lambda_{13} + i\lambda_{15}) + 2\alpha_3(-i\lambda_3 + \lambda_1) + 2\alpha_3(i\lambda_{11} - \lambda_{10}), \tag{33}$$

$$\dot{\lambda}_{24} = 2\alpha_3(i\lambda_1 + \lambda_3) + 2\alpha_3(-\lambda_{11} + i\lambda_{12}) + 2\alpha_1(-i\lambda_{16} - \lambda_{14}) + 2\beta_2(\lambda_{18} - \lambda_{17}), \tag{34}$$

$$\dot{\lambda}_{25} = 2\beta_1(-\lambda_2 - i\lambda_3) - 2\alpha_1(i\lambda_9 + \lambda_8) + 2\alpha_3(-\lambda_{16} - i\lambda_{15}) + 2\alpha_3(i\lambda_{19} + \lambda_{20}), \tag{35}$$

$$\dot{\lambda}_{26} = 2\beta_1(-i\lambda_2 + \lambda_3) + 2\alpha_1(i\lambda_7 - \lambda_9) + 2\alpha_3(-i\lambda_{13} - \lambda_{14}) + 2\alpha_3(\lambda_{19} - i\lambda_{20}), \tag{36}$$

$$\dot{\lambda}_{27} = 2\alpha_3(-i\lambda_2 + \lambda_3) + 2\alpha_3(-i\lambda_{10} - \lambda_{11}) - 2\alpha_1(i\lambda_{13} + \lambda_{15}) + 2\beta_2(\lambda_{19} + i\lambda_{20}), \tag{37}$$

$$\dot{\lambda}_{28} = 2\alpha_3(\lambda_2 + i\lambda_3) + 2\alpha_3(-i\lambda_{11} - i\lambda_{12}) - 2\alpha_1(\lambda_{14} - i\lambda_{16}) + 2\beta_2(-\lambda_{19} + i\lambda_{20}), \tag{38}$$

$$\dot{\lambda}_{29} = 2\beta_2(\lambda_4 + \lambda_6) + 2\alpha_2(-\lambda_{10} + i\lambda_{11}) + 2\alpha_3(i\lambda_{15} - \lambda_{13}) + 2\alpha_3(\lambda_{17} - i\lambda_{19}), \tag{39}$$

$$\dot{\lambda}_{30} = -2\alpha_3(\lambda_4 + i\lambda_6) + 2\alpha_2(-\lambda_{11} - i\lambda_{12}) + 2\alpha_3(-i\lambda_{16} - i\lambda_{14}) + 2\alpha_3(i\lambda_{17} + \lambda_{19}), \tag{40}$$

$$\dot{\lambda}_{31} = 2\alpha_3(i\lambda_4 + \lambda_6) + 2\alpha_2(i\lambda_8 + \lambda_9) + 2\alpha_2(-i\lambda_{16} - \lambda_{15}) + 2\beta_1(i\lambda_{17} + \lambda_{19}), \tag{41}$$

$$\dot{\lambda}_{32} = 2\alpha_3(\lambda_4 - i\lambda_6) + 2\alpha_3(\lambda_7 + i\lambda_9) + 2\alpha_2(i\lambda_{14} - \lambda_{13}) + 2\beta_1(\lambda_{17} - i\lambda_{19}), \tag{42}$$

$$\dot{\lambda}_{33} = 2\beta_2(\lambda_5 + \lambda_6) + 2\alpha_2(-\lambda_{10} - i\lambda_{11}) + 2\alpha_3(-i\lambda_{13} - \lambda_{15}) + 2\alpha_3(\lambda_{18} - i\lambda_{20}), \tag{43}$$

$$\dot{\lambda}_{34} = 2\beta_2(-\lambda_5 - \lambda_6) + 2\alpha_2(-i\lambda_{11} + \lambda_{12}) + 2\alpha_3(-\lambda_{14} - i\lambda_{16}) + 2\alpha_3(i\lambda_{18} + \lambda_{20}), \tag{44}$$

$$\dot{\lambda}_{35} = 2\beta_1(-\lambda_{18} - \lambda_{20}) + 2\alpha_2(-i\lambda_{13} - \lambda_{14}) + 2\alpha_3(-i\lambda_5 + \lambda_6) + 2\alpha_3(i\lambda_7 - \lambda_9), \tag{45}$$

$$\dot{\lambda}_{36} = 2\alpha_3(\lambda_5 + i\lambda_6) - 2\alpha_3(i\lambda_9 + \lambda_8) + 2\alpha_2(-\lambda_{16} + i\lambda_{15}) + 2\beta_1(i\lambda_{18} + \lambda_{20}). \tag{46}$$

As such the solution to these 36 coupled equation turns out to be difficult. Therefore, we make the following choices about λ's which facilitate to find solutions of above equations.
From eqs.(11), (12) and (13), we get $2\dot{\lambda}_3 = i\dot{\lambda}_1 - i\dot{\lambda}_2$. If we consider $\lambda_3 = c_3$ (a constant), and by taking $\lambda_1 = \lambda_2 = \eta_1(t)$; which immediately gives

$$\lambda_1 = \eta_1(t) + c_1; \qquad \lambda_2 = \eta_1(t) + c_2. \tag{47}$$

From eqs.(14), (15) and (16), we get $2\dot{\lambda}_6 = i\dot{\lambda}_4 - i\dot{\lambda}_5$. If we consider $\lambda_6 = c_6$ (a constant), and by taking $\lambda_4 = \lambda_5 = \eta_2(t)$; which immediately gives

$$\lambda_4 = \eta_2(t) + c_4; \qquad \lambda_5 = \eta_2(t) + c_5. \tag{48}$$

Notes

© 2013 Global Journals Inc. (US)

Notes

Again From eqs.(17), (18) and (19), we get $2\dot{\lambda}_9 = i\dot{\lambda}_7 - i\dot{\lambda}_8$. If we set $\lambda_9 = c_9$ (a constant), and consider $\lambda_7 = \lambda_8 = \eta_3(t)$; which immediately gives

$$\lambda_7 = \eta_3(t) + c_7; \qquad \lambda_8 = \eta_3(t) + c_8. \tag{49}$$

From eqs.(20), (21) and (22), we get $2\dot{\lambda}_{11} = i\dot{\lambda}_{10} - i\dot{\lambda}_{12}$. If we set $\lambda_{11} = c_{11}$ (a constant), and consider $\lambda_{12} = \lambda_{10} = \eta_4(t)$; which immediately gives

$$\lambda_{12} = \eta_4(t) + c_{12}; \qquad \lambda_{10} = \eta_4(t) + c_{10}. \tag{50}$$

Now, in order to find solutions for λ_{13}, λ_{14}, λ_{15} and λ_{16} we have to make simplification for complications of above set of 24 eqs (23-46). (i.e. $\alpha_1 = \alpha_2 = \alpha_3$, and $\beta_1 = \beta_2 = \alpha_3$). From eqs.(17), (20) and (23), we get $2\dot{\lambda}_{13} = \dot{\lambda}_7 + i\dot{\lambda}_{10}$. If we consider $\lambda_{13} = c_{13}$ (a constant), and considering the eqn from above relation (with $\lambda_{10} = \eta_4(t) + c_{10},\ \lambda_7 = \eta_3(t) + c_7$;) gives

$$\lambda_{13} = \eta(t) + c_{13}. \tag{51}$$

where $\eta(t) = \frac{1}{2i}[\eta_4(t) + \eta_3(t);]$ is an another function of time, and ($c_{13} = c_7 + c_{10}$; a constant.)
From eqs.(18), (20) and (25), we get $2i\dot{\lambda}_{15} = \dot{\lambda}_{10} + i\dot{\lambda}_8$, If we consider $\lambda_{15} = c_{15}$ (a constant), and considering the eqn from above relation (with $\lambda_{10} = \eta_4(t) + c_{10},\ \lambda_8 = \eta_3(t) + c_8$;) gives

$$\lambda_{15} = \eta(t) + c_{15}, \tag{52}$$

where $\eta(t) = \frac{1}{2i}[\eta_4(t) + \eta_3(t)]$ is an another function of time and $c_{15} = c_{10} + c_8$; a constant.
In order to find solutions for λ_{16}, from eqs.(18), (22) and (26), we get $2i\dot{\lambda}_{16} = \dot{\lambda}_8 + \dot{\lambda}_{12}$. If we consider $\lambda_{16} = c_{16}$; (a constant), and considering the relation (with $\lambda_8 = \eta_3(t) + c_8,\ \lambda_{12} = \eta_4(t) + c_{12}$;) gives

$$\lambda_{16} = \eta(t) + c_{16}, \tag{53}$$

where $\eta(t) = \frac{1}{2i}[\eta_4(t) + \eta_3(t)]$; and $c_{16} = c_8 + c_{12}$; a constant.
From eqs.(17), (22) and (24), we get $2i\dot{\lambda}_{14} = \dot{\lambda}_7 + \dot{\lambda}_{12}$, If we consider $\lambda_{14} = c_{14}$; (a constant), and considering relation (with $\lambda_7 = \eta_3(t) + c_7,\ \lambda_{12} = \eta_4(t) + c_{12}$;) gives

$$\lambda_{14} = \eta(t) + c_{14}, \tag{54}$$

where $\eta(t) = \frac{1}{2i}[\eta_4(t) + \eta_3(t)]$; and $c_{16} = c_7 + c_{12}$ a constant.
Now, to find solutions for λ_{17}, λ_{18}, λ_{19} and λ_{20}, refer from eqs.(11), (14) and (27), we get $2\dot{\lambda}_{17} = \dot{\lambda}_1 + \dot{\lambda}_4$; and considering the relation $\lambda_1 = \eta_1(t) + c_1,\ \ \lambda_4 = \eta_2(t) + c_4$; gives

$$\lambda_{17} = \phi(t) + c_{17}, \tag{55}$$

where $\phi(t) = \frac{1}{2}\int[\dot{\eta}_1(t) + \dot{\eta}_2(t)]dt$; and $c_{17} = c_1 + c_4$, a constant. From eqs.(11), (15) and (28), we get $2\dot{\lambda}_{18} = i\dot{\lambda}_1 - i\dot{\lambda}_5$; and considering the relation $\lambda_1 = \eta_1(t) + c_1,\ \lambda_5 = \eta_2(t) + c_5$; will results

$$\lambda_{18} = \varphi(t) + c_{18}, \tag{56}$$

where $\varphi(t) = \frac{1}{2}\int[i\dot{\eta}_1(t) - i\dot{\eta}_2(t)]dt$; and $c_{18} = c_1 + c_5$, a constant. Similarly to find solutions of λ_{19}, from eqs.(12), (14) and (29), we get $2\dot{\lambda}_{19} = i\dot{\lambda}_2 - i\dot{\lambda}_4$; and considering the relation $\lambda_2 = \eta_1(t) + c_2,\ \ \lambda_4 = \eta_2(t) + c_4$; gives

$$\lambda_{19} = \chi(t) + c_{19}, \tag{57}$$

where $\chi(t) = \frac{1}{2}\int[i\dot{\eta}_1(t) - i\dot{\eta}_2(t)]dt$ and $c_{19} = c_2 + c_4$, a constant.
Again from eqs.(12), (15) and (30), we get $2\dot{\lambda}_{20} = \dot{\lambda}_2 + \dot{\lambda}_4$; and considering the relation $\lambda_2 = \eta_1(t) + c_1,\ \ \lambda_4 = \eta_2(t) + c_4$; gives

$$\lambda_{20} = \quad (t) + c_{20}, \tag{58}$$

where $\quad (t) = \frac{1}{2}\int[\dot{\eta}_1(t) + \dot{\eta}_2(t)]dt$; and $c_{20} = c_2 + c_4$, a constant.
Solutions for $(\lambda_{21} - \lambda_{28})$ can be obtained respectively as from eqs.(31-38), we obtain following equations

© 2013 Global Journals Inc. (US)

$$i\dot{\lambda}_{21} + \dot{\lambda}_{22} = 2(\alpha_1\lambda_7 - \alpha_1\lambda_8 - \alpha_3\lambda_{13} + i\alpha_3\lambda_{14} - i\alpha_3\lambda_{15} + \alpha_3\lambda_{16}) = 0. \tag{59}$$

$$\dot{\lambda}_{23} + i\dot{\lambda}_{24} = 2(-\alpha_3\lambda_{10} + \alpha_3\lambda_{12} - \alpha_1\lambda_{13} + i\alpha_1\lambda_{15} - i\alpha_1\lambda_{14} + \alpha_1\lambda_{16}) = 0. \tag{60}$$

$$\dot{\lambda}_{25} - i\dot{\lambda}_{26} = 2(\alpha_1\lambda_7 - \alpha_1\lambda_8 - \alpha_3\lambda_{13} + i\alpha_3\lambda_{14} - i\alpha_3\lambda_{15} + \alpha_3\lambda_{16}) = 0. \tag{61}$$

$$\dot{\lambda}_{27} + i\dot{\lambda}_{28} = 2i(-\alpha_3\lambda_{10} + \alpha_3\lambda_{12} - \alpha_1\lambda_{13} + i\alpha_1\lambda_{15} - i\alpha_1\lambda_{14} + \alpha_1\lambda_{16}) = 0. \tag{62}$$

Notes

Since

$$\lambda_7 = \lambda_8, \lambda_{10} = \lambda_{12}; \qquad \lambda_{13} = \lambda_{14} = \lambda_{15} = \lambda_{16} = \eta(t), \tag{63}$$

or if we set

$$\begin{aligned} &\dot{\lambda}_{21} = -i\dot{\lambda}_{22} = \dot{\xi}(t); \quad \dot{\lambda}_{23} = -i\dot{\lambda}_{24} = \dot{\theta}(t); \\ &\dot{\lambda}_{25} = i\dot{\lambda}_{26} = \dot{\delta}(t); \quad \dot{\lambda}_{27} = -i\dot{\lambda}_{28} = \dot{\zeta}(t). \end{aligned} \tag{64}$$

which immediately gives

$$\begin{aligned} &\lambda_{21} = \xi(t) + c_{21}; \quad \lambda_{22} = -i\xi(t) + c_{22}; \quad \lambda_{23} = \theta(t) + c_{23}; \quad \lambda_{24} = -i\theta(t) + c_{24}; \\ &\lambda_{25} = \delta(t) + c_{25}; \quad \lambda_{26} = -i\delta(t) + c_{26}; \quad \lambda_{27} = \zeta(t) + c_{27}; \quad \lambda_{28} = i\zeta(t) + c_{28}. \end{aligned} \tag{65}$$

Solutions for $(\lambda_{29} - \lambda_{36})$ can be obtained respectively as, from eqs.(39-46), we obtain following equations

$$\dot{\lambda}_{31} - i\dot{\lambda}_{32} = -2(-i\alpha_3\lambda_7 + i\alpha_3\lambda_8 + i\alpha_2\lambda_{13} + \alpha_2\lambda_{14} - \alpha_2\lambda_{15} - i\alpha_2\lambda_{16}) = 0. \tag{66}$$

$$\dot{\lambda}_{29} + i\dot{\lambda}_{30} = 2(-\alpha_2\lambda_{10} + \alpha_2\lambda_{12} - \alpha_3\lambda_{13} + i\alpha_3\lambda_{15} - i\alpha_3\lambda_{14} + \alpha_3\lambda_{16}) = 0. \tag{67}$$

$$\dot{\lambda}_{33} + i\dot{\lambda}_{34} = 2i(-\alpha_2\lambda_{10} + \alpha_2\lambda_{12} - \alpha_3\lambda_{13} + i\alpha_3\lambda_{15} - i\alpha_3\lambda_{14} + \alpha_3\lambda_{16}) = 0. \tag{68}$$

$$\dot{\lambda}_{35} + i\dot{\lambda}_{36} = -2(-i\alpha_3\lambda_7 + i\alpha_3\lambda_8 + i\alpha_2\lambda_{13} + \alpha_2\lambda_{14} - \alpha_2\lambda_{15} - i\alpha_2\lambda_{16}) = 0. \tag{69}$$

Since

$$\lambda_7 = \lambda_8, \lambda_{10} = \lambda_{12}; \qquad \lambda_{13} = \lambda_{14} = \lambda_{15} = \lambda_{16} = \eta(t), \tag{70}$$

or if we set

$$\dot{\lambda}_{29} = -i\dot{\lambda}_{30} = \dot{\gamma}(t); \quad \dot{\lambda}_{31} = i\dot{\lambda}_{32} = \dot{\mu}(t); \quad \dot{\lambda}_{33} = -i\dot{\lambda}_{34} = \dot{\rho}(t); \quad \dot{\lambda}_{35} = -i\dot{\lambda}_{36} = \dot{\sigma}(t). \tag{71}$$

Which immediately gives

$$\begin{aligned} &\lambda_{29} = \gamma(t) + c_{29}; \quad \lambda_{30} = i\gamma(t) + c_{30}; \quad \lambda_{31} = \mu(t) + c_{31}; \quad \lambda_{32} = -i\mu(t) + c_{32}; \\ &\lambda_{33} = \rho(t) + c_{33}; \quad \lambda_{34} = i\rho(t) + c_{34}; \quad \lambda_{35} = \sigma(t) + c_{35}; \quad \lambda_{36} = i\sigma(t) + c_{36}. \end{aligned} \tag{72}$$

where the arbitrary function η's, $\phi, \varphi, \chi, \psi, \xi, \theta, \delta, \zeta, \gamma, \mu, \rho$ and σ's and integration constants, c_i's, $(i = 1, ...36)$. are obtained through the equations, when we have solved eqs.[(11)-(46)]. Therefore, substitution of the solutions for λ_i's yield the, complex invariant for a two dimensional complex oscillator systems as

$$\begin{aligned} I = &\frac{1}{2}\eta_1(p_1^2 + x_3^2) + \frac{1}{2}\eta_2(p_2^2 + x_4^2) + \frac{1}{2}\eta_4(x_1^2 + p_3^2) + \frac{1}{2}\eta_3(x_2^2 + p_4^2) + \eta(x_1x_3 + p_3p_4 + x_2p_3 + x_1p_4) \\ &+\phi p_1p_2 + \varphi p_1x_4 + \chi x_3p_2 + \quad x_3x_4 + \xi(p_1p_3 - ip_1x_1) + \theta(p_1x_2 - ip_1p_4) + \delta(p_3x_3 - ix_1x_3) \\ &+\zeta(ip_4x_3 + x_2x_3) + \gamma(ip_2p_4 + p_2x_2) + \mu(p_3p_2 - ix_1p_2) + \rho(ip_4x_4 + x_2x_4) + \sigma(ip_3x_4 + x_1x_4). \end{aligned} \tag{73}$$

is our desired invariant.

© 2013 Global Journals Inc. (US)

II. Conclusion

We have shown in this paper, the coupled harmonic oscillator system with two space and one time variable share the same mathematical framework as the coupled harmonic oscillators in one dimension ECPS. The role of a linear invariant designed, however, for a rotating TD harmonic oscillator in N-dimensions is investigated by Malkin and Man'ko [16] in the context of coherent states. While the use of the quantum analogue of such TD systems in one dimension has been known [17, 18] for more than three decades.

Notes

References Références Referencias

1. T. D. Lee, Some special examples in renormalizable field theory, Phys. Rev. **95**, 1329 (1954).
2. H. Van Dam, Y. J. Ng, and L. C. Biedenharn, A comment on fermionic tachyons and Poincre representations, Phys. Lett. B **158**, (1985) 227-258.
3. J. Fu, S. Jimnez, Y. Tang, L. Vzquez, Construction of exact invariants of time-dependent linear nonholonomic dynamical systems, J. Phys. A: Math. Gen. **372**, (2008) 10-23.
4. H Goldstien, C Poole and J Safco, "*Classical mechanics*" (Pearson Edu. Inc., Singapore, 2002).
5. L. A. Pipes and L. R. Harvill, "*Applied Mathematics for Engineers and Physicists*" (New York: McGraw-Hill: 1970).
6. B. Yurke and D. Stoler, Bell's-inequality experiment employing four harmonic oscillators Phys. Rev. A **51**, (1995) 3437-3454.
7. L. O. Chua, C. A. Desoer, E. S. Kuh, "*Linear and Nonlinear Circuits*" (Singapore: McGraw-Hill).
8. M. Blasone, P. Jizba and G. Vitiello, "*Quantum Field Theory and Its Macroscopic Manifestations*" (Imperial collge press: London 2011).
9. M. S. Abdalla, A.I. Elkasapy, Lie algebraic approach of a charged particle in presence of a constant magnetic field via the quadratic invariant, Annals of Physics **325**, (2010) 16671678; R.K. Colegrave, P. Croxson and M.A. Mannan, Complex invariants for the time-dependent harmonic oscillator, Phy. Lett. A. **131** (1988) 407-411: J.S. Virdi, F. Chand, C.N. Kumar, and S.C. Mishra, Complex dynamical invariants for two-dimensional nonhermitian Hamiltonian systems, Can. Jour. Phys. **90**, (2012) 151-157.
10. Kusenko A. and Shrock R., General determination of phases in quark mass matrices, Phy Lett B **323**, (1994) 18-24; Phys. Rev. D **50**, (1994) R30-54.
11. J. F. Gunion and H. E. Haber, Conditions for CP violation in the general two-Higgs-doublet model, Phys. Rev. D **72**, (2005) 095002-095117.
12. J.S. Virdi, and S.C. Mishra, Exact complex integrals in two dimensions for shifted harmonic oscillators, Pramana J. Phys. **79** (2012) 19-40; J.S. Virdi, F. Chand, C.N. Kumar, and S.C. Mishra, Complex dynamical invariants for two-dimensional complex potentials, Pramana J. Phys. **79**, (2012) 173-183.
13. R. S. Kaushal and H J Korsch, Some remarks on complex Hamiltonian systems, Phys. Lett. A **276**, (2000) 47-57; R. S. Kaushal and Singh S, Construction of complex invariants for classical dynamical systems, Ann. Phys. **288**, (2001) 253-274; R. S. Kaushal and Singh S,, Complex dynamical invariants for one-dimensional classical systems, Phys. Scr. **67**, (2003) 181-186; R. S. Kaushal, Classical and quantum mechanics of complex hamiltonian systems: An extended complex phase space approach, Pramana J. Phys. **73**, (2009) 287-299.
14. R. S. Kaushal, On the quantum mechanics of complex Hamiltonian systems in one dimension, J. Phys. A: Math. Gen. **34** (2001) L709-721; R. S. Kaushal and Parthasarathi, Quantum mechanics of complex Hamiltonian systems in one dimension, J. Phys. A **35**, (2002) 8743-8754.
15. R. S. Kaushal, "*Classical and Quantum Mechanics of Noncentral Potentials*": A Survey of Two-Dimensional Systems" (New Delhi: Narosa and Berlin: Springer 1998).
16. I. A. Malkin and V. I. Man'ko, Linear adiabatic invariants and coherent states, Phys. Lett. A **22**, (1970) 243-254; I. A. Malkin and V. I. Man' ko, and D. A. Trifonov, Coherent states and transition probabilities in a time-dependent electromagnetic field, Phys. Rev. D **2**, (1970) 1371-1184.
17. H. N. Hollinhorst, Quantum limits on resonant-mass gravitational-radiation detectors, Phys. Rev. D **19**, (1979) 1669-1685.
18. V. V. Dodonov, I. A. Malkin and V. I. Man'ko, Even and odd coherent states and excitations of a singular oscillator, Physica **72**, (1974) 597-612; Int. Journ. Theo. Phys. **14**, (1975) 37-56.

© 2013 Global Journals Inc. (US)

Notes

This page is intentionally left blank

© 2013 Global Journals Inc. (US)

Global Journal of Science Frontier Research
Mathematics and Decision Sciences
Volume 13 Issue 7 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Sensitivity Analysis for the Physiographic Parameters of Overland as well as Cannel Flow Elements Based on Kinematic Wave (Kw) Theory

By Dr. M. M. Hossain, J. Ferdous & M. Motateb Hossain
University of Dhaka, Bangladesh

Abstract - In the recent past, in many developing countries a good deal of research has been carried to solve the problems of 'large basins' whereas not much has been done with regard to the hydrologic problems of small watersheds. Small watersheds to play important roles e.g. a village pond catered by its own small watershed; in hilly watersheds, the generated runoff causes flash flood, resulting into disruptions of communication lines etc. Therefore, it is necessary to look into these aspects of the hydrologic problems with greater attention. The hydrologic responses of a small watershed depend upon the mechanics of surface runoff, which is primarily a nonlinear process. We have discussed the sensitivity of physiographic parameters through lumped kinematics wave models and found that the overland roughness and overland slope are more sensitive than other physiographic parameters of overland and channel flows. Appropriate discussions of results and conclusions as arrived at in different parts have been summarized.

Keywords : kinematic wave, overland flow, channel flow, lumped physiographic models, watershed, sensitivity analysis.

GJSFR-F Classification : MSC 2010: 53A17

SENSITIVITY ANALYSIS FOR THE PHYSIOGRAPHIC PARAMETERS OF OVERLAND AS WELL AS CANNEL FLOW ELEMENTS BASED ON KINEMATIC WAVE KWTHEORY

Strictly as per the compliance and regulations of :

© 2013. Dr. M. M. Hossain, J. Ferdous & M. Motateb Hossain. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Sensitivity Analysis for the Physiographic Parameters of Overland as well as Cannel Flow Elements Based on Kinematic Wave (Kw) Theory

Dr. M. M. Hossain[α], J. Ferdous [σ] & M. Motateb Hossain [ρ]

Abstract - In the recent past, in many developing countries a good deal of research has been carried to solve the problems of 'large basins' whereas not much has been done with regard to the hydrologic problems of small watersheds. Small watersheds to play important roles e.g. a village pond catered by its own small watershed; in hilly watersheds, the generated runoff causes flash flood, resulting into disruptions of communication lines etc. Therefore, it is necessary to look into these aspects of the hydrologic problems with greater attention. The hydrologic responses of a small watershed depend upon the mechanics of surface runoff, which is primarily a nonlinear process. We have discussed the sensitivity of physiographic parameters through lumped kinematics wave models and found that the overland roughness and overland slope are more sensitive than other physiographic parameters of overland and channel flows. Appropriate discussions of results and conclusions as arrived at in different parts have been summarized.

Keywords : kinematic wave, overland flow, channel flow, lumped physiographic models,watershed, sensitivity analysis.

I. Introduction

Water is one of the most important natural resources vitally needed for sustenance of life. The use of water is continuously increasing with the increase in population. This increased demand for water has forced the planners, scientists and hydrologist not only to concentrate on major 'basin planning' but also to pay sufficient attention to small watersheds. The watershed planning may help in micro-level water budgeting to meet the localized demands of water for small population concentrations scattered over the entire basin. In Bangladesh, in majority cases, each village has its own pond (or tank) from which the water is drawn for domestic and other needs. These ponds have their own watersheds from which the rainwater is fed to them. Prior to onset of monsoon, after the sustained dry period most of the ponds either get dries up or are left with insignificant quantities of water. The fine silt or clay, which is accumulated at the bottom of the tank, is removed by the local people for plastering of huts and other general repairs of the houses. Thus, the lost capacity of the pond due to accumulation of silt is recouped. Though is may appear quite insignificant, yet the role of small watersheds is very important is most of the developing tropical counties is general and in the Indian subcontinent is particular. The micro-climatic influences onto the small watersheds are of

Authors α σ ρ : Department of Mathematics, University of Dhaka, Dhaka-1000, Bangladesh. E-mail : ema38000@gmail.com

© 2013 Global Journals Inc. (US)

particular significance. Response of a watershed to small, medium heavy and very heavy rainfall leasing to near cloud burst conditions is of importance not only for meeting the water demands but also for the protection of life and property. In hilly and mountainous catchments, the situations may get worse. Heavy intensity rains and resulting runoffs may cause flash floods and landslides and thus disrupt the communication lines of vital importance. These disruptions may result into major economic problems as one region gets completely cutoff from rest of the country.

Notes

To find suitable solutions to these problems, it is necessary that micro-level in-depth studies be carried out for computations of watershed responses to vastly varying meteorological conditions. There are different types of watershed models, which have been developed in the past to estimate the peak flows and runoff hydrographs for small watersheds. With the advent of powerful high-speed digital computers, it is now possible to study through mathematical models the basic physical processes viz. transformation of rain into the runoff land also the movement of water in the surface as well as in channels. These models rely on mathematical statements to represent the system.

In a small watershed, the generation of runoff can better be studied through the 'overland phase' and 'channel phase'. The conversion of rainfall excess into surface runoff and then the movement of flood flows in the channel is recognized as a nonlinear process. These days this process is generally modeled through two approaches, viz. the hydrodynamic approaches and the system's approach. For small watersheds the hydrodynamic approaches have been found more useful for the transformation of rainfall excess into the surface runoff (viz. in the overland phase) and the movement of flood flows in the channel (viz. in the channel phase).

a) Objective of the Study

The objectives of the present work may thus be summarized as below:

(i) To develop suitable mathematical models based on KW theory, best suited to the typical problems of some small watersheds.

(ii) To confirm the applicability of finite difference schemes which are most suitable for small watersheds under study.

(iii) To carry out the sensitivity analysis for the physiographic parameters of overland as well as cannel flow elements and to identify the parameters which are most sensitive. Further, to find out the role of effective overland roughness in the context of natural high slope watersheds.

Computer can solve only a finite number of digits to represent each number at each step of a calculation. Hence, round-off errors are incorporated. The computation is stable if the growth of these errors is within reasonable limits or controlled.

A numerical model with consistent equations, convergent solutions and stable error propagation forms a computationally stable scheme and gives results, which are quite close to the exact solution.

b) Nature of Hydrologic System

The nature of the hydrologic system is completely defined, through one property each, mentioned in the following three sets of possible system behaviors.

(1) Linear or nonlinear

(2) Lumped or distributed

(3) Time-invariant or time-variant.

A brief description of these three characteristics is given in the following subsections.

© 2013 Global Journals Inc. (US)

Notes

i. *The Linear and Nonlinear Systems*

A hydrologic system is said to be linear, if a step input fed to the system produces an output, which is directly proportional to their input. A linear system may be represented mathematically by a linear equation. The principle of superposition applies to it may be defined mathematically as under:

$$f(\Psi_1)_t + f(\Psi_2)_t + --- + f(\Psi_n)_t = f(\Psi_1 + \Psi_2 + --- + \Psi_n)_t$$

where Ψ_i is any response function and t is the time of its applicability.

The principle of homogeneity, which is a particular case of principle of superposition, thus may be described as below $f(k\Psi)_t = kf(\Psi)_t$ where k is a constant function.

A nonlinear system is represented by a nonlinear function. The extent of nonlinearity depends upon the system itself.

The hydrologic system may be defined by a general type of differential equation and can be written as follows:

$$f(\Psi) = a_n \frac{d^n \Psi}{dt^n} + a_{n-1} \frac{d^{n-1} \Psi}{dt^{n-1}} + --- + a_1 \frac{d^1 \Psi}{dt^1} + a_0 \Psi .$$

The system would be nonlinear if any on these coefficients a_0, a_1, a_2, - - -, a_n etc. are the function of Ψ or the function Ψ carries an exponent other than unity.

ii. *The Lumped and Distributed System*

The hydrologic system is said to be lumped if its input functions or parameters do not vary with respect to spatial coordinates (Fig.2.3 (a)). For the lumped systems, average conditions or values of input and parameters are applicable. Thus, lumped systems are represented by ordinary differential equations.

The system is defined as distributed if the input or the transfer function and other parameters do vary with the spatial coordinates (Fig.2.3 (b)). Such systems are mathematically represented by the partial differential equations. The theoretical solution of such systems (a differential equation) this requires completes knowledge of the boundary conditions.



(a) LUMPED SYSTEM

$\Phi(x)$

© 2013 Global Journals Inc. (US)



(b) DISTRIBUTED SYSTEM

Figure : The Lumped and Distributed Systems

iii. *Time-Invariant and Time-Variant Systems*

A time-variant system is one in which the input-output relationship is not dependent upon the time at which the input is applied to the system. The concept of time-invariant makes the analysis simpler.

A time-variant system is one in which the input-output relationship is a dependent function of time. It may be concluded that a lumped, linear and time-invariant system is easiest to work with. But in surface hydrology, the hydrologic system happens to be distributed, nonlinear and time-variant in its behavior. This behavior of the system is the most complex one and quite difficult not only to formulate mathematically but also to solve. Therefore, a compromise has to be reached so that the complicated natural hydrologic system may be solved satisfactorily by making suitable assumptions with respect to the aspects stated above.

c) Numerical Technique (Finite Difference Methods) for Solving the Kinematic Wave Equations

The computational schemed will require the initial values for the entire domain (X_i's) and the upstream boundary conditions for all (t_j's). Solutions of the governing equations through the finite difference scheme will be obtained at each of this grid point. Computations advance along the downstream direction for a time step Δt, until all the discharges as well as water areas are computed at all the grid points in the entire longitudinal length L.

Next, the computations are advance ahead in time by another time step Δt and the computations proceed likewise.

The rainfall excess intensity (i_e) is assumed constant within a time step Δt. But it may change from one time step to the next time step, to account for the variation in rates of rainfall excess intensities occurring within a storm event.

Notes

© 2013 Global Journals Inc. (US)

Notes



Figure 3.3 : Flow Domain Representation Through a Fixed X− t Grid

d) Physiographic Models Employed

i. *Lumped Physiographic Model*



Figure 3.5 : Lumped Physiographic Models

© 2013 Global Journals Inc. (US)

a. Initial Conditions

$$\left.\begin{aligned} Q(x,0) &= 0 \\ A(x,0) &= 0 \end{aligned}\right\} \text{for all x}$$

For the overland flows,

$$\left.\begin{aligned} q(x,0) &= 0 \\ y_0(x,0) &= 0 \end{aligned}\right\} \text{for all x}$$

b. Boundary Conditions

For channel flow:

$$\left.\begin{aligned} Q(0,t) &= 0 \\ A(0,t) &= 0 \end{aligned}\right\} \text{for all t}$$

And for the overland flow:

$$\left.\begin{aligned} q(0,t) &= 0 \\ y_0(0,t) &= 0 \end{aligned}\right\} \text{for all t}$$

e) Physiographic Details of Watershed of Railway Bridge No. 319.

The small watershed of Railway Bridge No. 319 selected for this study is situated in Bangalore district of Karnataka of India. This bridge is located on Arsikere Bangalore Section of the Indian Railways. The index maps giving details of the watershed is given in Fig. Some important features, general information and the physiographic data as extracted from the available records and the topographic map of this bridge is given in Table.

Table 1 : General Information and Salient Features of Watershed of Railway Bridge No. 319

Sl. No.	Particulars	Units
1.	a. Name of the zone b. Name of the sub zone	Deccan Plateau 3i
2.	Geographical location	i. Longitude(app.) $77^0 10'$ East ii. Latitude (app.) $13^0 18'$ North
3.	Terrain	Hilly
4.	Shape of the basin	Oblong
5.	Climate	Humid
6.	Type of soil	i. Rocky ii. Red Earth

Notes

© 2013 Global Journals Inc. (US)

Notes

7.	Land use	Partially Dry cultivation
8.	No. of rain gauge station	1
9.	No. of discharge location	1
10.	Altitude (average elevation from MSL)	833 meter
11.	Watershed area	82.0 hectare
12.	Length of the main channel(longest)	1650.0metre
13.	Total average annual rainfall	500-1000(mm)



Figure 4.1 : The Index Map of Bridge no. 319

f) Estimation of Physiographic Parameters for Bridge No. 319

The topographic details of this watershed are shown in Fig. 4.1. In this natural watershed, only one main drainage channel exists that too in the central part of the watershed. As discussed the lumped physiographic model is used to compute the model parameters for the application of KW theory. For this purpose, a lumped model of the type given in Fig. (a) is adopted for the estimation of parameters. The equivalent watershed has been obtained by dividing equally the total drainage area into the two sides 1650 meters long main channel. The schematic representation of this model is shown in Fig.

As discussed in Section, the physiographic parameters were computed. The values of physiographic parameters are given in Table.

© 2013 Global Journals Inc. (US)



Figure 4.2 : Schmatic Representation Through a Lumped Model (Bridge No. 319)

Table 2 : The Lumped Physiographic Parameter Values of Bridge No. 319

Sl.No.	Particular	Unit
1.	Area	82.0 hectares
2.	Overland (Plane): (a). Average length (each side) (b). Average slope (each side) (c). Overland roughness	 248.0 meters 0.092 0.052
3.	Channel: (a). Average length (b). Average slope (c). Average roughness (d). Average bed width (e). Average side slope	 1650.0 meters 0.072 0.035 3.0 meters 2.5 H: IV

g) Overland Flow of Watershed of Bridge No. 319 for Different Values of Overland Roughness (An)

In overland there are two physiographic parameters land roughness and slope. The effect of overland roughness on overland flow of watershed of bridge no. 319 is shown in the following Fig.

Notes

© 2013 Global Journals Inc. (US)

Notes



Figure : Overland Flow for Different Values of Overland Rougheness

From the above figure, we observed that if the overland roughness is increasing then peak flow is decreasing. Therefore, the overland roughness (AN) is sensitive.

h) Overland Flow of Watershed of Bridge No. 319 for Different Values of Overland Slop (So)

Different overland has different overland slope. It has an effect on overland flow. The effect of overland slope on overland flow of watershed of bridge no. 319 is shown in the following Fig. 4.4



Figure : Overland Flow for Different Values of Overland Slope

© 2013 Global Journals Inc. (US)

From the above figure, we observe that if the overland slope is increasing then the peak flow is increasing for high slope. Therefore, the overland slope is sensitive.

i) Channel Flow Routing

A channel is a conduit in which water flows with a free surface. Classified according to its origin a cannel may be either natural or artificial. Natural channels include all watercourse that exist naturally on the earth, varying is size from tiny hillside rivulets through brooks, streams, small and large rivers, to tidal estuaries. Underground streams carrying water with a free surface are also considered natural channels.

Notes

j) Channel Flow Routing of Watershed of Bridge No. 319 for Different Value of Overland Roughness (An)

Flow of watershed is made of overland and channel flow. Overland roughness is a important factor for channel flow. The effect of overland roughness on channel flow of watershed of bridge no. 319 is shown in the Fig. 4.5



Figure : Channel Flow Routing for Different Value of Overland Roughness (An)

From above Fig, we observed that if overland roughness is increasing then the peak flow is decreasing and the flow routing changes is shapes quickly. So the overland roughness on channel flow is sensitive.

k) Channel Flow Routing of Watershed of Bridge No. 319 for Different Value of Channel Roughness (An1)

Different channel has different channel roughness. It is a factor of channel flow. The effect of channel roughness on channel flow of watershed of bridge no.319 is shown in the following Fig.4.6.

© 2013 Global Journals Inc. (US)

Notes



Figure : Channel Flow Routing for Different Value of Channel Roughness (An1)

From the above figure we easily observed that the effect of channel roughness on channel flow peak is sensitive but is not like overland roughness. So channel roughness is very sensitive but not like as overland roughness on channel flow.

1) *Channel Flow Routing of Watershed of Bridge No.319 for Different Value of Side Slope (Bz)*

Channel flow depends on different physiographic parameters of overland and channel. Here the effect of side slope (BZ) on channel flow of watershed of bridge no. 319 is shown in Fig.



Figure : Channel Flow Routing for Different Value of Side Slope (Bz)

© 2013 Global Journals Inc. (US)

From the above figure we observed that if side slope is increasing then peak flow is decreasing very slowly. Therefore, the effect of side slope on the flow routing of channel flow is not important factor and hence side slope is not sensitive so much.

m) Channel Flow Routing of Watershed of Bridge No. 319 for Different Value of Overland Slope (So)

Channel flow is constituted of direct rainfall and overland flow. Channel flows depend on different physiographic parameters of overland and channel. Here the effect of overland slope (SO) on channel flow of watershed of bridge no. 319 is shown in the following Fig.



Figure : Channel Flow Routing for Different Value of Overland Slope

From the above figure, we observe that the effect of overland slope (SO) on the peak flow routing of channel flow if SO is increasing then flow routing is increasing, but after a period of flow, the flow is not increasing. So the overland slope is not so sensitive for channel flow.

II. Conclusion

From the above discussion we come in conclusion that the overland roughness and overland slope and channel roughness are more sensitive than the other parameters on overland and channel flow.

III. Scope of Furthur Study

The present study can be extended as follows:

(i) Distributed method can be used for channel flow and overland flow routing.

(ii) Three fully off-center first order explicit finite difference methods have been used for the solution of proposed KW models. A single step, second explicit method can also be used for above-mentioned models. Also, implicit finite difference methods and the finite element methods may also be used for this purpose.

(iii) In the proposed models, the contributing surface plane elements have assumed rectangular and perpendicular to the elements. The shape and the alignment of

Notes

© 2013 Global Journals Inc. (US)

these planes can be oriented according to the prevailing overland slopes. An altogether, different approach will have to be developed for such cases of modeling.

References Références Referencias

Notes

1. **Amien, M.**, and **Chu, H.L**. (1975). Implicit Numerical Modeling of Unsteady Flows. *J. of Hydr. Div., Proc. ASCE., Vol. 101(HY1):* 717-731.
2. **Amien, M.,** and **Fang, C.S**. (1970). Implicit Flood routing in Natural Channels. *J. of Hyd. Div., Proc. ASCE, Vol. 96 (HY12):* 2481-2500.
3. **Brakensick, D.L.** (1967a). A Simulated Watershed Flow System for Hydrograph Prediction: A Kinematics Application. Paper No.3, *Proc. of the Intl. Hydrology Symp, Fort Colino, Colardo.*
4. **Cambell, S.Y., Parlange, J.Y.** and **Rose, C.W**. (1984). Overland Flow on Converging and Diverging Surfaces-Kinematics model and Similarly Solutions. *J. Hydrology, Vol. (67): 367-374.*
5. **Carnahan, B., Luther, H.A.,** and **Wilkes, J.O.** (1969). *Applied Numerical Methods. John Willy, New York.*
6. **Cow, V.T**. (1959). Open channel Hydraulics. *Mc Graw-Hill Book Company. New York.*
7. **Clark, C.O**. (1945). Storage and the Unit Hydrograph. *Proc. ASCE, Vol. 69:* 1333-1360.
8. **Clarke, R.T** (1973). Mathematical Models in Hydrology. *Food and Agri. Organ. of the United Nations, Irrigation and Drainage paper No. 19, Rome.*
9. **Cooley, R.L.,** and **Moin, S.A.** (1976). Finite Element Solution of St. Venant Equations. *J. of Hyd. Div., Proc. ASCE, Vol. 102 (NY6): 759-775.*
10. **Dooge, J.C.I**. (1959). A General Theory of the Unit Hydrograph. *Journal of Geophysical Res., AUG, 64(2):241-256*
11. **De Vriesm J.J.,** and **MacArthur, R. C**. (1979). Introduction and Application of Kinematics Wave Routing Techniques Using HEC-1. Training Doc. No. 10, *HEC, DAVIS, California, USA.*
12. **Dooge, J.C.I.** (1969). A General Theory of the Unit Hydrograph. *Journal of Geophysical Res., AUG, 64(2): 241-256.*
13. **Dooge, J.C.I.** (1973). Linear Theory of Hydrologic Systems. *U.S. Dept. Agri., Agri. Res. Serv., Tech. Bull. 1468, 327 pp.*
14. **Freeze, R.A.** (1978). Mathematical Models of Hill slope Hydrology, in Hill slope Hydrology, M.J. Kirkby, eds., *John Wiley & Sons Ltd., New York.*
15. **Hossain, M. M.** (1989). Application of Kinematics Wave Theory to Small Watersheds Ph.D. Thesis. Submitted in the *Roorkee University, UP, India.*
16. **King, I.P.** (1977). Finite Element Methods for Unsteady Flow Through Irregular Channels. *Proc. First Conf. on Finite Elements in Water Resoures, Princeton Univ., U.S.A., July 1976;* Finite Elements in Water Resources, *Pentech Press,* pp: 165-184.
17. **Mathur, B. S.** (2972). Runoff Hydrographs for Uneven Spatial Distribution of Rainfall Ph.D. thesis, *The Indian Institute of Tech., New Delhi, India.*
18. **Moore. I.D. et al**. (1985). Kinematics Overland Flow: Generalization of Rose's approximate Solution. *J. of Hydrology, Vol. 82:* 233-245.
19. **Moore. I.D. et al**. (1985). Kinematics Overland Flow: Generalization of Rose's approximate Solution. Part II, *J, of Hydrol., Vol. 92:* 351-362.
20. **Nash, J.E**. (1957). The Form of Instantaneous Unit Hydrograph. *IASH General Assembly, Toronto, Pub. 45:* 114-119.

© 2013 Global Journals Inc. (US)

21. **Nash, J.E.** (1960). A unit Hydrograph Study with Particular Reference to British Catchments. *Proc. Inst, Civ. Engrs., No. 17*, pp: 249-281.
22. **Noye, J**. (1982). Numerical Solutions of Partial Differential Equations. *North-Holland Publishing Company, Amsterdam.*
23. **Overton, D.E.** and **Meadows, M.E.** (1976). Storm water Modeling. *Academic Press, New York.*

Notes

© 2013 Global Journals Inc. (US)

Global Journal of Science Frontier Research
Mathematics and Decision Sciences
Volume 13 Issue 7 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

Optimal Discounted Portfolio, Expected Wealth and Strategic Consumption for a Defined Contribution Pension Scheme

By C. I. Nkeki
University of Benin, Nigeria

Abstract - This paper deals with optimal discounted portfolio, expected wealth and strategic consumption process for a defined contribution (DC) pension scheme. The aims of this paper are to find: the optimal discounted portfolio and optimal discounted consumption choice to be adopted by the pension plan member (PPM) up to retirement period; the expected discounted wealth and variance of the expected wealth for the plan member. The financial market is composed by a riskless and a risky assets, and the effective salary of the plan member is assume to be stochastic. The expected discounted wealth and its variance are obtained. The discounted portfolio and consumption processes of the plan member are obtained. It is find that part of the discounted portfolio value is proportional to the ratio of the present value of the discounted expected future contributions to the optimal discounted wealth value overtime. It is also find that there is the need for gradual transfer of part of the portfolio value in risky asset to the riskless one against unforeseen shocks.

Keywords : Strategic discounted consumption, optimal discounted portfolio, pension plan member, unforeseen shocks, stochastic salary, defined contribution, pension scheme.

GJSFR-F Classification : MSC 2010: 91G10

OPTIMAL DISCOUNTED PORTFOLIO, EXPECTED WEALTH AND STRATEGIC CONSUMPTION FOR A DEFINED CONTRIBUTION PENSION SCHEME

Strictly as per the compliance and regulations of :

© 2013. C. I. Nkeki. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Global Journal of Science Frontier Research (F) Volume XIII Issue VII Version I Year 2013

Optimal Discounted Portfolio, Expected Wealth and Strategic Consumption for a Defined Contribution Pension Scheme

C. I. Nkeki

Abstract - This paper deals with optimal discounted portfolio, expected wealth and strategic consumption process for a defined contribution (DC) pension scheme. The aims of this paper are to find: the optimal discounted portfolio and optimal discounted consumption choice to be adopted by the pension plan member (PPM) up to retirement period; the expected discounted wealth and variance of the expected wealth for the plan member. The financial market is composed by a riskless and a risky assets, and the effective salary of the plan member is assume to be stochastic. The expected discounted wealth and its variance are obtained. The discounted portfolio and consumption processes of the plan member are obtained. It is find that part of the discounted portfolio value is proportional to the ratio of the present value of the discounted expected future contributions to the optimal discounted wealth value overtime. It is also find that there is the need for gradual transfer of part of the portfolio value in risky asset to the riskless one against unforeseen shocks.

Keywords : strategic discounted consumption, optimal discounted portfolio, pension plan member, unforeseen shocks, stochastic salary, defined contribution, pension scheme.

I. Introduction

We consider optimal discounted portfolio and discounted consumption problem in a defined contributory pension scheme. This paper follows the work of [14] and [16]. In their paper, they considered the optimal portfolio and strategic consumption in the lifecycle of a PPM in pension scheme. But, the area of discounted portfolio, discounted consumption and the variance of expected discounted wealth of a PPM are yet to be considered to the best of my knowledge in the literature of financial mathematics and actuarial science. In this paper, we assume that the PPM consume continuously throughout his or her life time and consumption terminates when the individual dies.

In the literature, [9] considered a numerical solution as well as analytical results to the intertemporal consumption problem for portfolio management. [17] examined a tractable model of precautionary savings in continuous time and assumed that the uncertainty was about the timing of the income loss in addition to the assumption of non-stochastic asset return. [3] considered labour supply flexibility and portfolio choice of individual life cycle. They considered the objective of maximizing the expected discounted lifetime utility by assumed that the utility function has two argument (consumption and labour/leiture). [2] adopted the quadratic utility function that characterized a linear marginal utility function. They asserted that the utility function was not attractive in describing the behaviour of individual towards risk as it implies increasing absolute risk

Ref

2. Blanchard, O. J. and Fischer, S. (1989), "Lectures on Macroeconomics". The MIT press.

Author : Department of Mathematics, Faculty of Physical Sciences, University of Benin, P. M. B. 1154, Benin City, Edo State, Nigeria.
E-mail : nkekicharles2003@yahoo.com

© 2013 Global Journals Inc. (US)

aversion. [3] concluded that labour income induced the individual to invest an additional amount of wealth to the risky asset. They established that labour income and investment choices are related, but they failed to analyzed the optimal consumption process of the investor. [4] studied the generally the optimal management of a defined contribution pension plan where the guarantee depends on the level of interest rates at a fixed retirement date. [5] considered optimal dynamic asset allocation strategy for a DC pension plan by taking into account the stochastic features of the plan member's lifetime salary progression and the stochastic properties of the assets held in accumulating pension fund. They emphasised that salary risk (the fluctuation in the plan member's earning in response to economic shocks) is not fully hedgeable using existing financial assets. They further emphasized that wage-indexed bonds could be used to hedge productivity and inflation shocks. They further asserted that such bonds are not widely traded. They referred the optimal dynamic asset allocation strategy stochastic lifestyling. They compared it against various static and deterministic lifestyle strategies in order to calculate the costs of adopting suboptimal strategies. Their solution technique made use of the present value of future contribution premiums into the plan. This technique can be found in [1], [4], [8]. Deterministic lifestyling which is the gradual switch from equities to bonds according to present rules is a popular asset alloction strategy during the accumulation phase of DC pension plans and is designed to protect the pension fund from a catastrophic fall in the stock market just prior to retirement (see [1], [5], [6]). [5] and [9] analysed extensively the occupational DC pension funds, where the contribution rate is a fixed percentage of salary.

The classical dynamic lifetime portfolio selection in a continuous time model was developed by [10], [11]. [9] used a dynamic programming approach to derived a formula for optimal investment allocation in a DC scheme and compared three risk measures to analyzed the terminal net replacement ratio achieved by members. They suggested that when the choice of investment strategy is determined, risk profiles of individual and different risk measures are both important factors which should be taken into consideration.

In this paper, we aim at finding the optimal discounted portfolio and optimal discounted consumption choice to be adopted by the pesnion plan member (PPM) up to retirement period and the entire lifetime; and the expected discounted wealth and variance of the expected discounted wealth for the plan member.

The structure of the remainder parts of the paper is as follows. Section 2 presents the formulation of the problem which include the financial models, wealth process and stochastic salary of a PPM. In section 3, we present the discounted wealth and discounted consumption process of a PPM. Section 4 presents the present value of discounted future contribution of a PPM. In section 5, we presents the valuation of the discounted wealth process of a PPM. Section 6 present the optimal discounted portfolio and consumption process of a PPM. In section 7, we present the optimal expected discounted wealth valuation and discounted consumption process of a PPM. Section 8 presents the accumulated expected discounted consumption process of a PPM. Finally, section 9 concludes the paper.

II. Problem Formulation

We consider a continuous-time financial market where there are two investment instruments: a riskless and a risky assets. The price dynamics of the two assets are given, respectively, by

Ref

3. Bodie, Z., Merton, R. C., and Samuelson, W. F., (1992) "Labour supply flexibility and portfolio choice in a life cycle model". *Journal of Economic Dynamics and Control*, 16, 427-449.

© 2013 Global Journals Inc. (US)

Notes

$$dB(t) = rB(t)dt,\ B(0) = 1, \tag{1}$$

$$dS(t) = S(t)\big(\mu dt + \sigma dW(t)\big),\ S(0) = s_0, \tag{2}$$

$t \in \Re$. We allow $r > 0, \mu > r$, to be constants. μ is the predictable 1-dimensional process of excess appreciation rate in relation to the stock, $\sigma = (\sigma_1, \sigma_2)$ is a predictable process of volatility vector of stock, σ_1 is volatility of stock arising from inflation and σ_2 is the volatility stock arising from the stock market. $\big(W(t) = (W_1(t), W_2(t))'; t \in \Re_+\big)$ is a standard 2- dimensional Brownian motion on a filtered probability space $\big(\Omega, \Im, \{\Im_t\}_{t\in[0,T]}, P\big)$ with $W(0)$ a null vector almost surely. $W_1(t)$ is the source of risk of inflation and $W_2(t)$ is the source of risk of the stock market. We assume that the filtration $\{\Im_t\}_{t\in[0,T]}$ is generated by the Brownian motion and is right continuous, and that each $\Im_t$ contains all the $P-$ null sets of $\Im$ We denote by $L^2_{\Im}$ the set of square integrable $\{\Im_t\}_{t\in[0,T]}-$ adapted processes,

$$L^2_{\Im} = \left\{ \begin{array}{l} \Delta \,|\, The\ \ process\ \ \Delta = \{\Delta(t)\}_{t\in[0,T]} \\ adapted\ \ \ process\ \ such\ \ that \\ \int_0^T E\left[\Delta^2(t)\right]dt < \infty \end{array} \right\}$$

and by $L^2_{\Im_T}$ the set of square integrable $\Im_T-$ measurable random variables,

$$L^2_{\Im_T} = \left\{ \Delta \left| \begin{array}{l} \Delta\ \ is\ \ an\ \ F_T - measurable\ \ random \\ \text{var}\,iable\ \ such\ \ that\ \ E\left[\Delta^2\right] < \infty \end{array} \right. \right\}$$

.From (1), we have

$$B(t) = \exp(rt),\ B(0) = 1; \tag{3}$$

We assume that the financial market is arbitrage-free, complete and continuously open between time 0 and T, i.e., there is only one process θ satisfying

$$\theta = \begin{pmatrix} \theta_1 \\ \theta_2 \end{pmatrix},$$

where, θ_1 is the market price of inflation risk and and

$$\theta_2 = \frac{\mu - r - \sigma_1\theta_1}{\sigma_2},$$

is the market price of stock risk.
The exponential process

$$Z(t) = \exp\left[-\theta' W(t) - \frac{1}{2}\|\theta\|^2 t\right], \tag{4}$$

is assumed to be a martingale
From (1) and (4), we have the discouted factor to be

© 2013 Global Journals Inc. (US)

$$\Lambda(t) = B(t)^{-1} Z(t) \tag{5}$$

Taking the differential (5), we have

$$d\Lambda(t) = -\Lambda(t)\big(rdt + \theta' dW(t)\big) \tag{6}$$

The dynamics of the PPM effective salary is given by

$$\begin{aligned} dY(t) &= Y(t)\big(\omega dt + \sigma^I dW(t)\big), \\ Y(0) &= y > 0, \end{aligned} \tag{7}$$

where $Y(t)$ is the salary of the PPM at time t, ω is the expected growth rate of salary of PPM and $\sigma^I = (\sigma_{Y1}, \sigma_{Y2})$ is the volatility vector of salary which is driven by the source of uncertainty of inflation, $W_1(t)$ and the stock market, $W_2(t)$. The two sources of risk are partial correlated. We assume that $\omega > 0$ is a constant and and σ^I is a constant vector.

III. Discounted Wealth and Discounted Consumption Process of a Ppm

In this section, we consider the discounted wealth process of the PPM in pension scheme.

Let $\Delta(t)$ portfolio process invested in risky asset at time t and $C(t)$ the consumption process at time t, then the pair (Δ, C) is said to be self-financing if the corresponding wealth process $X^{\Delta,C}(t), t \in [0,T]$, satisfies

$$dX^{\Delta,C}(t) = \Delta(t) X^{\Delta,C}(t) \frac{dS(t)}{S(t)} + \big(1 - \Delta(t)\big) X^{\Delta,C}(t) \frac{dB(t)}{B(t)} + \big(cY(t) - C(t)\big)dt, \tag{8}$$

where $\Delta_0 = 1 - \Delta(t)$ represents the proportion of the portfolio invested in cash account at time t.

Substituting the assets returns in (1) and (2) into (8), we obtain the following

$$dX^{\Delta,C}(t) = (X^{\Delta,C}(t)\big(r + \Delta(t)(\mu - r)\big) + cY(t) - C(t))dt + \sigma\Delta(t) X^{\Delta,C}(t) dW(t) \tag{9}$$

This is our stochastic differential equation which represents the wealth process of the PPM at time t.

Using (6) and (9), we have the discounted wealth process to be

$$d\big(\Lambda(t) X^{\Delta,C}(t)\big) = \big(c\Lambda(t)Y(t) - \Lambda(t)C(t)\big)dt + \Lambda(t) X^{\Delta,C}(t)(\Delta(t)\sigma' - \theta)' dW(t) \tag{10}$$

IV. Present Value of Discounted Future Contribution

In this section, we present the present value of pension plan member future contribution in pension scheme.

Definition 1: The present value of the expected discounted future contributions process is defined as

$$\widetilde{\Phi}(t) = E_t\left[\int_t^T \frac{\Lambda(s)}{\Lambda(t)} cY(s)ds\right], \tag{11}$$

Notes

© 2013 Global Journals Inc. (US)

where $c > 0$ is the proportion of the salary contributed into the pension funds and E_t is the conditional expectation with respect to the Brownian filtration $\{\Im(t)\}_{t\geq 0}$

$$\Lambda(t) \equiv Z(t)\exp[-rt], 0 \leq t \leq T. \tag{12}$$

is the stochastic discount factor which adjusts for nominal interest rate and market price of risk.

Notes

Definition 2: The expected discounted future consumption process is defined by

$$\Psi(t) = E\left[\int_t^{\infty} \frac{\Lambda(u)}{\Lambda(t)} C(u)du\right], t \in [T, \infty). \tag{13}$$

We take limits of integration in Eq. (13) from $t = T$ to $t = \infty$ because, we assume that consumption starts when the PPM retired and consume his/her investment (benefits) till he/she is dead.

Theorem 1: Let $\widetilde{\Phi}(t)$ be the present value of expected future contributions into the pension funds, then

$$\widetilde{\Phi}(t) = \frac{cY(t)}{\varphi}\left(\exp[\varphi(T-t)] - 1\right) \tag{14}$$

where $\varphi = \omega - r - \sigma^I \theta$

Proof: (see Nwozo and Nkeki (2011)).

Find the differential of bothsides of Eq. (14), we obtain

$$d\widetilde{\Phi}(t) = \widetilde{\Phi}(t)\left[\left(r + \sigma^I \theta\right)dt + \sigma^I dW(t)\right] - cY(t)dt \tag{15}$$

Using (6) and (15), we have

$$d\left(\Lambda(t)\widetilde{\Phi}(t)\right) = -c\Lambda(t)Y(t)dt + \Lambda(t)\widetilde{\Phi}(t)(\sigma^I{}' - \theta)' dW(t) \tag{16}$$

V. Valuation of the Discounted Wealth Process of a Ppm

Definition 3: The value of wealth of the PPM is define as

$$V(t) = X(t) + \widetilde{\Phi}(t). \tag{17}$$

So that the discounted value of wealth becomes

$$\Lambda(t)V(t) = \Lambda(t)X(t) + \Lambda(t)\widetilde{\Phi}(t).$$

Then, dynamics of the value of the discounted wealth of the PPM is obtain as

$$d\left(\Lambda(t)V(t)\right) = d\left(\Lambda(t)X(t)\right) + d\left(\Lambda(t)\widetilde{\Phi}(t)\right). \tag{18}$$

Substituting in (10) and (16) into (18), we obtain the change in wealth of the PPM as follows:

$$d\left(\Lambda(t)V(t)\right) = -\Lambda(t)C(t)dt + (\Lambda(t)X^{\Delta,C}(t)\left(\Delta(t)\sigma' - \theta\right) + \Lambda(t)\widetilde{\Phi}(t)\left(\sigma^I{}' - \theta\right))' dW(t) \tag{19}$$

© 2013 Global Journals Inc. (US)

VI. Optimal Portfolio and Consumption Process of a Ppm

In this section, we derived the optimal discounted portfolio and discounted consumption process of a PPM under dynamic programming principle. We assume that the PPM chooses power utility function in maximizing the expected utility of the terminal discounted wealth and consumption process. The choice of the power utility (which is a linear combination of two power utility functions. The first term is with respect to the wealth process while the second is with respect the consumption process) is motivated by the fact that pension scheme are in general large investment conpanies whose strategic plans are with respect to the size of funds they are managing.

We now give the optimal discounted portfolio and discounted consumption process for pension funds planners at time t.

The PPM's problem is to choose an admissible strategy so as to maximize the expected utility of accumulative consumptions and terminal discounted wealth,

$$U(t,V) = \sup_{\substack{\Delta \in \Pi(\ ,\Phi) \\ C \in \Re}} E_t^{x,\Phi} J(t,x,\Phi,\widetilde{C})$$

where

$$J(t,\ ,\Phi,\widetilde{C}) = \int_t^T e^{-\rho t} U(u,\widetilde{C}(u))du + U(V(T)),$$

We assume that U is concave and

$U((V,t) \in C^{1,2}(\Re \times [0,T])$, ρ is the PPM's preference discount factor, $\Pi(x,\Phi)$ set of admissible portfolio strategy that are F_V - progressively measureable, that satisfy the integrability condition

$$E\left[\int_0^T \Delta(s)^2 ds\right] < \infty,\ E\left[\int_0^T \widetilde{C}(s)^2 ds\right] < \infty,$$

$E_0^{x,\Phi}$ denotes the conditional expectation at time t given the initial endowment,

$$\Lambda(t)X^{\Delta,C}(t) = x,\ \Lambda(t)\widetilde{\Phi}(t) = \Phi,$$

$$\Lambda(t)C(t) = \widetilde{C}(t),$$

and the utility function is taken as

$$U(V) = \frac{V^\gamma}{\gamma}, 0 < \gamma < 1.$$

It turns out that $U(t,V)$ satisfies the following Hamilton–Jacobi–Bellman equation

$$\begin{aligned} U_t + \frac{1}{2}\ {}^2(\Delta(t)\sigma' - \theta)'(\Delta(t)\sigma' - \theta)U_{xx} + \frac{1}{2}\Phi^2(\sigma^I{}' - \theta)'(\sigma^I{}' - \theta)U_{\Phi\Phi} \\ \Phi(\sigma^I{}' - \theta)'(\Delta(t)\sigma' - \theta)U_{x\Phi} - \widetilde{C}(t)U_x + U(\widetilde{C}(t))\exp(-\rho t) = 0. \end{aligned} \tag{20}$$

Notes

© 2013 Global Journals Inc. (US)

Notes

This yields the HJB equation for the value function

$$\left(U_t V\right),t \ + \mathrm{H}(t,\Delta,C) = 0,$$

Subject to:

$$U(V) = \frac{(x+\Phi)^{\gamma}}{\gamma} - \frac{\widetilde{C}(t)^{\gamma}}{\gamma}, 0 < \gamma < 1. \tag{21}$$

where

$$\mathrm{H}(t,\Delta,\widetilde{C}) = \frac{1}{2}x^2\left(\Delta(t)\sigma' - \theta\right)'\left(\Delta(t)\sigma' - \theta\right)U_{xx}$$

$$+\frac{1}{2}\Phi^2\left(\sigma^I{}' - \theta\right)'\left(\sigma^I{}' - \theta\right)U_{\Phi\Phi}$$

$$x\Phi\left(\sigma^I{}' - \theta\right)'\left(\Delta(t)\sigma' - \theta\right)U_{x\Phi}$$

$$-\widetilde{C}(t)U_x + U(\widetilde{C}(t))\exp\left(-\rho t\right)$$

Hence, we have the following optimal discounted portfolio and optimal discounted consumption, respectively as follows:

$$\Delta^*(t) = \frac{\sigma^{-1}\theta}{x} - \frac{\Phi\sigma^{-1}\left(\sigma^I{}' - \theta\right)U_{x\Phi}}{xU_{xx}}, \tag{22}$$

$$\widetilde{C}*(t) = I\left(U_x \exp[\rho t]\right) \tag{29}$$

where, $I = \left(\frac{dU(\widetilde{C})}{d\widetilde{C}}\right)^{-1}$.

Substituting Eq. (22), and (29) into Eq.(20), we have

$$U_t + \frac{1}{2}\theta'\theta(1-x^2)U_{xx} -$$

$$\frac{1}{2}\Phi^2(\sigma^I{}' - \theta)'(\sigma^I{}' - \theta)\frac{U_{x\Phi}^2}{U_{xx}}$$

$$+\frac{1}{2}\Phi^2(\sigma^{I'} - \theta)'(\sigma^{I'} - \theta)U_{\Phi\Phi}$$

$$-I\left(U_x \exp(\rho t)\right)U_x +$$

$$\exp(-\rho t)U\left(I\left(U_x \exp(\rho t)\right)\right) = 0, \tag{36}$$

© 2013 Global Journals Inc. (US)

We assume that in this paper that

$$U(t,V)=\frac{(x+\Phi)^{\gamma}Q(t)^{\gamma}}{\gamma} \tag{23}$$

and

$$U(t,\widetilde{C})=\frac{\left(\widetilde{C}(t)Q(t)\right)^{\gamma}}{\gamma}.$$

Finding the partial derivative of (23) with respect to $t, x, xx, x\Phi, \Phi\Phi$, we have the following:

$$U_t=(x+\Phi)^{\gamma}Q(t)^{\gamma-1}Q'(t)$$

$$U_x=(x+\Phi)^{\gamma-1}Q(t)^{\gamma}$$

$$U_{\Phi}=(x+\Phi)^{\gamma-1}Q(t)^{\gamma}$$

$$U_{xx}=(\gamma-1)(x+\Phi)^{\gamma-2}Q(t)^{\gamma}$$

$$U_{x\Phi}=(\gamma-1)(x+\Phi)^{\gamma-2}Q(t)^{\gamma}$$

$$U_{\Phi\Phi}=(\gamma-1)(x+\Phi)^{\gamma-2}Q(t)^{\gamma}$$

We observe that the assumption of concavity of U turns out to be true, as

$$U_{xx}=U_{\Phi\Phi}=U_{x\Phi}=(\gamma-1)V^{\gamma-2}<0 \text{ since, } \gamma<1.$$

Substituting the partial derivatives into (36) we have

$$Q'(t)+\frac{1}{2V^2}\theta'\theta(1-x^2)(\gamma-1)Q(t)+\frac{1-\gamma}{\gamma V}\left[\left(Q(t)^{\frac{\gamma^2}{\gamma-1}}\exp\left[\frac{\rho t}{\gamma-1}\right]\right)\right]=0 \tag{24}$$

(24) can be solve numerically.

Suppose that

$$U(t,v_0)=\frac{(x_0+\Phi_0)^{\gamma}Q(t)^{\gamma}}{\gamma} \tag{25}$$

Then (25) becomes

$$Q'(t)+\frac{1}{2v_0^2}\theta'\theta(1-x_0^2)(\gamma-1)Q(t)+\frac{1-\gamma}{\gamma v_0}\left[\left(Q(t)^{\frac{\gamma^2}{\gamma-1}}\exp\left[\frac{\rho t}{\gamma-1}\right]\right)\right]=0 \tag{26}$$

Notes

© 2013 Global Journals Inc. (US)

Solving (26), we have

$$Q(t) = \widetilde{Q}(t)^{\frac{1-\gamma}{1-\gamma+\gamma^2}},$$

where,

Notes

$$\widetilde{Q}(t) = \frac{2v_0 e^{\frac{\rho t}{\gamma-1}}\big((\gamma-1)-2\gamma(\gamma-1)\big)}{\psi\gamma(\gamma-1)} + \frac{2v_0 e^{\frac{\rho t}{\gamma-1}}(\gamma^2(\gamma-1)+\gamma^2+2\gamma^3+\gamma^4)}{\psi\gamma(\gamma-1)} + \frac{2v_0 e^{\frac{\rho t}{\gamma-1}+\frac{r(T-t)}{\gamma-1}}\big((\gamma-1)(1+(\gamma-1)\gamma)\big)}{\psi\gamma} \times e^{\frac{1}{1-\gamma}\left(r-\rho+\frac{(x_0^2-1)(1+(\gamma-1)\gamma)(1-\gamma)\theta'\theta}{2v_0^2}\right)(T-t)}$$

and

$$\psi = (x_0^2-1)(\gamma-1)(1+(\gamma-1)\gamma)\theta'\theta + 2v_0^2\rho.$$

Therefore, the optimal discounted protfolio for the PPM at time t is obtained as

$$\Delta^*(t) = \frac{\sigma^{-1}\theta\big(\Lambda(t)X^*(t)+\Lambda(t)\Phi(t)\big)}{\Lambda(t)X^*(t)} - \frac{\sigma^{-1}\sigma^{I}{}'\Phi(t)}{X^*(t)},$$

$$\Delta_0 = 1 + \frac{\sigma^{-1}\sigma^{I}{}'\Phi(t)}{X^*(t)} - \frac{\sigma^{-1}\theta\big(\Lambda(t)X^*(t)+\Lambda(t)\Phi(t)\big)}{\Lambda(t)X^*(t)}, \tag{27}$$

the optimal discounted consumption process $C^*(t)$ of the PPM at time t, given that (25) holds, is obtained

$$\widetilde{C}^*(t) = (x_0+\Phi_0)(g(t)+h(t)\times e^{\frac{1}{1-\gamma}\left(r-\rho+\frac{(x_0^2-1)(1+(\gamma-1)\gamma)(1-\gamma)\theta'\theta}{2v_0^2}\right)(T-t)}) \tag{28}$$

where

$$g(t) = \frac{2v_0 e^{\frac{\rho t}{\gamma-1}}\big((\gamma-1)-2\gamma(\gamma-1)\big)}{\psi\gamma(\gamma-1)} + \frac{2v_0 e^{\frac{\rho t}{\gamma-1}}(\gamma^2(\gamma-1)+\gamma^2+2\gamma^3+\gamma^4)}{\psi\gamma(\gamma-1)}$$

© 2013 Global Journals Inc. (US)

and

$$h(t) = \frac{2v_0 e^{\frac{\rho t}{\gamma-1}+\frac{r(T-t)}{\gamma-1}}\left((\gamma-1)(1+(\gamma-1)\gamma)\right)}{\psi\gamma}.$$

At time $t = 0$, (28) becomes

$$\widetilde{C}^*(0) = (x_0 + \Phi_0)(g(0) + h(0) \times e^{\frac{1}{1-\gamma}\left(r-\rho+\frac{(\,{}^2_0-1)(1+(\gamma-1)\gamma)(1-\gamma)\theta'\theta}{2v_0^2}\right)T}) \tag{29}$$

where $$g(0) = \frac{2v_0\left((\gamma-1)-2\gamma(\gamma-1)\right)}{\psi\gamma(\gamma-1)} + \frac{2v_0(\gamma^2(\gamma-1)+\gamma^2+2\gamma^3+\gamma^4)}{\psi\gamma(\gamma-1)},$$

and $$h(0) = \frac{2v_0 e^{\frac{rT}{\gamma-1}}\left((\gamma-1)(1+(\gamma-1)\gamma)\right)}{\psi\gamma}$$

At time $t = T$, (28) becomes

$$\widetilde{C}^*(T) = (x_0 + \Phi_0)(g(T) + h(T)), \tag{29}$$

where

$$g(T) = \frac{2v_0 e^{\frac{\rho T}{\gamma-1}}\left((\gamma-1)-2\gamma(\gamma-1)\right)}{\psi\gamma(\gamma-1)} + \frac{2v_0 e^{\frac{\rho T}{\gamma-1}}(\gamma^2(\gamma-1)+\gamma^2+2\gamma^3+\gamma^4)}{\psi\gamma(\gamma-1)}$$

and $$h(T) = \frac{2v_0 e^{\frac{\rho T}{\gamma-1}}\left((\gamma-1)(1+(\gamma-1)\gamma)\right)}{\psi\gamma}.$$

Now, for $0 < \gamma < 1$, we have intuitively that the growth rate (GRC) of the optimal expected discounted consumption is obtain as

$$GRC = \frac{1}{1-\gamma}\left(r-\rho+\frac{(x_0^2-1)(1+(\gamma-1)\gamma)(1-\gamma)\theta'\theta}{2v_0^2}\right). \tag{30}$$

This is referred to as the Euler equation for the intertemporal maximization of discounted consumption under uncertainty. The positive term $\theta'\theta$ captures the uncer-

Notes

© 2013 Global Journals Inc. (US)

tainty of the financial martet. When the financial market is risky, it will induce investors to shift consumption over time. Suppose there is no initial wealth, then (30) becomes

$$GRC = \frac{1}{1-\gamma}(r - \rho + \frac{(1+(\gamma-1)\gamma)(\gamma-1)\theta'\theta}{2\Phi_0^2}).$$

Notes

From (27), the first term is the variational form of the classical portfolio strategy while the last term is an intertemporal hedging strategy that offset any shock to the stochastic salary of the PPM. At $t = 0$, we have

$$\Delta^*(0) = \frac{\sigma^{-1}\theta(x_0 + \Phi_0)}{x_0} - \frac{\sigma^{-1}\sigma^{I}{}'\Phi_0}{x_0},$$

$$\Delta_0 = 1 + \frac{\sigma^{-1}\sigma^{I}{}'\Phi_0}{x_0} - \frac{\sigma^{-1}\theta(x_0 + \Phi_0)}{x_0}.$$

VII. Optimal Expected Discounted Wealth Valuation and Discounted Consumption Process of a Ppm

In this section, we present the optimal expected value of wealth of the PPM in pension plan at time t. From (19), we have that

$$d(\Lambda(t)V(t)) = -\Lambda(t)C(t)dt + \theta(1 - \Lambda(t)V(t) + \Lambda(t)\Phi(t))dW(t) \tag{31}$$

$$\begin{aligned} d(\Lambda(t)V(t))^2 &= (-2\Lambda(t)V(t)\Lambda(t)C(t)dt + \\ &\theta'\theta(\Lambda(t)V(t))^2 - \theta'\theta\Lambda(t)V(t) - \\ &\theta'\theta\Lambda(t)V(t)\Lambda(t)\Phi(t) + \\ &\theta'\theta(1 + (\Lambda(t)\Phi(t))^2 + (\Lambda(t)\Phi(t))dt \\ &+ 2\theta'\Lambda(t)V(t)(1 - \Lambda(t)V(t) + \Lambda(t)\Phi(t))dW(t) \end{aligned} \tag{32}$$

$$dE(\Lambda(t)V(t)) = -E(\Lambda(t)C(t))dt \tag{33}$$

$$\begin{aligned} dE(\Lambda(t)V(t))^2 &= E(-2\Lambda(t)V(t)\Lambda(t)C(t) + \\ &\theta'\theta(\Lambda(t)V(t))^2 - \theta'\theta\Lambda(t)V(t) - \\ &\theta'\theta\Lambda(t)V(t)\Lambda(t)\Phi(t) + \\ &\theta'\theta(1 + (\Lambda(t)\Phi(t))^2 + (\Lambda(t)\Phi(t))dt \end{aligned} \tag{34}$$

Solving, (33) and (34), we have

$$E(\Lambda(t)V(t)) = v_0 - E\left(\int_0^t (\Lambda(s)C(s))ds\right) \tag{35}$$

© 2013 Global Journals Inc. (US)

$$E(\Lambda(t)V(t))^2 = v_0^2 e^{\theta'\theta t} + \int_0^t \mathrm{K}(s) e^{\theta'\theta(t-s)} ds \tag{36}$$

where,

$$\begin{aligned}
&\mathrm{K}(t) = 2E\left(\Lambda(t)C(t)(v_0 - \textstyle\int_0^t \Lambda(s)C(s)ds)\right) + \\
&\theta'\theta E\left(v_0 - \textstyle\int_0^t \Lambda(s)C(s)ds\right) + \\
&\frac{c\theta'\theta e^{\omega t}(e^{\varphi(T-t)}-1)}{\varphi} E\left(v_0 - \textstyle\int_0^t \Lambda(s)C(s)ds\right) + \\
&\theta'\theta\left(1 + \frac{c^2 e^{2\omega t}(e^{\varphi(T-t)}-1)^2}{\varphi^2} + \frac{c e^{\omega t}(e^{\varphi(T-t)}-1)}{\varphi}\right).
\end{aligned}$$

$$\begin{aligned}
&Var(\Lambda(t)V(t)) = E(\Lambda(t)V(t))^2 - (E(\Lambda(t)V(t)))^2 \\
&= v_0^2 e^{\theta'\theta t} + \int_0^t \mathrm{K}(s) e^{\theta'\theta(t-s)} ds - \\
&\left(v_0 - E\left(\int_0^t (\Lambda(s)C(s))ds\right)\right)^2
\end{aligned} \tag{37}$$

At $t = T$, we have

$$E(\Lambda(T)V(T)) = v_0 - E\left(\int_0^T (\Lambda(s)C(s))ds\right) \tag{38}$$

$$\begin{aligned}
&Var(\Lambda(T)V(T)) = E(\Lambda(T)V(T))^2 - (E(\Lambda(T)V(T)))^2 = \\
&v_0^2 e^{\theta'\theta T} + \int_0^T \mathrm{K}(s) e^{\theta'\theta(T-s)} ds \\
&-\left(v_0 - E\left(\int_0^T (\Lambda(s)C(s))ds\right)\right)^2.
\end{aligned}$$

VIII. Accumulated Expected Discounted Consumption Process of a PPM

The accumulated expected discounted consumption process of the PPM up to retirement period is given by

$$\begin{aligned}
\widetilde{\Psi}(T) &= E\left[\int_0^T \widetilde{C}(u)du\right] = \int_0^T (x_0 + \Phi_0)(g(t) + h(t)e^{GRC(T-t)})dt \\
&= (x_0 + \Phi_0)\int_0^T g(t)dt + (x_0 + \Phi_0)\int_0^T h(t)e^{GRC(T-t)})dt
\end{aligned} \tag{39}$$

Notes

© 2013 Global Journals Inc. (US)

Notes

Solving (39), we have

$$\widetilde{\Psi}(T)=\frac{2v^2{}_0\big((\gamma-1)-2\gamma(\gamma-1)\big)}{\psi\gamma\rho}(1-e^{\frac{\rho T}{\gamma-1}})$$

$$+\frac{2v^2{}_0(\gamma^2(\gamma-1)+\gamma^2+2\gamma^3+\gamma^4)}{\psi\gamma\rho}(1-e^{\frac{\rho T}{\gamma-1}})$$

$$+\frac{2v^2{}_0\big((1-\gamma)(1+(\gamma-1)\gamma)\big)e^{\frac{(\rho+\alpha)T}{\gamma-1}}}{\alpha\psi\gamma}(1-e^{\frac{\alpha T}{1-\gamma}})$$

where $$\alpha=\frac{(x_0^2-1)(1+(\gamma-1)\gamma)(1-\gamma)\theta'\theta}{2v_0^2}.$$

If we allow T to tend to infinity, we have

$$\lim_{T\to\infty}E_0\left[\int_0^T\widetilde{C}(u)du\right]=E_0\left[\int_0^\infty\widetilde{C}(u)du\right]=$$

$$\lim_{T\to\infty}\widetilde{\Psi}(T)=\widetilde{\Psi}_\infty=\frac{2v^2{}_0\big((\gamma-1)-2\gamma(\gamma-1)\big)}{\psi\gamma\rho}$$

$$+\frac{2v^2{}_0(\gamma^2(\gamma-1)+\gamma^2+2\gamma^3+\gamma^4)}{\psi\gamma\rho}.$$

This shows that throughout the life-cycle of the PPM, his or her accumulated expected discounted consumption will remain nonempty. If $\widetilde{\Psi}_\infty$ is nonnegative, it implies that the PPM is unable to finished his or her consumption before he or she dies. If $\widetilde{\Psi}_\infty$ is negative, it implies that the PPM is finished his or her consumption why still alive. If $\widetilde{\Psi}_\infty$ is zero, it implies that the period the PPM finishes his or her consumption was the period he or she dies. Interestingly, the nature of $\widetilde{\Psi}_\infty$ absolutely depend on the value of γ,ρ. This implies that the level of risk the PPM takes will depend on how large or how small the value of the accumulated discounted consumption will be and consumption rate. We observe that at $\rho=0$ or

$$\rho=\frac{(x_0^2-1)(\gamma-1)(1+(\gamma-1)\gamma)\theta'\theta}{2v_0^2},$$

optimal expected discounted consumption will be unbounded. The optimal expected discounted consumption will be negative if

$$\rho<\frac{(x_0^2-1)(\gamma-1)(1+(\gamma-1)\gamma)\theta'\theta}{2v_0^2},$$

© 2013 Global Journals Inc. (US)

and non negative if

$$\rho > \frac{(x_0^2 - 1)(\gamma - 1)(1 + (\gamma - 1)\gamma)\theta'\theta}{2v_0^2}.$$

IX. Conclusion

This paper dealt with optimal discounted portfolio, expected wealth and strategic life-time consumption process for a defined contributory pension scheme. The expected discounted wealth and its variance are obtained. The discounted variational classical portfolio and consumption process of the plan member are established. The accumulated discounted consumption process of the PPM throughout his or her life-cycle was established. It was found that part of the discounted portfolio value is proportional to the ratio of the present value of the discounted expected future contributions to the optimal discounted portfolio value overtime. It was found that there is the need for gradual transfer of part of the portfolio value in risky asset to the riskless one against unforeseen shocks.

Notes

References Références Referencias

1. Blake, D., Wright, D. and Zhang, Y., (2008), "Optimal funding and investment strategies in defined contribution pension plans under Epstein-Zin utility". Discussion paper, The pensions Institute, Cass Business School, City University, UK.
2. Blanchard, O. J. and Fischer, S. (1989), "Lectures on Macroeconomics". The MIT press.
3. Bodie, Z., Merton, R. C., and Samuelson, W. F., (1992) "Labour supply flexibility and portfolio choice in a life cycle model". *Journal of Economic Dynamics and Control*, 16, 427-449.
4. Boulier, J. F, Huang, S. J. and Taillard, G., (2001), "Optimal management under stochasticinterest rates: The case of a protected defined contribution pension fund". *Insurance: Mathematics and Economics*, Vol. 28, pp. 173-189.
5. Cairns, A. J. G., Blake, D. and Dowd, K., (*2003),* "Stochastic lifestyling: Optimal dynamic asset allocation for defined contribution pension plans". *CRIS Discussion paper series, The University of 2ottingham.*
6. Cajueiro, D. O. and Yoneyama, T., (*2003)*, "Optimal portfolio, Optimal consumption and the Markowitz mean-variance analysis in a switching diffusion market". Working paper.
7. Fleming, W. and Rishel, R., (1975), "Deterministic and stochastic optimal control". Springer-Verlag, New York.
8. Korn, R. and Krekel, M., (2001), "Optimal portfolios with fixed consumption or income streams". *Working paper, University of Kaiserslautern.*
9. Mason, R. and Wright, S., (2001), "The effects of uncertainty on optimal consumption", *Journal of Economic Dynamics and Control,* 25, 185-212.

© 2013 Global Journals Inc. (US)

Notes

10. Merton, R. C., "Lifetime portfolio selection under uncertainty: the continuous-time case". *Review of Economics and Statistics,* 1969, Vol. 51, pp. 247-257.
11. Merton, R. C., (1971), "Optimum consumption and portfolio rules in a continuous-time model". *Journal of Economic Theory*, Vol. 3, pp. 373-413.
12. Nkeki, C. I. (2011), On optimal portfolio management of the accumulation phase of a defined contributory pension scheme, Ph.D thesis, Department of Mathematics, University of Ibadan, Ibadan, Nigeria.
13. Nkeki C. I.and Nwozo C. R, (2012), Variational form of classical portfolio strategy and expected wealth for a defined contributory pension scheme, Journal of Mathematical Finance, 2(1), pp.132-139.
14. Nwozo C. R. and Nkeki, C. I. (2011), Optimal portfolio and strategic consumption planning in a life-cycle of a pension plan member in a defined contributory pension scheme, IAENG International Journal of Applied Mathematics, 41(4), pp.299-309.
15. Toche, P., "A tractable model of precautionary saving in continuous time". *Economics letters,* 2005, 87, 267-272.

© 2013 Global Journals Inc. (US)

Notes

This page is intentionally left blank

© 2013 Global Journals Inc. (US)

Global Journal of Science Frontier Research
Mathematics and Decision Sciences
Volume 13 Issue 7 Version 1.0 Year 2013
Type : Double Blind Peer Reviewed International Research Journal
Publisher: Global Journals Inc. (USA)
Online ISSN: 2249-4626 & Print ISSN: 0975-5896

On the Right Perfect Ring

By R.H. Sallam
Helwan University, Egypt

Abstract - We prove that a right self right perfect algebra which is at most countable dimensional modulo their Jacobson radical is right artinian.

Keywords : *self injective rings, right perfect rings, quasi-frobenius rings.*

GJSFR-F Classification : *MSC 2010: 16D50*

ON THE RIGHT PERFECT RING

Strictly as per the compliance and regulations of :

© 2013. R.H. Sallam. This is a research/review paper, distributed under the terms of the Creative Commons Attribution-Noncommercial 3.0 Unported License http://creativecommons.org/licenses/by-nc/3.0/), permitting all non commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

On the Right Perfect Ring

R.H. Sallam

Abstract - We prove that a right self right perfect algebra which is at most countable dimensional modulo their Jacobson radical is right artinian.

Keywords : self injective rings, right perfect rings, quasi-frobenius rings.

Section 1

I. Introduction

In this note we present a proof for algebra R over a field K which is at most countable imensional modulo their Jacobson radical i.e. R/Rad(R) is at most N_o

This includes for example the important situation when R/Rad R is not only semisimple but also finite dimensional' Let R be a ring with identity and RadR is its Jacobson radical. If R is right perfect (i.e. R/RadR is semisimple and RadR is left T-nilpotent) then R/RadR is semisimple and hence $(R/RadR)_R$ and $_R(R/RadR)$ are semisimple right and left R-module.

Note that since R/RadR is semisimple then an R-module M is semisimple iff it is cancelled by RadR. Indeed every simple is cancelled by Rad R and if RadM=0 then M has R/RadR –module structure which is semisimple, therefore M is semisimple as the lattice of R-submodules and R/RadR sub-modules coincide in this case.

$(R/RadR)_R$ is semisimple and finitely generated, so it has a composition series i.e.

$$J/J^2 \subset J^2/J^3 \subset \ldots \subset J^{n-1}/J^n$$

is of length n-1 where each J^k/J^{k+1} is semisimple for all k

Note that if R is right self injective then each decomposable (eR) has simple socle. Indeed if we have a nontrivial decomposition of the socle $socle(eR) = M \oplus N$

Then we can find E(M) and E(N)injective hulls of M, N contained in eR and we obtain eR= E(M) + E(N) nontrivial decomposition that is a contradiction.

Note also that if R is right self injective right perfect then for each simple Right R-module, the left module Hom (S, R) is simple.

First Hom (S, R) is non zero. Looking for the isomorphism types of indecomposable modules eR, these are projective, local and the cover of some simple R-module.

The number of isomorphism types of such modules equals the number of isomorphism types of simple modules equal t, say.

Moreover since the indecomposable eR's also injective with simple socle we see that they are isomorphic if and only if their socle isomorphic. This shows that the distinct

Author : Mathematics Department, Faculty of Science, Helwan University, Cairo, Egypt. E-mail : rsallams@hotmail.com

© 2013 Global Journals Inc. (US)

types of isomorphism of simples occurring as socle of some eR is also t, and so each simple S must appears as a socle of some l eR (i.e. it embeds in R) this shows that

$Hom(S,R) \neq 0$, for each simple R-modules S. If f, $g \in Hom(S,R)$ and $f \neq 0$, *then* $f : S \to R$

Is a monomorphism and since R_R is injective then there is some $h : R \to R$

Such that $h \circ f = g$. If $f(x) = xc \ \forall x \in R$ we get $f(x).c = g(x)$ i.e. $f.c = g$ in $Hom(S, R)$, this shows that $Hom(S, R)$ is generated by $f \neq 0$ so it is simple.

In particular since each simple module embeds in R which is right self injective, it follows that R is injective cogenerator in the category of right R-module, i.e. it is a right PF (Pseudo- Frobonius) ring.

Section 2

Let S be a set of repesenatives for simple right R-modules and let t= S and $R/RadR = \oplus S^{K_s}$. Let W= socle (R_R) be the right socle of R so W is an R-sub-bimodule of R.

Since each indecomposable module (Re) has simple socle, we have Length (W) equals the number of terms in the indecomposable decomposition $R = \bigoplus_S$ Re which equals length $(R/RadR)_R$, since each Indecomposable (Re) is local

Let $W = \oplus S^{P_s}$ we have $\sum_{s \in S} P_s = \sum_{s \in S} K_s$

Proposition 1

Let R be a right self injective right perfect ring then the set $\{Hom(S, R); S \in S\}$ is a set of representatives for simple left R-modules.

In particular $Hom(S, R)$ and $Hom(T, R)$ are nonisomorphic for nonisomorphic, $S, T \in R$

Proof

Since R is right self injective the monomorphism

$0 \to W \to R$, gives rise to the epimorphism

$$Hom(R, R) \to Hom(W.R) \to 0$$

Note that

$$Hom(W, R) = \oplus Hom(S, R)$$

Since $Hom(S, R) \neq 0 \ \forall \ S \in S$, we have

$$Hom(W, R) = \oplus Hom(S, R)^{P_s},$$

has length equal to the length $W = \sum_{s \in S} P^s = length(R/RadR)_R$

By classical Wedderburn –Artin Theorem:

$$length(R/RadR)_R = length_R(R/RadR)$$

Since $Hom(W,R)$ is semisimple, the kernel of $R \to Hom(W,R)$

Contains $RadR$ and therefore since

$$length_R(R/RadR) = length \ Hom_R(W, R),$$

© 2013 Global Journals Inc. (US)

we obtain $R/RadR \cong Hom(W, R)$ as left R-module.

This shows that all types of isomorphism of simple left R-modules are found among components of $Hom(W,R)$ and the statement is proved

Note: we note that the above proof further shows that there is an exact sequence of left R-modules

Notes

$$0 \to RadR \to R \to Hom(W, R) \to 0$$

This means that

$$\{ r \in R, \text{ suchthat}, rW = 0 \} = RadR. \text{ That is } ann({}_R W) = RadR$$

This shows that W is also semisimple as a left R-module, i.e. the right socle of R is contained in the left socle, and hence the left and the right socle or the right PF ring coincide.

For a right R-module M, denote $M^* = Hom(M,R)$ this is a left R-module.

Proposition 2

Let R be a right injective ring and let M be a right R-module such that there is an exact sequence

$$0 \to S_R \to M_R \to L_R^{(\alpha)} \to 0$$

With S, L simple modules and assume S= socle (M), $L_R^{(\alpha)}$ denote the coproduct of α copies of L_R. Then M^* is local left R-module with unique maximal ideal

$$S^{\perp} = \{ f \in Hom(M, R); f|_S \neq 0 \}$$

Which is semisimple isomorphic to $(L_R^*)^{(\alpha)}$

Proof

Since

$$0 \to S_R \to M_R \to L_R^{(\alpha)} \to 0$$

is an exact sequence, so we get the exact sequence,

$$0 \to Hom(L^{(\alpha)}, R) \to Hom(M, R) \to Hom(S, R) \to 0$$

That is

$$0 \to (L^*)^{\alpha} \to M^* \to S^* \to 0$$

The kernel of the morphism

$$M^* = Hom(M, R) \xrightarrow{\gamma^*} S^* = Hom(S, R)$$

$$\text{Is} \quad \{ f \in Hom(M, R); f|_S \neq 0 \} = S^{\perp}$$

Hence $S^{\perp} \cong (L^*)^{\alpha}$ which is left semisimple module since it is cancelled by $RadR$.

© 2013 Global Journals Inc. (US)

Now since M has simple socle, and its socle embeds in R which is injective it follows that M embeds in R. We note that M^* is generated by any $f \notin S^{\perp}$ which will show that M is cyclic. Indeed such f must be a monomorphism and given any other $h : M \to R$.

By injectivity of R_R there is $g \in Hom(R,R)$ such that $g \circ f = h$. If $g(x) = cx$ for $c \in R$ then we have $h = c.f$ in M^*.This shows that $f..R = M$ which shows that $S^{\perp}$ is the only maximal submodules of the cyclic left Rmodule M^*, that ends of the proof.

Note that the fact that M^* is local can also be proved by embedding M in some indecomposable (eR) for an indecomposable idempotent e, and then, by applying the exact functor $Hom(-,R)$ one obtain an epimorphism $Hom(eR, R) = eR \to M$, and so M^* is local because eR is.

Let α be the largest cardinality for which there is a right R-module M with simple socle and such that $M/(socle(M)) \cong L^{\alpha}$ for some simple module L.

Such cardinality exists, since any such module is contained in R because R is injective. In fact if W_1 is the second socle of R, then $\alpha \leq length(W_2 / W)$ we note that if α is infinite this is an equality.

Indeed if for each simple modules S, L we denote by $\alpha_{S,L} = [(E(S)/S) : L]$ – the multiplicity of L in the second socle of the injective hull E(S) of S, then $\alpha = \max_{S,L}(\alpha_{S,L})$.

Therefore $\alpha \leq \sum_{S,L \in S} \alpha_{S,L} \leq n\alpha = \alpha$ if α is infinite.

We note also that if $\sum_k$ is the k'th socle then the $length(W_k / W_{k-1}) \leq \alpha$; this follows by induction on k.
If this is true for k, then there is an embedding

$$W_k / W_{k-1} \hookrightarrow R^{(\alpha)}$$

And therefore we have

$$length\ (W_{k+1} / W_k) \leq length\ (W_1 / W_0)^{(\alpha)} = \alpha \times \alpha = \alpha ,$$

since α is infinite cardinal. So we have the following theorem:

Theorem

Let R be a right self injective right perfect algebra such that the dimension of each simple R-module is at most countable (equivalently the dimension of $R / RadR$ is at most countable) hence R is countable, and hence right artinian.

Proof

With the above notation, assume α is infinite. The length of each W_k / W_{k-1} is at most α, so since the dimension of each simple is at most $\aleph_0$, its dimension is at most $\aleph_0$, × α = α (since α is infinite).

Thus the dimension of R is at most α, and so it equals α (since $length(W_1 / W_0) = \alpha$).

On the other hand, by proposition (2.2) there is a local left R-module M^*, with socle L^{α} for some simple L.

Notes

© 2013 Global Journals Inc. (US)

Note that $\dim(L^{\alpha}) \geq 2^{\alpha}$ and that there is an epimorphism $R \to M^{*}$
So $\dim(R) \geq 2^{\alpha}$, this is a contradiction.

By the result of Lawrence [3] R is right self injective countably generated algebra over a field so it is quasi -frobenius and hence right artinian.

References Références Referencias

Notes

1. F. W. Anderson and K. R. Fuller, Rings and Categories of Modules Springer-Verlag, Heidelberg, New York, (1991).
2. F. Kasch, Modules and Rings, London Mathematical Society Monographs" Vol.17, Academic Press, New York, (1982).
3. J. Lawrence, a countable self injective ring is quasi-frobenius, proceeding of the American mathematical society, volume 65, No.2, august 1977.
4. M. C. Iovanov, semiprimary selfinjective algebras with at most countable dimensional Jacobson quotient are quasi frobenius, arxiv: 1111.2901, 2011.
5. T-Y. Lam, lectures on modules and rings, Graduate Texts in Mathematics, No.189. Berlin-New York, Springer-Verlag (1999).

© 2013 Global Journals Inc. (US)

Global Journals Inc. (US) Guidelines Handbook 2013

www.GlobalJournals.org

FELLOW OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (FARSS)

- 'FARSS' title will be awarded to the person after approval of Editor-in-Chief and Editorial Board. The title 'FARSS" can be added to name in the following manner. eg. Dr. John E. Hall, Ph.D., FARSS or William Walldroff Ph. D., M.S., FARSS
- Being FARSS is a respectful honor. It authenticates your research activities. After becoming FARSS, you can use 'FARSS' title as you use your degree in suffix of your name. This will definitely will enhance and add up your name. You can use it on your Career Counseling Materials/CV/Resume/Visiting Card/Name Plate etc.
- 60% Discount will be provided to FARSS members for publishing research papers in Global Journals Inc., if our Editorial Board and Peer Reviewers accept the paper. For the life time, if you are author/co-author of any paper bill sent to you will automatically be discounted one by 60%
- FARSS will be given a renowned, secure, free professional email address with 100 GB of space eg.johnhall@globaljournals.org. You will be facilitated with Webmail, SpamAssassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.
- FARSS member is eligible to become paid peer reviewer at Global Journals Inc. to earn up to 15% of realized author charges taken from author of respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account or to your PayPal account.
- Eg. If we had taken 420 USD from author, we can send 63 USD to your account.
- FARSS member can apply for free approval, grading and certification of some of their Educational and Institutional Degrees from Global Journals Inc. (US) and Open Association of Research,Society U.S.A.
- After you are FARSS. You can send us scanned copy of all of your documents. We will verify, grade and certify them within a month. It will be based on your academic records, quality of research papers published by you, and 50 more criteria. This is beneficial for your job interviews as recruiting organization need not just rely on you for authenticity and your unknown qualities, you would have authentic ranks of all of your documents. Our scale is unique worldwide.
- FARSS member can proceed to get benefits of free research podcasting in Global Research Radio with their research documents, slides and online movies.
- After your publication anywhere in the world, you can upload you research paper with your recorded voice or you can use our professional RJs to record your paper their voice. We can also stream your conference videos and display your slides online.
- FARSS will be eligible for free application of Standardization of their Researches by Open Scientific Standards. Standardization is next step and level after publishing in a journal. A team of research and professional will work with you to take your research to its next level, which is worldwide open standardization.

- FARSS is eligible to earn from their researches: While publishing his paper with Global Journals Inc. (US), FARSS can decide whether he/she would like to publish his/her research in closed manner. When readers will buy that individual research paper for reading, 80% of its earning by Global Journals Inc. (US) will be transferred to FARSS member's bank account after certain threshold balance. There is no time limit for collection. FARSS member can decide its price and we can help in decision.

MEMBER OF ASSOCIATION OF RESEARCH SOCIETY IN SCIENCE (MARSS)

- 'MARSS' title will be awarded to the person after approval of Editor-in-Chief and Editorial Board. The title 'MARSS" can be added to name in the following manner. eg. Dr. John E. Hall, Ph.D., MARSS or William Walldroff Ph. D., M.S., MARSS
- Being MARSS is a respectful honor. It authenticates your research activities. After becoming MARSS, you can use 'MARSS' title as you use your degree in suffix of your name. This will definitely will enhance and add up your name. You can use it on your Career Counseling Materials/CV/Resume/Visiting Card/Name Plate etc.
- 40% Discount will be provided to MARSS members for publishing research papers in Global Journals Inc., if our Editorial Board and Peer Reviewers accept the paper. For the life time, if you are author/co-author of any paper bill sent to you will automatically be discounted one by 60%
- MARSS will be given a renowned, secure, free professional email address with 30 GB of space eg.johnhall@globaljournals.org. You will be facilitated with Webmail, SpamAssassin, Email Forwarders, Auto-Responders, Email Delivery Route tracing, etc.
- MARSS member is eligible to become paid peer reviewer at Global Journals Inc. to earn up to 10% of realized author charges taken from author of respective paper. After reviewing 5 or more papers you can request to transfer the amount to your bank account or to your PayPal account.
- MARSS member can apply for free approval, grading and certification of some of their Educational and Institutional Degrees from Global Journals Inc. (US) and Open Association of Research,Society U.S.A.
- MARSS is eligible to earn from their researches: While publishing his paper with Global Journals Inc. (US), MARSS can decide whether he/she would like to publish his/her research in closed manner. When readers will buy that individual research paper for reading, 40% of its earning by Global Journals Inc. (US) will be transferred to MARSS member's bank account after certain threshold balance. There is no time limit for collection. MARSS member can decide its price and we can help in decision.

Auxiliary Memberships

ANNUAL MEMBER

- Annual Member will be authorized to receive e-Journal GJSFR for one year (subscription for one year).
- The member will be allotted free 1 GB Web-space along with subDomain to contribute and participate in our activities.
- A professional email address will be allotted free 500 MB email space.

PAPER PUBLICATION

- The members can publish paper once. The paper will be sent to two-peer reviewer. The paper will be published after the acceptance of peer reviewers and Editorial Board.

Process of submission of Research Paper

The Area or field of specialization may or may not be of any category as mentioned in 'Scope of Journal' menu of the GlobalJournals.org website. There are 37 Research Journal categorized with Six parental Journals GJCST, GJMR, GJRE, GJMBR, GJSFR, GJHSS. For Authors should prefer the mentioned categories. There are three widely used systems UDC, DDC and LCC. The details are available as 'Knowledge Abstract' at Home page. The major advantage of this coding is that, the research work will be exposed to and shared with all over the world as we are being abstracted and indexed worldwide.

The paper should be in proper format. The format can be downloaded from first page of 'Author Guideline' Menu. The Author is expected to follow the general rules as mentioned in this menu. The paper should be written in MS-Word Format (*.DOC,*.DOCX).

The Author can submit the paper either online or offline. The authors should prefer online submission.Online Submission: There are three ways to submit your paper:

(A) (I) First, register yourself using top right corner of Home page then Login. If you are already registered, then login using your username and password.

(II) Choose corresponding Journal.

(III) Click 'Submit Manuscript'. Fill required information and Upload the paper.

(B) If you are using Internet Explorer, then Direct Submission through Homepage is also available.

(C) If these two are not conveninet , and then email the paper directly to dean@globaljournals.org.

Offline Submission: Author can send the typed form of paper by Post. However, online submission should be preferred.

Preferred Author Guidelines

MANUSCRIPT STYLE INSTRUCTION (Must be strictly followed)

Page Size: 8.27" X 11'"

- Left Margin: 0.65
- Right Margin: 0.65
- Top Margin: 0.75
- Bottom Margin: 0.75
- Font type of all text should be Swis 721 Lt BT.
- Paper Title should be of Font Size 24 with one Column section.
- Author Name in Font Size of 11 with one column as of Title.
- Abstract Font size of 9 Bold, "Abstract" word in Italic Bold.
- Main Text: Font size 10 with justified two columns section
- Two Column with Equal Column with of 3.38 and Gaping of .2
- First Character must be three lines Drop capped.
- Paragraph before Spacing of 1 pt and After of 0 pt.
- Line Spacing of 1 pt
- Large Images must be in One Column
- Numbering of First Main Headings (Heading 1) must be in Roman Letters, Capital Letter, and Font Size of 10.
- Numbering of Second Main Headings (Heading 2) must be in Alphabets, Italic, and Font Size of 10.

You can use your own standard format also.
Author Guidelines:

1. General,

2. Ethical Guidelines,

3. Submission of Manuscripts,

4. Manuscript's Category,

5. Structure and Format of Manuscript,

6. After Acceptance.

1. GENERAL

Before submitting your research paper, one is advised to go through the details as mentioned in following heads. It will be beneficial, while peer reviewer justify your paper for publication.

Scope

The Global Journals Inc. (US) welcome the submission of original paper, review paper, survey article relevant to the all the streams of Philosophy and knowledge. The Global Journals Inc. (US) is parental platform for Global Journal of Computer Science and Technology, Researches in Engineering, Medical Research, Science Frontier Research, Human Social Science, Management, and Business organization. The choice of specific field can be done otherwise as following in Abstracting and Indexing Page on this Website. As the all Global

Journals Inc. (US) are being abstracted and indexed (in process) by most of the reputed organizations. Topics of only narrow interest will not be accepted unless they have wider potential or consequences.

2. ETHICAL GUIDELINES

Authors should follow the ethical guidelines as mentioned below for publication of research paper and research activities.

Papers are accepted on strict understanding that the material in whole or in part has not been, nor is being, considered for publication elsewhere. If the paper once accepted by Global Journals Inc. (US) and Editorial Board, will become the copyright of the Global Journals Inc. (US).

Authorship: The authors and coauthors should have active contribution to conception design, analysis and interpretation of findings. They should critically review the contents and drafting of the paper. All should approve the final version of the paper before submission

The Global Journals Inc. (US) follows the definition of authorship set up by the Global Academy of Research and Development. According to the Global Academy of R&D authorship, criteria must be based on:

1) Substantial contributions to conception and acquisition of data, analysis and interpretation of the findings.

2) Drafting the paper and revising it critically regarding important academic content.

3) Final approval of the version of the paper to be published.

All authors should have been credited according to their appropriate contribution in research activity and preparing paper. Contributors who do not match the criteria as authors may be mentioned under Acknowledgement.

Acknowledgements: Contributors to the research other than authors credited should be mentioned under acknowledgement. The specifications of the source of funding for the research if appropriate can be included. Suppliers of resources may be mentioned along with address.

Appeal of Decision: The Editorial Board's decision on publication of the paper is final and cannot be appealed elsewhere.

Permissions: It is the author's responsibility to have prior permission if all or parts of earlier published illustrations are used in this paper.

Please mention proper reference and appropriate acknowledgements wherever expected.

If all or parts of previously published illustrations are used, permission must be taken from the copyright holder concerned. It is the author's responsibility to take these in writing.

Approval for reproduction/modification of any information (including figures and tables) published elsewhere must be obtained by the authors/copyright holders before submission of the manuscript. Contributors (Authors) are responsible for any copyright fee involved.

3. SUBMISSION OF MANUSCRIPTS

Manuscripts should be uploaded via this online submission page. The online submission is most efficient method for submission of papers, as it enables rapid distribution of manuscripts and consequently speeds up the review procedure. It also enables authors to know the status of their own manuscripts by emailing us. Complete instructions for submitting a paper is available below.

Manuscript submission is a systematic procedure and little preparation is required beyond having all parts of your manuscript in a given format and a computer with an Internet connection and a Web browser. Full help and instructions are provided on-screen. As an author, you will be prompted for login and manuscript details as Field of Paper and then to upload your manuscript file(s) according to the instructions.

To avoid postal delays, all transaction is preferred by e-mail. A finished manuscript submission is confirmed by e-mail immediately and your paper enters the editorial process with no postal delays. When a conclusion is made about the publication of your paper by our Editorial Board, revisions can be submitted online with the same procedure, with an occasion to view and respond to all comments.

Complete support for both authors and co-author is provided.

4. MANUSCRIPT'S CATEGORY

Based on potential and nature, the manuscript can be categorized under the following heads:

Original research paper: Such papers are reports of high-level significant original research work.

Review papers: These are concise, significant but helpful and decisive topics for young researchers.

Research articles: These are handled with small investigation and applications

Research letters: The letters are small and concise comments on previously published matters.

5.STRUCTURE AND FORMAT OF MANUSCRIPT

The recommended size of original research paper is less than seven thousand words, review papers fewer than seven thousands words also.Preparation of research paper or how to write research paper, are major hurdle, while writing manuscript. The research articles and research letters should be fewer than three thousand words, the structure original research paper; sometime review paper should be as follows:

Papers: These are reports of significant research (typically less than 7000 words equivalent, including tables, figures, references), and comprise:

(a)Title should be relevant and commensurate with the theme of the paper.

(b) A brief Summary, "Abstract" (less than 150 words) containing the major results and conclusions.

(c) Up to ten keywords, that precisely identifies the paper's subject, purpose, and focus.

(d) An Introduction, giving necessary background excluding subheadings; objectives must be clearly declared.

(e) Resources and techniques with sufficient complete experimental details (wherever possible by reference) to permit repetition; sources of information must be given and numerical methods must be specified by reference, unless non-standard.

(f) Results should be presented concisely, by well-designed tables and/or figures; the same data may not be used in both; suitable statistical data should be given. All data must be obtained with attention to numerical detail in the planning stage. As reproduced design has been recognized to be important to experiments for a considerable time, the Editor has decided that any paper that appears not to have adequate numerical treatments of the data will be returned un-refereed;

(g) Discussion should cover the implications and consequences, not just recapitulating the results; conclusions should be summarizing.

(h) Brief Acknowledgements.

(i) References in the proper form.

Authors should very cautiously consider the preparation of papers to ensure that they communicate efficiently. Papers are much more likely to be accepted, if they are cautiously designed and laid out, contain few or no errors, are summarizing, and be conventional to the approach and instructions. They will in addition, be published with much less delays than those that require much technical and editorial correction.

The Editorial Board reserves the right to make literary corrections and to make suggestions to improve briefness.

It is vital, that authors take care in submitting a manuscript that is written in simple language and adheres to published guidelines.

Format

Language: The language of publication is UK English. Authors, for whom English is a second language, must have their manuscript efficiently edited by an English-speaking person before submission to make sure that, the English is of high excellence. It is preferable, that manuscripts should be professionally edited.

Standard Usage, Abbreviations, and Units: Spelling and hyphenation should be conventional to The Concise Oxford English Dictionary. Statistics and measurements should at all times be given in figures, e.g. 16 min, except for when the number begins a sentence. When the number does not refer to a unit of measurement it should be spelt in full unless, it is 160 or greater.

Abbreviations supposed to be used carefully. The abbreviated name or expression is supposed to be cited in full at first usage, followed by the conventional abbreviation in parentheses.

Metric SI units are supposed to generally be used excluding where they conflict with current practice or are confusing. For illustration, 1.4 l rather than 1.4 × 10-3 m3, or 4 mm somewhat than 4 × 10-3 m. Chemical formula and solutions must identify the form used, e.g. anhydrous or hydrated, and the concentration must be in clearly defined units. Common species names should be followed by underlines at the first mention. For following use the generic name should be constricted to a single letter, if it is clear.

Structure

All manuscripts submitted to Global Journals Inc. (US), ought to include:

Title: The title page must carry an instructive title that reflects the content, a running title (less than 45 characters together with spaces), names of the authors and co-authors, and the place(s) wherever the work was carried out. The full postal address in addition with the e-mail address of related author must be given. Up to eleven keywords or very brief phrases have to be given to help data retrieval, mining and indexing.

Abstract, used in Original Papers and Reviews:

Optimizing Abstract for Search Engines

Many researchers searching for information online will use search engines such as Google, Yahoo or similar. By optimizing your paper for search engines, you will amplify the chance of someone finding it. This in turn will make it more likely to be viewed and/or cited in a further work. Global Journals Inc. (US) have compiled these guidelines to facilitate you to maximize the web-friendliness of the most public part of your paper.

Key Words

A major linchpin in research work for the writing research paper is the keyword search, which one will employ to find both library and Internet resources.

One must be persistent and creative in using keywords. An effective keyword search requires a strategy and planning a list of possible keywords and phrases to try.

Search engines for most searches, use Boolean searching, which is somewhat different from Internet searches. The Boolean search uses "operators," words (and, or, not, and near) that enable you to expand or narrow your affords. Tips for research paper while preparing research paper are very helpful guideline of research paper.

Choice of key words is first tool of tips to write research paper. Research paper writing is an art.A few tips for deciding as strategically as possible about keyword search:

- One should start brainstorming lists of possible keywords before even begin searching. Think about the most important concepts related to research work. Ask, "What words would a source have to include to be truly valuable in research paper?" Then consider synonyms for the important words.
- It may take the discovery of only one relevant paper to let steer in the right keyword direction because in most databases, the keywords under which a research paper is abstracted are listed with the paper.
- One should avoid outdated words.

Keywords are the key that opens a door to research work sources. Keyword searching is an art in which researcher's skills are bound to improve with experience and time.

Numerical Methods: Numerical methods used should be clear and, where appropriate, supported by references.

Acknowledgements: Please make these as concise as possible.

References

References follow the Harvard scheme of referencing. References in the text should cite the authors' names followed by the time of their publication, unless there are three or more authors when simply the first author's name is quoted followed by et al. unpublished work has to only be cited where necessary, and only in the text. Copies of references in press in other journals have to be supplied with submitted typescripts. It is necessary that all citations and references be carefully checked before submission, as mistakes or omissions will cause delays.

References to information on the World Wide Web can be given, but only if the information is available without charge to readers on an official site. Wikipedia and Similar websites are not allowed where anyone can change the information. Authors will be asked to make available electronic copies of the cited information for inclusion on the Global Journals Inc. (US) homepage at the judgment of the Editorial Board.

The Editorial Board and Global Journals Inc. (US) recommend that, citation of online-published papers and other material should be done via a DOI (digital object identifier). If an author cites anything, which does not have a DOI, they run the risk of the cited material not being noticeable.

The Editorial Board and Global Journals Inc. (US) recommend the use of a tool such as Reference Manager for reference management and formatting.

Tables, Figures and Figure Legends

Tables: Tables should be few in number, cautiously designed, uncrowned, and include only essential data. Each must have an Arabic number, e.g. Table 4, a self-explanatory caption and be on a separate sheet. Vertical lines should not be used.

Figures: Figures are supposed to be submitted as separate files. Always take in a citation in the text for each figure using Arabic numbers, e.g. Fig. 4. Artwork must be submitted online in electronic form by e-mailing them.

Preparation of Electronic Figures for Publication

Even though low quality images are sufficient for review purposes, print publication requires high quality images to prevent the final product being blurred or fuzzy. Submit (or e-mail) EPS (line art) or TIFF (halftone/photographs) files only. MS PowerPoint and Word Graphics are unsuitable for printed pictures. Do not use pixel-oriented software. Scans (TIFF only) should have a resolution of at least 350 dpi (halftone) or 700 to 1100 dpi (line drawings) in relation to the imitation size. Please give the data for figures in black and white or submit a Color Work Agreement Form. EPS files must be saved with fonts embedded (and with a TIFF preview, if possible).

For scanned images, the scanning resolution (at final image size) ought to be as follows to ensure good reproduction: line art: >650 dpi; halftones (including gel photographs) : >350 dpi; figures containing both halftone and line images: >650 dpi.

Color Charges: It is the rule of the Global Journals Inc. (US) for authors to pay the full cost for the reproduction of their color artwork. Hence, please note that, if there is color artwork in your manuscript when it is accepted for publication, we would require you to complete and return a color work agreement form before your paper can be published.

Figure Legends: Self-explanatory legends of all figures should be incorporated separately under the heading 'Legends to Figures'. In the full-text online edition of the journal, figure legends may possibly be truncated in abbreviated links to the full screen version. Therefore, the first 100 characters of any legend should notify the reader, about the key aspects of the figure.

6. AFTER ACCEPTANCE

Upon approval of a paper for publication, the manuscript will be forwarded to the dean, who is responsible for the publication of the Global Journals Inc. (US).

6.1 Proof Corrections

The corresponding author will receive an e-mail alert containing a link to a website or will be attached. A working e-mail address must therefore be provided for the related author.

Acrobat Reader will be required in order to read this file. This software can be downloaded

(Free of charge) from the following website:

www.adobe.com/products/acrobat/readstep2.html. This will facilitate the file to be opened, read on screen, and printed out in order for any corrections to be added. Further instructions will be sent with the proof.

Proofs must be returned to the dean at dean@globaljournals.org within three days of receipt.

As changes to proofs are costly, we inquire that you only correct typesetting errors. All illustrations are retained by the publisher. Please note that the authors are responsible for all statements made in their work, including changes made by the copy editor.

6.2 Early View of Global Journals Inc. (US) (Publication Prior to Print)

The Global Journals Inc. (US) are enclosed by our publishing's Early View service. Early View articles are complete full-text articles sent in advance of their publication. Early View articles are absolute and final. They have been completely reviewed, revised and edited for publication, and the authors' final corrections have been incorporated. Because they are in final form, no changes can be made after sending them. The nature of Early View articles means that they do not yet have volume, issue or page numbers, so Early View articles cannot be cited in the conventional way.

6.3 Author Services

Online production tracking is available for your article through Author Services. Author Services enables authors to track their article - once it has been accepted - through the production process to publication online and in print. Authors can check the status of their articles online and choose to receive automated e-mails at key stages of production. The authors will receive an e-mail with a unique link that enables them to register and have their article automatically added to the system. Please ensure that a complete e-mail address is provided when submitting the manuscript.

6.4 Author Material Archive Policy

Please note that if not specifically requested, publisher will dispose off hardcopy & electronic information submitted, after the two months of publication. If you require the return of any information submitted, please inform the Editorial Board or dean as soon as possible.

6.5 Offprint and Extra Copies

A PDF offprint of the online-published article will be provided free of charge to the related author, and may be distributed according to the Publisher's terms and conditions. Additional paper offprint may be ordered by emailing us at: editor@globaljournals.org .

Before start writing a good quality Computer Science Research Paper, let us first understand what is Computer Science Research Paper? So, Computer Science Research Paper is the paper which is written by professionals or scientists who are associated to Computer Science and Information Technology, or doing research study in these areas. If you are novel to this field then you can consult about this field from your supervisor or guide.

TECHNIQUES FOR WRITING A GOOD QUALITY RESEARCH PAPER:

1. Choosing the topic: In most cases, the topic is searched by the interest of author but it can be also suggested by the guides. You can have several topics and then you can judge that in which topic or subject you are finding yourself most comfortable. This can be done by asking several questions to yourself, like Will I be able to carry our search in this area? Will I find all necessary recourses to accomplish the search? Will I be able to find all information in this field area? If the answer of these types of questions will be "Yes" then you can choose that topic. In most of the cases, you may have to conduct the surveys and have to visit several places because this field is related to Computer Science and Information Technology. Also, you may have to do a lot of work to find all rise and falls regarding the various data of that subject. Sometimes, detailed information plays a vital role, instead of short information.

2. Evaluators are human: First thing to remember that evaluators are also human being. They are not only meant for rejecting a paper. They are here to evaluate your paper. So, present your Best.

3. Think Like Evaluators: If you are in a confusion or getting demotivated that your paper will be accepted by evaluators or not, then think and try to evaluate your paper like an Evaluator. Try to understand that what an evaluator wants in your research paper and automatically you will have your answer.

4. Make blueprints of paper: The outline is the plan or framework that will help you to arrange your thoughts. It will make your paper logical. But remember that all points of your outline must be related to the topic you have chosen.

5. Ask your Guides: If you are having any difficulty in your research, then do not hesitate to share your difficulty to your guide (if you have any). They will surely help you out and resolve your doubts. If you can't clarify what exactly you require for your work then ask the supervisor to help you with the alternative. He might also provide you the list of essential readings.

6. Use of computer is recommended: As you are doing research in the field of Computer Science, then this point is quite obvious.

7. Use right software: Always use good quality software packages. If you are not capable to judge good software then you can lose quality of your paper unknowingly. There are various software programs available to help you, which you can get through Internet.

8. Use the Internet for help: An excellent start for your paper can be by using the Google. It is an excellent search engine, where you can have your doubts resolved. You may also read some answers for the frequent question how to write my research paper or find model research paper. From the internet library you can download books. If you have all required books make important reading selecting and analyzing the specified information. Then put together research paper sketch out.

9. Use and get big pictures: Always use encyclopedias, Wikipedia to get pictures so that you can go into the depth.

10. Bookmarks are useful: When you read any book or magazine, you generally use bookmarks, right! It is a good habit, which helps to not to lose your continuity. You should always use bookmarks while searching on Internet also, which will make your search easier.

11. Revise what you wrote: When you write anything, always read it, summarize it and then finalize it.

12. Make all efforts: Make all efforts to mention what you are going to write in your paper. That means always have a good start. Try to mention everything in introduction, that what is the need of a particular research paper. Polish your work by good skill of writing and always give an evaluator, what he wants.

13. Have backups: When you are going to do any important thing like making research paper, you should always have backup copies of it either in your computer or in paper. This will help you to not to lose any of your important.

14. Produce good diagrams of your own: Always try to include good charts or diagrams in your paper to improve quality. Using several and unnecessary diagrams will degrade the quality of your paper by creating "hotchpotch." So always, try to make and include those diagrams, which are made by your own to improve readability and understandability of your paper.

15. Use of direct quotes: When you do research relevant to literature, history or current affairs then use of quotes become essential but if study is relevant to science then use of quotes is not preferable.

16. Use proper verb tense: Use proper verb tenses in your paper. Use past tense, to present those events that happened. Use present tense to indicate events that are going on. Use future tense to indicate future happening events. Use of improper and wrong tenses will confuse the evaluator. Avoid the sentences that are incomplete.

17. Never use online paper: If you are getting any paper on Internet, then never use it as your research paper because it might be possible that evaluator has already seen it or maybe it is outdated version.

18. Pick a good study spot: To do your research studies always try to pick a spot, which is quiet. Every spot is not for studies. Spot that suits you choose it and proceed further.

19. Know what you know: Always try to know, what you know by making objectives. Else, you will be confused and cannot achieve your target.

20. Use good quality grammar: Always use a good quality grammar and use words that will throw positive impact on evaluator. Use of good quality grammar does not mean to use tough words, that for each word the evaluator has to go through dictionary. Do not start sentence with a conjunction. Do not fragment sentences. Eliminate one-word sentences. Ignore passive voice. Do not ever use a big word when a diminutive one would suffice. Verbs have to be in agreement with their subjects. Prepositions are not expressions to finish sentences with. It is incorrect to ever divide an infinitive. Avoid clichés like the disease. Also, always shun irritating alliteration. Use language that is simple and straight forward. put together a neat summary.

21. Arrangement of information: Each section of the main body should start with an opening sentence and there should be a changeover at the end of the section. Give only valid and powerful arguments to your topic. You may also maintain your arguments with records.

22. Never start in last minute: Always start at right time and give enough time to research work. Leaving everything to the last minute will degrade your paper and spoil your work.

23. Multitasking in research is not good: Doing several things at the same time proves bad habit in case of research activity. Research is an area, where everything has a particular time slot. Divide your research work in parts and do particular part in particular time slot.

24. Never copy others' work: Never copy others' work and give it your name because if evaluator has seen it anywhere you will be in trouble.

25. Take proper rest and food: No matter how many hours you spend for your research activity, if you are not taking care of your health then all your efforts will be in vain. For a quality research, study is must, and this can be done by taking proper rest and food.

26. Go for seminars: Attend seminars if the topic is relevant to your research area. Utilize all your resources.

27. Refresh your mind after intervals: Try to give rest to your mind by listening to soft music or by sleeping in intervals. This will also improve your memory.

28. Make colleagues: Always try to make colleagues. No matter how sharper or intelligent you are, if you make colleagues you can have several ideas, which will be helpful for your research.

29. Think technically: Always think technically. If anything happens, then search its reasons, its benefits, and demerits.

30. Think and then print: When you will go to print your paper, notice that tables are not be split, headings are not detached from their descriptions, and page sequence is maintained.

31. Adding unnecessary information: Do not add unnecessary information, like, I have used MS Excel to draw graph. Do not add irrelevant and inappropriate material. These all will create superfluous. Foreign terminology and phrases are not apropos. One should NEVER take a broad view. Analogy in script is like feathers on a snake. Not at all use a large word when a very small one would be sufficient. Use words properly, regardless of how others use them. Remove quotations. Puns are for kids, not grunt readers. Amplification is a billion times of inferior quality than sarcasm.

32. Never oversimplify everything: To add material in your research paper, never go for oversimplification. This will definitely irritate the evaluator. Be more or less specific. Also too, by no means, ever use rhythmic redundancies. Contractions aren't essential and shouldn't be there used. Comparisons are as terrible as clichés. Give up ampersands and abbreviations, and so on. Remove commas, that are, not necessary. Parenthetical words however should be together with this in commas. Understatement is all the time the complete best way to put onward earth-shaking thoughts. Give a detailed literary review.

33. Report concluded results: Use concluded results. From raw data, filter the results and then conclude your studies based on measurements and observations taken. Significant figures and appropriate number of decimal places should be used. Parenthetical remarks are prohibitive. Proofread carefully at final stage. In the end give outline to your arguments. Spot out perspectives of further study of this subject. Justify your conclusion by at the bottom of them with sufficient justifications and examples.

34. After conclusion: Once you have concluded your research, the next most important step is to present your findings. Presentation is extremely important as it is the definite medium though which your research is going to be in print to the rest of the crowd. Care should be taken to categorize your thoughts well and present them in a logical and neat manner. A good quality research paper format is essential because it serves to highlight your research paper and bring to light all necessary aspects in your research.

Informal Guidelines of Research Paper Writing

Key points to remember:

- Submit all work in its final form.
- Write your paper in the form, which is presented in the guidelines using the template.
- Please note the criterion for grading the final paper by peer-reviewers.

Final Points:

A purpose of organizing a research paper is to let people to interpret your effort selectively. The journal requires the following sections, submitted in the order listed, each section to start on a new page.

The introduction will be compiled from reference matter and will reflect the design processes or outline of basis that direct you to make study. As you will carry out the process of study, the method and process section will be constructed as like that. The result segment will show related statistics in nearly sequential order and will direct the reviewers next to the similar intellectual paths throughout the data that you took to carry out your study. The discussion section will provide understanding of the data and projections as to the implication of the results. The use of good quality references all through the paper will give the effort trustworthiness by representing an alertness of prior workings.

Writing a research paper is not an easy job no matter how trouble-free the actual research or concept. Practice, excellent preparation, and controlled record keeping are the only means to make straightforward the progression.

General style:

Specific editorial column necessities for compliance of a manuscript will always take over from directions in these general guidelines.

To make a paper clear

· Adhere to recommended page limits

Mistakes to evade

- Insertion a title at the foot of a page with the subsequent text on the next page
- Separating a table/chart or figure - impound each figure/table to a single page
- Submitting a manuscript with pages out of sequence

In every sections of your document

· Use standard writing style including articles ("a", "the," etc.)

· Keep on paying attention on the research topic of the paper

· Use paragraphs to split each significant point (excluding for the abstract)

· Align the primary line of each section

· Present your points in sound order

· Use present tense to report well accepted

· Use past tense to describe specific results

· Shun familiar wording, don't address the reviewer directly, and don't use slang, slang language, or superlatives

· Shun use of extra pictures - include only those figures essential to presenting results

Title Page:

Choose a revealing title. It should be short. It should not have non-standard acronyms or abbreviations. It should not exceed two printed lines. It should include the name(s) and address (es) of all authors.

Abstract:

The summary should be two hundred words or less. It should briefly and clearly explain the key findings reported in the manuscript--must have precise statistics. It should not have abnormal acronyms or abbreviations. It should be logical in itself. Shun citing references at this point.

An abstract is a brief distinct paragraph summary of finished work or work in development. In a minute or less a reviewer can be taught the foundation behind the study, common approach to the problem, relevant results, and significant conclusions or new questions.

Write your summary when your paper is completed because how can you write the summary of anything which is not yet written? Wealth of terminology is very essential in abstract. Yet, use comprehensive sentences and do not let go readability for briefness. You can maintain it succinct by phrasing sentences so that they provide more than lone rationale. The author can at this moment go straight to shortening the outcome. Sum up the study, with the subsequent elements in any summary. Try to maintain the initial two items to no more than one ruling each.

- Reason of the study - theory, overall issue, purpose
- Fundamental goal
- To the point depiction of the research
- Consequences, including definite statistics - if the consequences are quantitative in nature, account quantitative data; results of any numerical analysis should be reported
- Significant conclusions or questions that track from the research(es)

Approach:

- Single section, and succinct
- As a outline of job done, it is always written in past tense
- A conceptual should situate on its own, and not submit to any other part of the paper such as a form or table
- Center on shortening results - bound background information to a verdict or two, if completely necessary
- What you account in an conceptual must be regular with what you reported in the manuscript
- Exact spelling, clearness of sentences and phrases, and appropriate reporting of quantities (proper units, important statistics) are just as significant in an abstract as they are anywhere else

Introduction:

The **Introduction** should "introduce" the manuscript. The reviewer should be presented with sufficient background information to be capable to comprehend and calculate the purpose of your study without having to submit to other works. The basis for the study should be offered. Give most important references but shun difficult to make a comprehensive appraisal of the topic. In the introduction, describe the problem visibly. If the problem is not acknowledged in a logical, reasonable way, the reviewer will have no attention in your result. Speak in common terms about techniques used to explain the problem, if needed, but do not present any particulars about the protocols here. Following approach can create a valuable beginning:

- Explain the value (significance) of the study
- Shield the model - why did you employ this particular system or method? What is its compensation? You strength remark on its appropriateness from a abstract point of vision as well as point out sensible reasons for using it.
- Present a justification. Status your particular theory (es) or aim(s), and describe the logic that led you to choose them.
- Very for a short time explain the tentative propose and how it skilled the declared objectives.

Approach:

- Use past tense except for when referring to recognized facts. After all, the manuscript will be submitted after the entire job is done.
- Sort out your thoughts; manufacture one key point with every section. If you make the four points listed above, you will need a least of four paragraphs.

- Present surroundings information only as desirable in order hold up a situation. The reviewer does not desire to read the whole thing you know about a topic.
- Shape the theory/purpose specifically - do not take a broad view.
- As always, give awareness to spelling, simplicity and correctness of sentences and phrases.

Procedures (Methods and Materials):

This part is supposed to be the easiest to carve if you have good skills. A sound written Procedures segment allows a capable scientist to replacement your results. Present precise information about your supplies. The suppliers and clarity of reagents can be helpful bits of information. Present methods in sequential order but linked methodologies can be grouped as a segment. Be concise when relating the protocols. Attempt for the least amount of information that would permit another capable scientist to spare your outcome but be cautious that vital information is integrated. The use of subheadings is suggested and ought to be synchronized with the results section. When a technique is used that has been well described in another object, mention the specific item describing a way but draw the basic principle while stating the situation. The purpose is to text all particular resources and broad procedures, so that another person may use some or all of the methods in one more study or referee the scientific value of your work. It is not to be a step by step report of the whole thing you did, nor is a methods section a set of orders.

Materials:

- Explain materials individually only if the study is so complex that it saves liberty this way.
- Embrace particular materials, and any tools or provisions that are not frequently found in laboratories.
- Do not take in frequently found.
- If use of a definite type of tools.
- Materials may be reported in a part section or else they may be recognized along with your measures.

Methods:

- Report the method (not particulars of each process that engaged the same methodology)
- Describe the method entirely
- To be succinct, present methods under headings dedicated to specific dealings or groups of measures
- Simplify - details how procedures were completed not how they were exclusively performed on a particular day.
- If well known procedures were used, account the procedure by name, possibly with reference, and that's all.

Approach:

- It is embarrassed or not possible to use vigorous voice when documenting methods with no using first person, which would focus the reviewer's interest on the researcher rather than the job. As a result when script up the methods most authors use third person passive voice.
- Use standard style in this and in every other part of the paper - avoid familiar lists, and use full sentences.

What to keep away from

- Resources and methods are not a set of information.
- Skip all descriptive information and surroundings - save it for the argument.
- Leave out information that is immaterial to a third party.

Results:

The principle of a results segment is to present and demonstrate your conclusion. Create this part a entirely objective details of the outcome, and save all understanding for the discussion.

The page length of this segment is set by the sum and types of data to be reported. Carry on to be to the point, by means of statistics and tables, if suitable, to present consequences most efficiently.You must obviously differentiate material that would usually be incorporated in a study editorial from any unprocessed data or additional appendix matter that would not be available. In fact, such matter should not be submitted at all except requested by the instructor.

Content

- Sum up your conclusion in text and demonstrate them, if suitable, with figures and tables.
- In manuscript, explain each of your consequences, point the reader to remarks that are most appropriate.
- Present a background, such as by describing the question that was addressed by creation an exacting study.
- Explain results of control experiments and comprise remarks that are not accessible in a prescribed figure or table, if appropriate.
- Examine your data, then prepare the analyzed (transformed) data in the form of a figure (graph), table, or in manuscript form.

What to stay away from

- Do not discuss or infer your outcome, report surroundings information, or try to explain anything.
- Not at all, take in raw data or intermediate calculations in a research manuscript.
- Do not present the similar data more than once.
- Manuscript should complement any figures or tables, not duplicate the identical information.
- Never confuse figures with tables - there is a difference.

Approach

- As forever, use past tense when you submit to your results, and put the whole thing in a reasonable order.
- Put figures and tables, appropriately numbered, in order at the end of the report
- If you desire, you may place your figures and tables properly within the text of your results part.

Figures and tables

- If you put figures and tables at the end of the details, make certain that they are visibly distinguished from any attach appendix materials, such as raw facts
- Despite of position, each figure must be numbered one after the other and complete with subtitle
- In spite of position, each table must be titled, numbered one after the other and complete with heading
- All figure and table must be adequately complete that it could situate on its own, divide from text

Discussion:

The Discussion is expected the trickiest segment to write and describe. A lot of papers submitted for journal are discarded based on problems with the Discussion. There is no head of state for how long a argument should be. Position your understanding of the outcome visibly to lead the reviewer through your conclusions, and then finish the paper with a summing up of the implication of the study. The purpose here is to offer an understanding of your results and hold up for all of your conclusions, using facts from your research and generally accepted information, if suitable. The implication of result should be visibly described. Infer your data in the conversation in suitable depth. This means that when you clarify an observable fact you must explain mechanisms that may account for the observation. If your results vary from your prospect, make clear why that may have happened. If your results agree, then explain the theory that the proof supported. It is never suitable to just state that the data approved with prospect, and let it drop at that.

- Make a decision if each premise is supported, discarded, or if you cannot make a conclusion with assurance. Do not just dismiss a study or part of a study as "uncertain."
- Research papers are not acknowledged if the work is imperfect. Draw what conclusions you can based upon the results that you have, and take care of the study as a finished work
- You may propose future guidelines, such as how the experiment might be personalized to accomplish a new idea.
- Give details all of your remarks as much as possible, focus on mechanisms.
- Make a decision if the tentative design sufficiently addressed the theory, and whether or not it was correctly restricted.
- Try to present substitute explanations if sensible alternatives be present.
- One research will not counter an overall question, so maintain the large picture in mind, where do you go next? The best studies unlock new avenues of study. What questions remain?
- Recommendations for detailed papers will offer supplementary suggestions.

Approach:

- When you refer to information, differentiate data generated by your own studies from available information
- Submit to work done by specific persons (including you) in past tense.
- Submit to generally acknowledged facts and main beliefs in present tense.

Administration Rules Listed Before Submitting Your Research Paper to Global Journals Inc. (US)

Please carefully note down following rules and regulation before submitting your Research Paper to Global Journals Inc. (US):

Segment Draft and Final Research Paper: You have to strictly follow the template of research paper. If it is not done your paper may get rejected.

- The **major constraint** is that you must independently make all content, tables, graphs, and facts that are offered in the paper. You must write each part of the paper wholly on your own. The Peer-reviewers need to identify your own perceptive of the concepts in your own terms. NEVER extract straight from any foundation, and never rephrase someone else's analysis.

- Do not give permission to anyone else to "PROOFREAD" your manuscript.

- Methods to avoid Plagiarism is applied by us on every paper, if found guilty, you will be blacklisted by all of our collaborated research groups, your institution will be informed for this and strict legal actions will be taken immediately.)
- To guard yourself and others from possible illegal use please do not permit anyone right to use to your paper and files.

Criterion for Grading a Research Paper (Compilation) by Global Journals Inc. (US)

Please note that following table is only a Grading of "Paper Compilation" and not on "Performed/Stated Research" whose grading solely depends on Individual Assigned Peer Reviewer and Editorial Board Member. These can be available only on request and after decision of Paper. This report will be the property of Global Journals Inc. (US).

Topics	Grades		
	A-B	**C-D**	**E-F**
Abstract	Clear and concise with appropriate content, Correct format. 200 words or below	Unclear summary and no specific data, Incorrect form Above 200 words	No specific data with ambiguous information Above 250 words
Introduction	Containing all background details with clear goal and appropriate details, flow specification, no grammar and spelling mistake, well organized sentence and paragraph, reference cited	Unclear and confusing data, appropriate format, grammar and spelling errors with unorganized matter	Out of place depth and content, hazy format
Methods and Procedures	Clear and to the point with well arranged paragraph, precision and accuracy of facts and figures, well organized subheads	Difficult to comprehend with embarrassed text, too much explanation but completed	Incorrect and unorganized structure with hazy meaning
Result	Well organized, Clear and specific, Correct units with precision, correct data, well structuring of paragraph, no grammar and spelling mistake	Complete and embarrassed text, difficult to comprehend	Irregular format with wrong facts and figures
Discussion	Well organized, meaningful specification, sound conclusion, logical and concise explanation, highly structured paragraph reference cited	Wordy, unclear conclusion, spurious	Conclusion is not cited, unorganized, difficult to comprehend
References	Complete and correct format, well organized	Beside the point, Incomplete	Wrong format and structuring

Index

A

Algebraic · 31

B

Bogoliubov · 31

C

Cabrezizo · 5, 6

E

Eigenvalue · 12, 16
Epimorphism · 90, 92, 94

G

Gronwall · 16, 20

H

Hamiltonian · 31, 39
Hildebrand · 26, 30

I

Isomorphism · 88, 90, 91

K

Kinematic · 42, 48

L

Laplacian · 7
Lorentzian · 1, 3

M

Monomorphism · 90, 92

N

Nilpotent · 88

O

Orthogonal · 6, 16, 18
Oscillator · 31

P

Precautionary · 60, 86

R

Renormalizable · 39

T

Theanalysis · 31
Twomomentum · 31

W

Wedderburn · 90

save our planet

Global Journal of Science Frontier Research

Visit us on the Web at www.GlobalJournals.org | www.JournalofScience.org
or email us at helpdesk@globaljournals.org

ISSN 9755896

© Global Journals